

Prevalence and Development of Psychiatric Disorders in Childhood and Adolescence

E. Jane Costello, PhD; Sarah Mustillo, PhD; Alaattin Erkanli, PhD; Gordon Keeler, MS; Adrian Angold, MRCPsych

Background: This longitudinal community study assessed the prevalence and development of psychiatric disorders from age 9 through 16 years and examined homotypic and heterotypic continuity.

Methods: A representative population sample of 1420 children aged 9 to 13 years at intake were assessed annually for *DSM-IV* disorders until age 16 years.

Results: Although 3-month prevalence of any disorder averaged 13.3% (95% confidence interval [CI], 11.7%-15.0%), during the study period 36.7% of participants (31% of girls and 42% of boys) had at least 1 psychiatric disorder. Some disorders (social anxiety, panic, depression, and substance abuse) increased in prevalence, whereas others, including separation anxiety disorder and attention-deficit/hyperactivity disorder (ADHD), decreased. Lagged analyses showed that children with a history of psychiatric disorder were 3 times more likely than those with no previous disorder to have a diagnosis

at any subsequent wave (odds ratio, 3.7; 95% CI, 2.9-4.9; $P < .001$). Risk from a previous diagnosis was high among both girls and boys, but it was significantly higher among girls. Continuity of the same disorder (homotypic) was significant for all disorders except specific phobias. Continuity from one diagnosis to another (heterotypic) was significant from depression to anxiety and anxiety to depression, from ADHD to oppositional defiant disorder, and from anxiety and conduct disorder to substance abuse. Almost all the heterotypic continuity was seen in girls.

Conclusions: The risk of having at least 1 psychiatric disorder by age 16 years is much higher than point estimates would suggest. Concurrent comorbidity and homotypic and heterotypic continuity are more marked in girls than in boys.

Arch Gen Psychiatry. 2003;60:837-844

From the Departments of Psychiatry and Behavioral Sciences (Drs Costello, Mustillo, and Angold and Mr Keeler) and Biostatistics and Bioinformatics (Dr Erkanli), Duke University Medical School, Durham, NC.

STUDIES THAT follow the same subjects as they grow up are the best source of information about the prevalence and causes of continuity and discontinuity of psychiatric disorders. A review¹ of the few studies that cover both child and adolescent psychiatric disorders²⁻⁹ showed that between 23% and 61% of children with a diagnosis at one wave had a diagnosis, although not necessarily the same one, at a subsequent wave. This suggests quite a high level of continuity. However, few studies have the power to distinguish between homotypic continuity (the same diagnosis at different assessments) and heterotypic continuity (continuity of disorder but a different diagnosis). Homotypic continuity is evidence for a disorder that has a similar manifestation across the age range of the study, whereas heterotypic continuity suggests an underlying vulnerability to psychiatric illness that may expose children to different disorders at different ages

or an underlying disorder that has different manifestations at different ages. The clinical and research implications of homotypic and heterotypic continuity are quite different.

In this article we address the following questions about continuity for the age range 9 to 16 years: What is the prevalence of *DSM-IV* disorders at different ages? (Because *DSM-IV* is inconsistent in whether impairment is required for a diagnosis, we include the prevalence of serious emotional disturbance (SED), the term used by the federal government¹⁰ for psychiatric disorder accompanied by significant impairment in the child's functioning.¹¹) Does the prevalence of these disorders increase or decrease as children grow up? What are the patterns of comorbidity? Is there continuity of disorder across this developmental period? Which disorders predict which other disorders? And are there significant sex differences in any of these estimates?

Table 1. Participant Age and Date of Interviews by Cohort

Cohort	Age, y	Wave, y							
		1, 1993	2, 1994	3, 1995	4, 1996	5, 1997	6, 1998	7, 1999	8, 2000
A	9	A1							
	10		A2						
B	11	B1		A3					
	12		B2		A4				
C	13	C1		B3					
	14		C2		B4		A5*		
	15			C3		B5		A6	
	16				C4		B6		A7
Response rate, %		94.2	90.5	88.0	78.0	80.3	75	75.2	80.7

*Financial constraints prevented us from interviewing any of cohort A during 1997 and the first half of 1998. Because subjects were randomly selected and interviewed around their birthdays, subjects interviewed during the second half of 1998 were a random sample of the whole cohort.

METHODS

The Great Smoky Mountains Study (GSMS) is a longitudinal study of the development of psychiatric disorder and the need for mental health services.¹²⁻¹⁷ Three cohorts of children, aged 9, 11, and 13 years at intake, were recruited from 11 counties in western North Carolina. A multistage sampling design was used, with weighting that returned the weighted sample to its original size.¹⁸ Potential participants were randomly selected from the population of some 20000 children using a household equal probability, accelerated cohort design.¹⁹ This means that each cohort reaches a given age in a different year, thus controlling for cohort effects.²⁰ The initial random sample of 4067 yielded 3896 screening questionnaires (95%) consisting mainly of the externalizing (behavioral) problems scale of the Child Behavior Checklist²¹ completed by a parent (usually the mother), by telephone or in person. All children scoring above a predetermined cutoff point (the top 25% of the total scores), plus a 1 in 10 random sample of the rest, were recruited for detailed interviews.

About 8% of area residents and the sample are African American, and fewer than 1% are Hispanic. American Indians make up only about 3% of the population of the study area, which is overwhelmingly white, but were oversampled from school records to constitute 25% of the study sample. This was done by using the same screening procedure but recruiting all American Indian children irrespective of screen score. Of the 456 American Indian children identified, screening questionnaires were obtained from 96%, and 81% (n=350) participated in the study. Although race was included in all analyses, no conclusions are drawn in this article about racial or ethnic similarities or differences, which are reported elsewhere.²²⁻²⁴

Table 1 presents the study design and the numbers of observations at each wave. Funding constraints prevented our interviewing the youngest cohort from January 1997 through June 1998. Because subjects were randomly selected, the cohort members interviewed between July and December 1998 are a random sample of the whole cohort. Data were collected on 1 cohort at ages 9 and 10 years, 2 cohorts at ages 11, 12, and 13 years, and all 3 cohorts at ages 14, 15, and 16 years. In the results presented in the current study, the data from the interviews of 9- and 10-year-olds are combined to provide a more reliable estimate.

MEASURES

The Child and Adolescent Psychiatric Assessment (CAPA) is an interviewer-based interview.^{14,25} The goal of interviews using this format, such as the adult Present State Examination²⁶

or Schedules for Clinical Assessment in Neuropsychiatry,²⁷ is to combine the advantages of clinical interviews with those of highly structured "epidemiologic" interview methods. A detailed glossary provides the operational rules for identifying clinically significant symptoms.

With the CAPA, parent and child are interviewed separately by different interviewers. In this article, with the exception of attention-deficit/hyperactivity disorder (ADHD) symptoms, about which only the parent was interviewed, we counted a symptom as present if it was reported by either the parent or the child or both, as is standard clinical practice.

The time frame of the CAPA for determining the presence of most psychiatric symptoms is the past 3 months. In the case of a few rare and severe acts, such as fire-setting or assault, a lifetime frame of reference is used, as required by DSM-IV. Two-week test-retest reliability of CAPA diagnoses in children aged 10 through 18 years is comparable to that of other highly structured child psychiatric interviews.^{16,25}

PROCEDURE

Interviewers were residents of the study area and had at least bachelor's-level degrees. They received 1 month of training, constant monitoring of quality control, and instruction by Department of Social Services staff in North Carolina's requirements for reporting abuse or neglect. All suspected cases were referred to the appropriate agency.

Interviews usually took place at home. The parent and child signed informed consent forms, and each was paid \$10.

DATA MANAGEMENT AND ANALYSIS

Scoring programs for the CAPA, written in SAS statistical software (SAS Institute Inc, Cary, NC), combined information about the date of onset, duration, and intensity of each symptom to create diagnoses according to the fourth edition of the DSM-IV.²⁸

GENERATING POPULATION PREVALENCE AND VARIANCE ESTIMATES

To generate population prevalence estimates from a multistage sampling design¹⁸ subjects were assigned a weight inversely proportional to their probability of selection. Correlation matrices were introduced to account for within-subject correlations. We used general estimation equations (GEEs)²⁹ to account for both the sampling design and within-subject correlation. In GEE, *subject* is introduced as a cluster (class) variable, and the sampling weights are introduced as a scale vec-

Table 2. Three-Month Prevalence of Selected DSM-IV Psychiatric Disorders*

Diagnosis	Total (N = 6674)	Age, y							Girls (n = 3005)	Boys (n = 3669)
		9-10 (n = 936)	11 (n = 901)	12 (n = 854)	13 (n = 833)	14 (n = 913)	15 (n = 1136)	16 (n = 1101)		
Any diagnosis†	13.3 (11.7-15.0)	19.5 (15.4-24.4)	12.7 (10.0-16.2)	8.3 (6.0-11.4)	12.7 (9.6-16.6)	9.7 (7.3-13.0)	14.2 (11.3-17.8)	12.7 (9.9-16.2)	10.6 (8.7-12.9)	15.8 (13.5-18.5)
Serious emotional disturbance‡	6.8 (5.8-7.9)	4.6 (3.4-6.3)	3.6 (2.7-4.9)	4.6 (3.1-6.7)	6.5 (4.5-9.3)	6.7 (4.8-9.3)	10.5 (8.0-13.6)	8.6 (6.5-11.5)	5.6 (4.3-7.3)	7.9 (6.5-9.5)
Any behavioral disorder	7.0 (6.0-8.2)	5.7 (4.2-7.8)	4.4 (3.1-6.1)	5.3 (3.5-7.9)	5.2 (3.6-7.5)	6.3 (4.5-8.8)	10.0 (7.6-13.0)	10.3 (7.8-13.5)	4.5 (3.5-5.7)	9.5 (7.9-11.5)
Conduct disorder	2.7 (2.1-3.5)	2.7 (1.8-4.1)	2.1 (1.3-3.4)	2.5 (1.3-4.8)	3.3 (2.0-5.4)	2.8 (1.7-4.8)	3.1 (1.9-5.0)	1.6 (0.7-3.3)	1.2 (0.7-2.1)	4.2 (3.1-5.6)
Oppositional defiant disorder	2.7 (2.2-3.3)	2.1 (1.5-2.9)	1.9 (1.4-2.6)	2.1 (1.2-3.5)	2.7 (1.6-4.6)	3.1 (2.0-4.7)	4.1 (2.7-6.2)	2.2 (1.3-3.9)	2.1 (1.6-2.9)	3.1 (2.4-4.2)
ADHD	0.9 (0.6-1.2)	2.2 (1.2-3.9)	1.4 (0.7-2.8)	1.4 (0.7-2.9)	0.5 (0.2-0.9)	0.3 (0.1-0.6)	0.4 (0.2-0.7)	0.3 (0.1-0.6)	0.3 (0.2-0.5)	1.5 (1.0-2.1)
SUDs	2.4 (1.8-3.1)	0	0	0.1 (<0.1-0.4)	0.3 (0.2-0.7)	1.4 (0.7-2.8)	5.3 (3.6-7.9)	7.6 (5.4-10.5)	2.0 (1.3-2.9)	2.8 (1.9-4.0)
Any anxiety disorder	2.4 (1.8-3.1)	4.6 (2.8-7.5)	2.6 (1.5-4.6)	0.9 (0.6-1.4)	2.0 (1.0-3.9)	1.8 (0.9-3.6)	2.8 (1.6-4.9)	1.6 (0.8-3.1)	2.9 (2.0-4.0)	2.0 (1.3-3.1)
Any depressive disorder	2.2 (1.6-3.0)	0.5 (0.3-1.0)	1.9 (1.0-3.5)	0.4 (0.2-0.8)	2.6 (1.3-5.3)	2.7 (1.4-5.0)	3.7 (2.3-6.0)	3.1 (1.8-5.3)	2.8 (1.8-4.3)	1.6 (1.0-2.5)

Abbreviations: ADHD, attention-deficit/hyperactivity disorder; SUDs, substance use disorders.

*Data are given as the percentage of participants (95% confidence intervals).

†One or more of separation anxiety disorder, generalized anxiety disorder, simple phobia, panic disorder, agoraphobia, major depression, dysthymia, depression not otherwise specified, bipolar disorder, ADHD, conduct disorder, oppositional disorder, anorexia nervosa, bulimia, mania, trichotillomania, enuresis, encopresis, SUDs, posttraumatic stress disorder, psychosis, or obsessive compulsive disorder.

‡Serious emotional disturbance is defined as any diagnosis accompanied by significant functional impairment.


tor that multiplies the subject's wave to wave correlation matrix. We also used the robust variance estimates (ie, sandwich type estimates), together with sampling weights, to adjust the standard errors of the parameter estimates to account for the multiphase sampling design. The use of multiwave data with the appropriate sample weights thus capitalized on the multiple observation points over time, while controlling for the effect on variance estimates of repeated measures. Identical results can be generated with SUDAAN³⁰ using primary sampling units as the subject identifiers, along with sampling weights. To obtain the prevalence of each of the various psychiatric disorders we fitted logistic regression models, again using GEE, with no predictors but including the intercept only. We used the Delta method (Taylor series expansion) to generate overall and time-averaged prevalence estimates and standard errors, corrected for within-subject correlations and design effects. Statistical tests such as comparison of prevalence between sexes used the same approach.

Lagged analyses were used to answer the questions about homotypic and heterotypic continuity. Two sets of analyses were done, one looking at the effect of a diagnosis in the year immediately preceding any annual interview and one looking at the effect of a diagnosis at any interview before the current one. Although the effects were stronger when the 1-year lag was used, results are more generalizable using the entire preceding period, and so the latter are reported here. Logistic regression analyses regressed this year's diagnosis on all past years, with other past and present psychiatric disorders entered as covariates.

RESULTS

THREE-MONTH PREVALENCE OF PSYCHIATRIC DISORDER

Table 2 presents the 3-month prevalence of a range of DSM-IV diagnoses by age and sex. A fuller version of the table can be found on our Web site (<http://www.devepi>



Three-month prevalence of any disorder and serious emotional disturbance (SED) by age and sex.

.mc.duke.edu; Web Table 1). In the case of depressive disorders, data are also presented on depressive disorder not otherwise specified, which includes the DSM-IV experimental category of minor depression. The overall prevalence of any disorder was highest in 9- to 10-year-olds, falling to its lowest level in 12-year-olds and then rising slowly (**Figure**). Twelve years was the age at which many of the disorders of childhood (ie, ADHD, separation anxiety disorder [SAD], enuresis and encopresis, and motor and verbal tic disorders) had almost disappeared, especially in boys, while those of adolescence and adulthood had not yet developed. Childhood disorders also decreased in girls across the same period but were less frequent to start with.

Table 3. Predicted Cumulative Prevalence of Psychiatric Disorders by Age 16 Years*

Diagnosis	Total	Girls	Boys
Any disorder	36.7 (2.7)	31.0 (2.3)	42.3 (3.1)
Any emotional disorder	15.0 (1.7)	17.1 (1.7)	13.0 (1.6)
Any anxiety disorder	9.9 (1.5)	12.1 (1.5)	7.7 (1.4)
Any depressive disorder	9.5 (1.1)	11.7 (1.2)	7.3 (1.0)
Any behavior disorder	23.0 (1.7)	16.1 (1.2)	29.9 (2.2)
Conduct disorder	9.0 (1.2)	3.8 (.7)	14.1 (1.8)
Oppositional defiant disorder	11.3 (1.0)	9.1 (1.0)	13.4 (1.0)
ADHD	4.1 (.7)	1.1 (.2)	7.0 (1.1)
SUDs	12.2 (.6)	10.1 (.5)	14.3 (.7)

Abbreviations: ADHD, attention-deficit/hyperactivity disorder; SUDs, substance use disorders.

*Data are given as the percentage of participants (SE).

The move into adolescence was marked by a rise in rates of depression and social phobia in girls that was not seen in boys. In middle adolescence the increase in substance use disorders (SUDs) in both sexes was dramatic, and there was also a modest increase in panic and generalized anxiety disorder (GAD).

The Figure also shows that SED increased as a proportion of psychiatric morbidity across adolescence, especially in boys. At age 9 and 10 years only 20% of boys and 31% of girls with a psychiatric disorder qualified as significantly impaired by SED. By age 16 years, 79% of diagnosed boys and 58% of diagnosed girls had SED.

ONSET OF DISORDERS BETWEEN AGES 9 AND 16 YEARS

We estimated the cumulative prevalence of disorders across the age range of the study; that is, the accumulation of new cases in previously unaffected children. (Note that the individual may have had an episode before entry into the study.) **Table 3** presents the predicted cumulative prevalence by age 16 years. By then, 36.7% of children had met *DSM-IV* criteria for 1 or more disorders. The estimate for boys greatly exceeded that for girls, as a result of a much higher cumulative prevalence of conduct disorder (CD) and ADHD. Girls accumulated more cases of depression and anxiety disorders.

CONCURRENT COMORBIDITY

Concurrent comorbidity refers to the co-occurrence of 2 or more diagnoses at the time of measurement,³¹ in this case, within the same 3-month period (A.A., A.E., E.J.C., and Helen Egger, MD, unpublished research, 2001). **Table 4** presents the major types of concurrent comorbidity by sex.

In bivariate analyses, 25 of 30 comparisons showed significant comorbidity. This fell to 16 of 30 after controlling for other comorbidities. There was significant comorbidity among the behavioral disorders and between anxiety and depression, as expected. The strong bivariate association between anxiety disorders and the behavioral disorders fell markedly and in most cases became nonsignificant when controlling for other forms of co-

morbidity. This was mainly because of the strong association between oppositional defiant disorder (ODD) and depression, on the one hand, and depression and anxiety, on the other.

There were 2 marked sex differences in patterns of concurrent comorbidity, both involving depression. After controlling for other comorbidity, depression was comorbid with CD in girls but not boys. Conversely, depression was comorbid with SUD in boys but not girls.

HOMOTYPIC AND HETEROTYPIC CONTINUITY

Children with a history of psychiatric disorder were 3 times more likely than those with no previous disorder to have a diagnosis at any subsequent wave of data collection (28.0% vs 9.3%; odds ratio [OR], 3.7; 95% confidence interval [CI], 2.9-4.9; $P < .001$). Risk from a previous diagnosis was high in girls (OR, 5.2; 95% CI, 3.4-7.9; $P < .001$) and boys (OR, 2.9; 95% CI, 2.0-4.1; $P < .001$), but it was significantly higher in girls (interaction OR, 1.8; 95% CI, 1.0-3.7; $P = .03$). Within this general picture, there were marked differences in continuity from one diagnosis to the same one (homotypic continuity) or to another (heterotypic continuity).

Looking first at homotypic continuity, this was significant for every diagnosis except for specific phobias. The details can be found on our Web site (<http://www.devepi.mc.duke.edu>; Web Table 2). The disorders showing the highest level of continuity were panic disorders, psychosis, verbal tics, enuresis and encopresis, and SUDs. Girls showed more continuity than boys, even though they had fewer psychiatric disorders. Prediction from past episodes was significantly higher in girls for depression, GAD, social phobia, and specific phobia. On the other hand, encopresis showed continuity only in boys. There were no significant sex differences in continuity for the behavioral disorders or SUDs, despite their greater frequency in boys.

We next tested whether there were cases in which the presence of one disorder earlier in life increased the risk of a different diagnosis later, net of the effects of concurrent comorbidity. Models included both homotypic and heterotypic continuity and then tested whether adding a term for concurrent comorbidity affected heterotypic continuity. Results are summarized in **Table 5**.

There was strong heterotypic continuity from depression to anxiety, which was not greatly affected by the high level of concurrent comorbidity between the 2 disorders. Prediction from anxiety to depression also occurred, and remained significant, although with a lower OR when controlling for concurrent comorbidity. Anxiety predicted later SUDs, but depression did not. Among the disruptive behavior disorders, ADHD showed a modest but significant prediction to later ODD, even controlling for the high level of concurrent comorbidity between the two. Conduct disorder predicted both ADHD and SUD in the bivariate model, but the very high level of concurrent comorbidity explained this.

Separate analyses by sex showed that heterotypic continuity was much more common in girls. Anxiety predicted depression, and vice versa, even controlling for concurrent comorbidity, but this was not the case for boys. Similarly, the link between past anxiety or CD and cur-

Table 4. Concurrent Comorbidity*†

Diagnosis	Any Anxiety Disorder	Any Depressive Disorder	ADHD	Conduct Disorder	Oppositional Defiant Disorder	SUDs
Any anxiety disorder		25.1 (9.9-63.3)	6.6 (2.9-15.4)	2.5 (1.1-5.5)‡	3.3 (1.7-6.5)	0.6 (0.2-1.9)
		27.9 (8.9-87.8) 	3.4 (1.2-9.8)‡	1.6 (0.7-3.9)	0.8 (0.5-2.5)	0.2 (0.04-0.8)‡
Any depressive disorder	28.4 (14.3-55.0)		8.0 (3.4-16.5)	5.0 (2.3-11.2)	20.7 (8.8-48.8)	9.9 (3.1-31.5)
	28.9 (13.8-60.7) 		2.2 (0.6-8.2)	0.7 (0.2-2.4)	16.7 (5.9-47.8) 	10.4 (2.7-40.3)
ADHD	7.7 (2.7-22.2)	3.3 (0.7-15.1)		3.7 (1.8-7.6)	8.7 (4.6-16.4)	NC
	6.0 (1.3-28.1)‡	0.1 (0.002-1.6)		1.9 (1.0-3.5)	6.6 (3.6-12.2) 	0.1 (0.01-1.1)
Conduct disorder	3.5 (1.3-5.9)	21.3 (9.0-50.5)	41.3 (14.0-121.9)		9.6 (5.3-17.3)	7.2 (3.1-16.7)
	0.3 (0.06-1.2)	10.6 (2.0-54.7)§	3.9 (1.1-13.6)		8.0 (4.5-14.0) 	5.7 (2.3-13.9)
Oppositional defiant disorder	7.7 (3.8-15.7)	15.1 (7.4-30.6)	79.0 (32.1-191.5)	44.1 (15.7-124.1)		4.1 (1.7-10.2)§
	2.4 (0.8-6.7)	7.1 (2.2-22.6) 	56.3 (10.6-153.4) 	29.4 (6.9-126.0) 		1.9 (0.7-4.6)
SUDs	0.4 (0.9-1.9)	3.5 (1.1-11.1)‡	0.3 (0.04-2.4)	23.4 (9.1-60.3)	2.4 (1.1-5.2)‡	
	NC	2.9 (0.6-12.9)	NC	30.7 (12.7-73.8) 	0.4 (0.04-2.7)	

Abbreviations: ADHD, attention-deficit/hyperactivity disorder; NC, model would not converge; SUDs, substance use disorders.

*Data are given as odds ratio (95% confidence interval). Data for girls are presented below the diagonal row of empty cells; boys, above the diagonal.

†Values in regular type are simple bivariate estimates, and values in boldface type are corrected for other possible comorbidities involving the 2 diagnoses under consideration (eg, anxiety and depression) controlling for comorbidity of depression with oppositional defiant disorder.

‡P<.05.

§P<.01.

||P<.001.

Table 5. Homotypic and Heterotypic Continuity With and Without Controls for Comorbidity*

Predicting to:	Past Depression	Past Anxiety	Past Conduct Disorder	Past ODD	Past ADHD	Past SUDs
Depression	7.0 (3.1-15.9)§	3.0 (1.7-5.4)§				
Controlling for comorbidity	4.2 (2.1-8.3)§	2.7 (1.8-5.2)‡				
Anxiety	5.7 (2.2-14.5)§	2.4 (1.6-3.7)§				
Controlling for comorbidity	2.8 (1.2-6.5)†	2.0 (1.2-3.4)†				
Conduct disorder			11.2 (5.9-21.1)§			
Controlling for comorbidity			10.3 (4.3-24.7)§			
ODD				3.7 (2.2-6.2)§	2.0 (1.1-3.8)†	
Controlling for comorbidity				4.7 (2.7-8.1)§	2.1 (1.1-4.2)†	
ADHD			2.2 (1.0-4.5)†		10.7 (5.2-22.3)§	
Controlling for comorbidity			1.8 (0.9-3.9)		9.6 (4.4-21.2)§	
SUDs		2.0 (1.2-3.5)†	2.7 (1.2-6.5)†			21.3 (6.3-72.5)§
Controlling for comorbidity		2.0 (1.1-3.7)†	1.7 (0.6-4.7)			25.7 (7.8-85.4)§

Abbreviations: ADHD, attention-deficit/hyperactivity disorder; ODD, oppositional defiant disorder; SUDs, substance use disorders.

*Data are given as the odds ratio (95% confidence interval). Empty cells indicate that continuity without controls for comorbidity was not significant. Boldface type indicates homotypic continuity.

†P<.05.

‡P<.01.

§P<.001.

rent SUDs (both of which persisted when controlled for concurrent comorbidity) were specific to girls. The link between past ADHD and current ODD was also specific to girls but appeared to be mainly the result of current comorbidity.

COMMENT

Although there has never been a representative population survey of child psychiatric disorders in the US comparable with the National Comorbidity Survey of 15- to 54-year-olds,³² or the recent British national survey of 5- to 15-year-olds,³³ researchers have patched together a picture of the prevalence of psychiatric disorders in American children of different ethnic groups and ages. This research has been reviewed several times,³⁴⁻³⁷ and a consensus has been reached that at any given time 1 child

in 5 will have a psychiatric disorder. This was also the conclusion that we reached on the basis of the first wave of data from GSMS, published in 1996.¹²

Analyses of further waves from the same data set, presented here, lead us to revise this picture. The 1-in-5 estimate was true of the youngest children, but as they grew up the 3-month prevalence fell to a low of 8.3% at age 12 years, before beginning to rise again in adolescence. This is consistent with the one other (cross-sectional) study that provides annual data by age group.³⁸

Earlier epidemiologic studies of children and adolescents in the 1980s and early 1990s tended to generate very varied, and sometimes very high, rates of some disorders, in particular anxiety disorders,^{6,39} ADHD,⁴⁰ and in some cases depression.^{32,41} However, recent studies have tended toward greater agreement and, on the whole, lower population estimates.^{33,38,42}

The GSMS sample comes from a small (11-county) and predominantly rural area of the southeastern United States. There were no Asian American subjects, and only 8% were African American. Thus, it does not represent the American population. A recent study of another part of North Carolina, a largely rural area including a large percentage of African American youth, found remarkably similar rates of disorder in white and African American children.³⁸ A recent British population study of youth aged 5 to 15 years produced quite similar results; for example, the prevalence of any anxiety disorder was 3.8% (GSMS, 2.4%); depression, 0.9% (GSMS, 2.2%); ADHD, 1.4% (GSMS, 0.9%); ODD, 2.9% (GSMS, 2.7%); and tic disorders, 0.1% (GSMS, 0.1%).

Despite the overlapping cohorts design, which used 3 age cohorts to permit age \times cohort comparisons, there is no escaping the fact that in the later data waves of GSMS the children's mean age is greater than it was in the earlier waves and the participants have been study subjects for longer. Thus, lower prevalence relative to cross-sectional studies may reflect (1) a tendency for subjects to report fewer symptoms in later interviews (ie, "symptom attenuation")⁴³⁻⁴⁵; (2) cohort differences, with the youngest cohort having more symptoms and the oldest cohort the fewest, controlling for age; and (3) differential dropout of children with and without psychiatric disorders. Another type of methodological challenge lies in the fact that the CAPA uses a 3-month window to estimate prevalence. If none of these methodological challenges is upheld, then we need to look for substantive reasons for the differences between this and earlier studies.

Symptom Attenuation

In a test-retest reliability study of 77 children aged 10 to 18 years using the CAPA,¹⁶ there was a very modest amount of attenuation between the first and second interviews; the difference was significant only for CD symptoms. The sample, however, had no 10-year-olds, so attenuation remains a possibility for the youngest children.

Cohort Differences

Analyses of symptom and diagnostic counts by cohort in GSMS (unpublished data) showed that the cohort differences were nonsignificant.

Differential Dropout

The response rates for the different waves of the study are reported in Table 1. Of 7944 possible interviews during the study period, 6675 (84%) were completed. Four subjects died, and 5.9% of participants only completed a single interview; these participants were no more or less likely than those who completed multiple interviews to have a diagnosis. Subjects who refused to be interviewed at one wave frequently returned later. This suggests that biased nonresponse is not a major reason for the observed prevalence rates.

Because the CAPA inquires only about the 3-month period preceding the annual interview, our estimate of cumulative prevalence may underestimate the number of cases occurring since the last interview, and thus the "burden of disease"⁴⁶ across childhood and adolescence.

There is no question that the CAPA underestimates cumulative prevalence. It misses cases that began and ended in the 9 months between interviews, including those for which some symptoms were present during the 3-month window, but not enough to reach the diagnostic threshold. Many psychiatric interviews adopt a 6-month,⁴⁷ 12-month,⁴⁸ or even lifetime⁴⁹ time frame for estimating the prevalence of psychiatric disorders. However, the evidence from studies using longer time frames suggests that a lot of forgetting occurs; for example, in the National Comorbidity Survey the lifetime rate for 15- to 54-year-olds (48.0%) was less than double the 12-month rate (29.5%).³² Clearly, both approaches underestimate the burden of psychiatric disorder. The CAPA uses a 3-month time frame because of the strong psychometric evidence that memory for the type of phenomena sought in psychiatric interviews is highly unreliable further back than 3 months.⁵⁰⁻⁵⁴ Ideally, study subjects would be interviewed every 3 months, but the burden imposed on subjects would introduce another set of biases.

Substantive Reasons

Controlling for wave, the 9- and 10-year-olds had significantly more symptoms and diagnoses than any other age. This was particularly true of the younger boys. As noted earlier, 9- and 10-year-old boys had very high rates of enuresis (9.5%), ADHD (3.6%), tic disorders (5.5%), and SAD (4.1%), all of them disorders of childhood that had fallen by more than 50% by age 11 years and disappeared by age 16 years. Excluding these diagnoses, the 3-month prevalence of psychiatric disorder for 9- and 10-year-old boys would fall to 8.3% (95% CI, 5.2%-12.9%).

In summary, bias is always a risk in survey research designed to estimate the population prevalence of a disorder.⁵⁵ We have guarded against it to some extent by choosing an interview that shows minimal attenuation effects, maintaining a good response rate over 8 years of data collection, and adopting appropriate statistical controls for the sampling design and the use of correlated data across waves. The result is an estimate of the 3-month prevalence of psychiatric disorder that is lower than our wave 1 estimate largely because of the fall in disorders of childhood: enuresis and encopresis, ADHD, tic disorders, and SAD.

PREVALENCE OVER TIME: THE BURDEN OF DISEASE

Multiple waves of data show that although only 13.3% of children, on average, had a diagnosis at any measurement point, almost 3 times this number had 1 or more disorders over the period of the study. This means that single-wave, cross-sectional studies are likely to under-

estimate the burden of disease⁴⁶ in the general population of children over time.

COMORBIDITY AND DEVELOPMENTAL CONTINUITY

As with other cross-sectional^{130,38,56} and longitudinal¹²⁻⁹ studies, we found considerable concurrent comorbidity: 25.5% of children with a diagnosis had 2 or more. It appears that much of the association between CD/ODD and ADHD is in fact carried by ODD rather than CD.³⁰ In both sexes, the high level of association between CD and ADHD fell noticeably when other diagnoses were included in the model, whereas the association between ADHD and ODD did not. Concurrent comorbidity between anxiety and depression was, as expected, strong in both sexes. Comorbidity between depression and ODD was also strong, but comorbidity between depression and CD was significant only for girls, when other types of comorbidity were controlled.

The data presented here on continuity of disorder across time suffer from the same shortcomings as do the cumulative prevalence figures: we have only a series of snapshots on which to base our estimates. Given that, the degree of homotypic continuity is remarkable. In bivariate analyses, every *DSM-IV* diagnosis examined showed significant homotypic continuity with the exception of specific phobias. Apart from encopresis, which rarely occurred in girls, continuity was higher in girls than in boys.

Compared with the level of homotypic continuity, there were few cases of heterotypic continuity. Heterotypic continuity was much stronger in girls than in boys. This suggests that the *DSM-IV* taxonomy may fit boys' developmental patterns better than those of girls. There is no evidence that boys with an emotional disorder were at increased risk of a behavioral disorder, or vice versa, whereas girls with anxiety disorders had increased risk for later SUDs. (See Kaplow et al⁵⁷ for a more detailed examination of this issue.) Robins and Price's prediction⁵⁸ that later SUDs will be predicted by CDs holds here for girls but not for boys. This sample is too young to test the prediction of Zoccolillo et al⁵⁹ that CD in girls is likely to lead to later depression and somatization disorders.

An important issue that exceeded the scope of this article is developmental continuity among the anxiety disorders and between specific anxiety disorders and other diagnoses. Continuity was much higher for some anxiety disorders (panic disorder and posttraumatic stress disorder) than for others (specific phobia and GAD) (<http://www.devepi.mc.duke.edu>; Web Table 2). Much more work needs to be done on comorbidity and heterotypic continuity among these disorders and with depression.

In summary, data on a representative population of children and adolescents growing up in the 1990s show that at any time 1 in 6 will have a psychiatric disorder and at least 1 in 3 will have 1 or more psychiatric disorders by age 16 years. As children grow older, psychiatric disorders are more and more likely to be accompanied by significant functional impairment. Once children, particularly girls, develop a psychiatric disorder their

chances of continuing to have one, or of developing another episode after remission, are much higher than those of their unaffected peers. By mid-adolescence, although some disorders of childhood have disappeared, impairing adult disorders such as depression, panic disorder, and SUDs are becoming the most prevalent problems. Much more work on the childhood antecedents of these disorders is needed if prevention programs are to be effective.

Submitted for publication September 4, 2002; final revision received January 13, 2003; accepted January 22, 2003.

This study was supported by grants DA11301 and MH57761 and Independent Scientist Award MH01167 from the National Institutes of Health, Bethesda, Md, and Faculty Scholar awards from the William T. Grant Foundation, New York, NY (Drs Costello and Angold).

Corresponding author and reprints: E. Jane Costello, PhD, Center for Developmental Epidemiology, Department of Psychiatry and Behavioral Sciences, Duke University Medical School, DUMC Box 3454, Durham, NC 27710 (e-mail: jcostell@psych.mc.duke.edu).

REFERENCES

1. Costello EJ, Angold A. Developmental epidemiology. In: Cicchetti D, Cohen D, eds. *Developmental Psychopathology*. Vol 1. New York, NY: John Wiley & Sons; 1995:23-56.
2. Kashani JH, McGee RO, Clarkson SE, Anderson JC, Walton LA, Williams S, Silva PA, Robins AJ, Cytryn L, McKnew DH. Depression in a sample of 9-year-old children: prevalence and associated characteristics. *Arch Gen Psychiatry*. 1983;40:1217-1223.
3. Anderson JC, Williams S, McGee R, Silva PA. *DSM-III* disorders in preadolescent children: prevalence in a large sample from the general population. *Arch Gen Psychiatry*. 1987;44:69-77.
4. McGee R, Feehan M, Williams S, Partridge F, Silva PA, Kelly J. *DSM-III* disorders in a large sample of adolescents. *J Am Acad Child Adolesc Psychiatry*. 1990;29:611-619.
5. Fergusson DM, Horwood LJ, Lynskey MT. Prevalence and comorbidity of *DSM-III-R* diagnoses in a birth cohort of 15 year olds. *J Am Acad Child Adolesc Psychiatry*. 1993;32:1127-1134.
6. Velez CN, Johnson J, Cohen P. A longitudinal analysis of selected risk factors of childhood psychopathology. *J Am Acad Child Adolesc Psychiatry*. 1989;28:861-864.
7. Cohen P, Cohen J, Kasen S, Velez CN, Hartmark C, Johnson J, Rojas M, Brook J, Streuning EL. An epidemiological study of disorders in late childhood and adolescence, 1: age- and gender-specific prevalence. *J Child Psychol Psychiatry*. 1993;34:851-867.
8. Reinherz HZ, Giaconia RM, Lefkowitz ES, Pakiz B, Frost AK. Prevalence of psychiatric disorders in a community population of older adolescents. *J Am Acad Child Adolesc Psychiatry*. 1993;32:369-377.
9. Lewinsohn PM, Rohde P, Klein DN, Seeley JR. Natural course of adolescent major depressive disorder, I: continuity into young adulthood. *J Am Acad Child Adolesc Psychiatry*. 1999;38:56-63.
10. 58 *Federal Register* 29425 (1993).
11. Canino G, Costello EJ, Angold A. Assessing functional impairment and social adaptation for child mental health services research: a review of measures. *J Ment Health Serv*. 1999;1:93-108.
12. Costello EJ, Angold A, Burns BJ, Stangl DK, Tweed DL, Erkanli A, Worthman CM. The Great Smoky Mountains Study of Youth: goals, designs, methods, and the prevalence of *DSM-III-R* disorders. *Arch Gen Psychiatry*. 1996;53:1129-1136.
13. Burns BJ, Costello EJ, Angold A, Tweed D, Stangl D, Farmer EMZ, Erkanli A. Children's mental health service use across service sectors. *Health Aff (Millwood)*. 1995;14:147-159.
14. Angold A, Prendergast M, Cox A, Harrington R, Simonoff E, Rutter M. The Child and Adolescent Psychiatric Assessment (CAPA). *Psychol Med*. 1995;25:739-753.
15. Costello EJ, Angold A, March J, Fairbank J. Life events and post-traumatic stress: the development of a new measure for children and adolescents. *Psychol Med*. 1998;28:1275-1288.

16. Angold A, Costello EJ. A test-retest reliability study of child-reported psychiatric symptoms and diagnoses using the Child and Adolescent Psychiatric Assessment (CAPA-C). *Psychol Med*. 1995;25:755-762.
17. Angold A, Erkanli A, Costello EJ, Rutter M. Precision, reliability and accuracy in the dating of symptom onsets in child and adolescent psychopathology. *J Child Psychol Psychiatry*. 1996;37:657-664.
18. Shrout PE, Newman SC. Design of two phase prevalence surveys of rare disorders. *Biometrics*. 1989;45:549-555.
19. Schaie KW. A general model for the study of developmental problems. *Psychol Bull*. 1965;64:92-107.
20. Kleinbaum DG, Kupper LL, Morgenstern H. *Epidemiologic Research: Principles and Quantitative Methods*. New York, NY: Van Nostrand Reinhold; 1982.
21. Achenbach TM, Edelbrock CS. Behavioral problems and competencies reported by parents of normal and disturbed children aged four through sixteen. *Monogr Soc Res Child Dev*. 1981;46:1-82.
22. Costello E, Farmer E, Angold A, Burns B, Erkanli A. Psychiatric disorders among American Indian and White youth in Appalachia: the Great Smoky Mountains Study. *Am J Public Health*. 1997;87:827-832.
23. Federman EB, Costello EJ, Angold A, Farmer EMZ, Erkanli A. Development of substance use and psychiatric comorbidity in an epidemiologic study of white and American Indian young adolescents: the Great Smoky Mountains Study. *Drug Alcohol Depend*. 1997;44:69-78.
24. Costello EJ, Farmer EMZ, Angold A. Same place, different children: White and American Indian children in the Appalachian Mountains. In: Cohen P, Robins L, Slomkowski C, eds. *Where and When: Historical and Geographical Aspects of Psychopathology*. Mahwah, NJ: Lawrence Erlbaum Associates; 1999:279-298.
25. Angold A, Fisher PW. Interviewer-based interviews. In: Shaffer D, Lucas C, Richters J, eds. *Diagnostic Assessment in Child and Adolescent Psychopathology*. New York, NY: Guilford Press; 1999:34-64.
26. Wing JK, Cooper JE, Sartorius N. Present State Examination (PSE). In: Wing JK, Cooper JE, Sartorius N, eds. *The Measurement and Classification of Psychiatric Symptoms*. Cambridge, England: Cambridge University Press; 1974:1-9.
27. Wing JK, Babor T, Brugha T, Burke J, Cooper JE, Giel R, Jablenski A, Regier D, Sartorius N. SCAN: Schedules for Clinical Assessment in Neuropsychiatry. *Arch Gen Psychiatry*. 1990;47:589-593.
28. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition*. Washington, DC: American Psychiatric Association; 1994.
29. Zeger SL, Liang KY, Albert PS. Models for longitudinal data: a generalized estimating equation approach. *Biometrics*. 1988;44:1049-1060.
30. Shah BV, Folsom RE, LaVange LM, Wheelless SC, Boyle KE, Williams RL. *SUDAAN Software for the Statistical Analysis of Correlated Data: Statistical Methods and Mathematical Algorithms Used in SUDAAN, 1995*. Research Triangle Park, NC: RTI International;
31. Angold A, Costello EJ, Erkanli A. Comorbidity. *J Child Psychol Psychiatry*. 1999; 40:57-87.
32. Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen HU, Kendler KS. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: results from the National Comorbidity Survey. *Arch Gen Psychiatry*. 1994;51:8-19.
33. Meltzer H, Gatward R, Goodman R, Ford T. *The Mental Health of Children and Adolescents in Great Britain*. London, England: Office for National Statistics; 1999.
34. Costello EJ. Developments in child psychiatric epidemiology. *J Am Acad Child Adolesc Psychiatry*. 1989;28:836-841.
35. Costello EJ, Angold A. Epidemiology. In: March J, ed. *Anxiety Disorders in Children and Adolescents*. New York, NY: Guilford Press; 1995:109-124.
36. Costello EJ, Angold A. Developmental epidemiology: a framework for developing psychopathology. In: Lewis M, Sameroff A, eds. *Handbook of Developmental Psychopathology*. 2nd ed. New York, NY: Kluwer Academic/Plenum Press; 2000:57-73.
37. Brandenburg NA, Friedman RM, Silver SE. The epidemiology of childhood psychiatric disorders. *J Am Acad Child Adolesc Psychiatry*. 1990;29:76-83.
38. Angold A, Erkanli A, Farmer EMZ, Fairbank JA, Burns BJ, Keeler G, Costello EJ. Psychiatric disorder, impairment, and service use in rural African American and White youth. *Arch Gen Psychiatry*. 2002;59:893-901.
39. Costello EJ, Costello AJ, Edelbrock C, Burns BJ, Dulcan MK, Brent D, Janiszewski S. Psychiatric disorders in pediatric primary care: prevalence and risk factors. *Arch Gen Psychiatry*. 1988;45:1107-1116.
40. Bird HR, Canino G, Rubio-Stipec M, Gould MS, Ribera J, Sesman M, Woodbury M, Huertas-Goldman S, Pagan A, Sanchez-Lacay A, Moscoso M. Estimates of the prevalence of childhood maladjustment in a community survey in Puerto Rico: the use of combined measures. *Arch Gen Psychiatry*. 1988;45:1120-1126.
41. Lewinsohn PM, Hops H, Roberts RE, Seeley JR, Andrews JA. Adolescent psychopathology, I: prevalence and incidence of depression and other DSM-III-R disorders in high school students. *J Abnorm Psychol*. 1993;102:133-144.
42. Shaffer D, Fisher PW, Dulcan M, Davies M, Piacentini J, Schwab-Stone M, Lahey BB, Bourdon K, Jensen P, Bird H, Canino G, Regier D. The NIMH diagnostic interview schedule for children (DISC 2.3): description, acceptability, prevalences, and performance in the MECA study. *J Am Acad Child Adolesc Psychiatry*. 1996;35:865-877.
43. Jensen PS, Shaffer D, Rae D, Canino G, Bird HR, Dulcan MK, Lahey BB, Rubio-Stipec M, Goodman S, Richters JE. Attenuation of the Diagnostic Interview Schedule for Children (DISC 2.1): sex, age and IQ relationships. Paper presented at: 39th Annual Meeting of the American Academy of Child and Adolescent Psychiatry; October 1992; Washington, DC.
44. Piacentini J, Jensen P, Shaffer D, Bird HR, Canino G, Dulcan M, Richters J, Roper M, Fisher PW, Davies M, Chen J, Goodman C, Rubio M, Bourdon K. Attenuation of child psychiatric diagnostic information: magnitude and relationship to severity. Paper presented at: Annual Meeting of the Society for Research in Child and Adolescent Psychopathology; July 1992; San Francisco, Calif.
45. Piacentini J, Roper M, Jensen P, Lucas C, Fisher P, Bird H, Bourdon K, Schwab-Stone M, Rubio-Stipec M, Davies M, Dulcan M. Informant-based determinants of symptom attenuation in structured child psychiatric interviews. *J Abnorm Child Psychol*. 1999;27:417-428.
46. Murray CJL, Lopez AD. *The Global Burden of Disease*. Geneva, Switzerland: World Health Organization; 1996.
47. Shaffer D, Fisher PW, Schwab-Stone M, Piacentini J, Wicks J. *The Diagnostic Interview Schedule for Children (DISC 2.1)*. Rockville, Md: National Institute of Mental Health; 1989.
48. Shaffer D, Fisher P, Dulcan MK, Davies M. The NIMH Diagnostic Interview Schedule for Children, Version 2.3 (DISC 2.3): description, acceptability, prevalence rates, and performance in the MECA study. *J Am Acad Child Adolesc Psychiatry*. 1996;35:865-877.
49. Lewinsohn P, Zinbarg J, Lewinsohn M, Sack W. Lifetime comorbidity among anxiety disorders and between anxiety disorders and other mental disorders in adolescents. *J Anxiety Disord*. 1997;11:377-394.
50. Som RK. *Recall Lapse in Demographic Enquiries*. New York, NY: Asia Publishing House; 1973.
51. Bjorklund DF, Muir JE. Children's development of free recall memory: remembering on their own. *Ann Child Dev*. 1988;5:79-123.
52. Harlow SD, Linet MS. Agreement between questionnaire data and medical records: the evidence for accuracy of recall. *Am J Epidemiol*. 1989;129:233-248.
53. Pillemer DB, White SH. Childhood events recalled by children and adults. *Adv Child Dev Behav*. 1989;21:297-340.
54. Angold A, Costello EJ. Toward establishing an empirical basis for the diagnosis of oppositional defiant disorder. *J Am Acad Child Adolesc Psychiatry*. 1996;35: 1205-1212.
55. Rothman KJ, Greenland S. *Modern Epidemiology*. Philadelphia, Pa: Lippincott-Raven; 1998.
56. Canino GJ, Bird HR, Shrout PE, Rubio-Stipec M, Bravo M, Martinez R, Sesman M, Guevara LM. The prevalence of specific psychiatric disorders in Puerto Rico. *Arch Gen Psychiatry*. 1987;44:727-735.
57. Kaplow JB, Curran PJ, Angold A, Costello EJ. The prospective relation between dimensions of anxiety and the initiation of adolescent alcohol use. *J Clin Child Psychol*. 2001;30:316-326.
58. Robins LN, Price RK. Adult disorders predicted by childhood conduct problems: results from the NIMH Epidemiologic Catchment Area Project. *Psychiatry*. 1991;54:116-132.
59. Zoccolillo M, Pickles A, Quinton D, Rutter M. The outcome of childhood conduct disorder: implications for defining adult personality disorder and conduct disorder. *Psychol Med*. 1992;22:971-986.