

Measurement of Phosphorylated Tau Epitopes in the Differential Diagnosis of Alzheimer Disease

A Comparative Cerebrospinal Fluid Study

Harald Hampel, MD; Katharina Buerger, MD; Raymond Zinkowski, PhD; Stefan J. Teipel, MD; Alexander Goernitz, MD; Niels Andreasen, MD, PhD; Magnus Sjoegren, MD; John DeBernardis, PhD; Daniel Kerkman, PhD; Koichi Ishiguro, PhD; Hideto Ohno, PhD; Eugene Vanmechelen, PhD; Hugo Vanderstichele, PhD; Cheryl McCulloch, BS; Hans-Jürgen Möller, MD; Peter Davies, PhD; Kaj Blennow, MD, PhD

Background: Abnormal hyperphosphorylation of the microtubule-associated protein tau and its incorporation into neurofibrillary tangles are major hallmarks of the pathogenesis of Alzheimer disease (AD). Different tau phosphoepitopes can be sensitively detected in cerebrospinal fluid (CSF).

Objective: To compare the diagnostic accuracy of CSF concentrations of tau proteins phosphorylated at 3 pathophysiologically important epitopes (p-tau) to discriminate among patients with AD, nondemented control subjects, and patients with other dementias.

Design and Setting: Cross-sectional, bicenter, memory clinic-based studies.

Participants: One hundred sixty-one patients with a clinical diagnosis of AD, frontotemporal dementia, dementia with Lewy bodies, or vascular dementia and 45 nondemented controls (N=206).

Main Outcome Measures: Levels of tau protein phosphorylated at threonine 231 (p-tau₂₃₁), threonine 181 (p-tau₁₈₁), and serine 199 (p-tau₁₉₉). The CSF p-tau pro-

tein levels were measured using 3 different enzyme-linked immunosorbent assays.

Results: The mean CSF levels of the studied p-tau proteins were significantly elevated in patients with AD compared with the other groups. Applied as single markers, p-tau₂₃₁ and p-tau₁₈₁ reached specificity levels greater than 75% between AD and the combined non-AD group when sensitivity was set at 85% or greater. Statistical differences between the assay performances are presented. Particularly, discrimination between AD and dementia with Lewy bodies was maximized using p-tau₁₈₁ at a sensitivity of 94% and a specificity of 64%, and p-tau₂₃₁ maximized group separation between AD and frontotemporal dementia with a sensitivity of 88% and a specificity of 92%. Combinations of the 3 markers did not add discriminative power compared with the application as single markers.

Conclusions: The p-tau proteins in CSF come closest to fulfilling the criteria of a biological marker of AD. There is a tendency for p-tau proteins to perform differently in the discrimination of primary dementia disorders from AD.

Arch Gen Psychiatry. 2004;61:95-102

ABNORMAL HYPERPHOSPHORYLATION of the microtubule-associated protein tau and its incorporation into neurofibrillary tangles are major components of the pathogenesis of Alzheimer disease (AD).

Using monoclonal antibodies specific for different phosphorylated epitopes of tau, enzyme-linked immunosorbent assays have been developed that sensitively measure concentrations of phosphorylated tau protein (p-tau) in cerebrospinal fluid (CSF). Statistically significant increases in CSF p-tau concentrations in patients with AD have recently been demonstrated in independent pilot studies, mainly using 3 different immunoassays specific for the phosphorylated epitopes threonine 231 (p-tau₂₃₁),^{1,2} threonine 181 (p-tau₁₈₁),^{3,4} and serine 199 (p-tau₁₉₉).⁵

Evidence from these pilot studies indicates that quantification of tau phosphorylated at these specific sites may improve early detection, differential diagnosis, and tracking of disease progression in AD (for a review see Blennow et al^{6,7}).

In a recent study,¹ CSF p-tau₂₃₁ distinguished between patients with AD and those with other neurologic disorders (ONDs) with a sensitivity of 85% and a specificity of 97%. Furthermore, p-tau₂₃₁ significantly improved differential diagnosis between AD and other non-AD groups, particularly frontotemporal dementia (FTD).⁸ In AD vs FTD, p-tau₂₃₁ correctly allocated 91% of patients vs only 66% using total tau.⁸ Itoh and colleagues⁵ showed that CSF p-tau₁₉₉ discriminates between AD and the combined non-AD groups with a sensitivity and a specificity of 85%. The level of CSF p-tau₁₈₁ was elevated in pa-

Author affiliations are listed at the end of this article.

Table 1. Characteristics of Patients and Controls and CSF p-Tau Values

Group	Age, Mean ± SD (Range), y	Sex, F/M, No.	MMSE Score, Mean ± SD (Range)	CSF Level, Median (25 th -75 th Percentile)		
				p-Tau ₂₃₁ , pg/mL	p-Tau ₁₈₁ , pmol/L	p-Tau ₁₉₉ , fmol/mL
AD (n = 108)	72.6 ± 5.7 (54-84)	65/43	22.2 ± 5.4 (5-30)	667.5 (361.0-907.2)	20.5 (16.5-26.2)	1.7 (1.3-2.2)
DLB + FTD + VaD (n = 53)	71.3 ± 8.8 (46-87)	26/27	20.5 ± 5.0 (10-30)	140.0 (50.5-285.5)*	11.3 (9.4-15.7)*	1.1 (0.9-1.5)*
DLB (n = 22)	76.1 ± 6.1† (65-87)	7/15	21.2 ± 5.4 (10-29)	213.5 (92.8-374.5)*	11.3 (9.0-16.9)*	1.1 (0.8-1.4)*
FTD (n = 24)	66.0 ± 0.9* (46-79)	15/9	19.8 ± 4.8† (13-30)	86.5 (9.0-174.5)*	10.7 (9.3-14.2)*	1.1 (0.9-1.6)*
VaD (n = 7)	74.1 ± 4.3 (65-77)	4/3	21.3 ± 4.9 (12-25)	201.0 (44.0-612.0)†	13.6 (12.1-19.0)†	1.3 (0.9-1.9)
OND (n = 22)	61.1 ± 9.0* (49-81)	16/6	27.8 ± 3.2* (17-30)	54.0 (13.3-97.0)*	11.3 (8.6-12.6)*	0.8 (0.6-1.0)*
Control (n = 23)	60.0 ± 10.1* (44-77)	14/9	29.2 ± 0.7* (27-30)	35.0 (18.0-70.0)*	11.0 (9.0-13.4)*	0.8 (0.6-1.1)*

Abbreviations: AD, Alzheimer disease; CSF, cerebrospinal fluid; DLB, dementia with Lewy bodies; FTD, frontotemporal dementia; MMSE, Mini-Mental State Examination; OND, other neurologic disorder; p-tau, phosphorylated tau protein; VaD, vascular dementia.

**P* < .001, AD group vs the other groups studied.

†*P* < .05, AD group vs the other groups studied.

tients with AD compared with patients with other dementias and controls,⁹ and p-tau₁₈₁ has been proposed as a potential marker for discriminating patients with AD from those with dementia with Lewy bodies (DLB).¹⁰ Furthermore, CSF p-tau₂₃₁ concentrations declined over time during the clinical progression of AD and correlated with cognitive performance at baseline.¹¹ Results of these first studies suggested that CSF p-tau proteins are promising biomarker candidates for AD.

Levels of CSF p-tau and the discriminative power of the 3 different p-tau assays among patients with AD, nondemented controls, and patients with other dementias, however, have not yet been assessed in the same set of patients. Investigation of the 3 p-tau assays would allow for direct comparison of the diagnostic performance of the 3 markers to differentiate AD from other relevant diseases. In the advent of large international multicenter trials (such as the National Institute on Aging Initiative on Neuroimaging in Alzheimer's Disease and the studies of the European Alzheimer's Disease Consortium) on the value of biological markers and neuroimaging in the diagnosis of AD, this is the first study to address the important clinical issue regarding the differential diagnostic performances of p-tau proteins as core biological marker candidates.

In particular, we tested the diagnostic accuracy of the 3 p-tau assays according to the recommendations of a consensus group¹² suggesting a sensitivity level of 85% or greater and a specificity level of at least 75% for useful biomarkers of AD. In addition, we asked whether use of a combination of the 3 markers might be superior to the application of single markers. To our knowledge, this is the first comparative study applying 3 developed and published immunoassays detecting different p-tau epitopes on the same set of controls and patients to compare individual and combined diagnostic accuracy. Concentrations of p-tau₂₃₁, p-tau₁₈₁, and p-tau₁₉₉ in the CSF were studied in the same group of patients with AD, DLB, FTD, or vascular dementia (VaD); patients with ONDs; and controls.

METHODS

PATIENT SELECTION

A total of 206 individuals were studied. One hundred eight patients had probable AD (National Institute of Neurological and

Communicative Disorders and Stroke–Alzheimer's Disease and Related Disorders Association criteria)¹³ and 53 had other dementia disorders (24 patients with FTD,¹⁴ 22 with DLB,¹⁵ and 7 with VaD).¹⁶ Structural and functional imaging results were consistent with the diagnoses in these patients. Twenty-two patients with ONDs were diagnosed as having mild psychiatric (eg, depressed mood) or neurologic (eg, dizziness) symptoms. One patient with OND experienced a bulbar syndrome of unknown etiology. We also studied 23 controls. Study participants were recruited at 2 academic expert centers: the Dementia Research Section and Memory Clinic, Alzheimer Memorial Center and Geriatric Psychiatry Branch, Department of Psychiatry, Ludwig-Maximilian University (22 patients with AD, 6 with FTD, 7 with VaD, 9 with DLB, 1 with OND, and 13 controls), and the Department of Clinical Neuroscience, University of Göteborg (86 patients with AD, 18 with FTD, 13 with DLB, 21 with OND, and 10 controls). Participants from the former center had been studied previously with a different objective, and the results of this study have been published.⁸ Characteristics of the patients and controls are given in **Table 1**. The protocol was approved by the local ethical committees and the institutional review boards of the 2 participating medical centers. Informed consent was obtained from all participants.

Examination of the controls included medical history, physical examination, routine blood tests (blood cell count; international normalized ratio; partial thromboplastin time; and sodium, potassium, creatinine, urea, and blood glucose levels), and a cognitive test using the Consortium to Establish a Registry for Alzheimer's Disease battery.¹⁷ Ten of the 23 controls were volunteers without any medical, neurologic, or psychiatric disorders. Samples of CSF were collected from 13 controls while they underwent spinal anesthesia for surgery of the urinary tract or lower extremities. They were cognitively normal according to the Consortium to Establish a Registry for Alzheimer's Disease battery (results within ±1 SD in all subtests). Three of these control subjects had diabetes mellitus as a somatic comorbidity.

CSF SAMPLING AND ANALYSES OF p-TAU PROTEINS

Samples of CSF were acquired via lumbar puncture between 9 and 11 AM according to a routine protocol (established by the 2 participating medical centers). Samples of CSF were collected in polypropylene tubes on ice in 0.5-mL aliquots. For this study, a total of 1 mL was taken. Aliquots were centrifuged at 4°C at 10000g for 10 minutes and stored at -80°C until analysis. The same procedures were performed at the 2 sites involved in the study. There was no effect of medical centers

on the variance of measured protein levels and no protein gradient in the CSF column for the markers (K.B., unpublished data, 2002).

Assay operators were masked to the diagnostic category of the samples. Levels of p-tau₂₃₁ were measured using an enzyme-linked immunosorbent assay (Applied NeuroSolutions Inc).¹ This assay uses a combination of CP27 (which recognizes amino acids 130-150 in normal tau and p-tau), Tau-1 (which recognizes amino acids 196-205 in nonphosphorylated tau), and CP9 (which recognizes phosphothreonine 231). Experiments describing the specificity of the detection antibody CP9 for phosphothreonine 231 have been reported previously.¹ Full-length recombinant tau (441 amino acids) phosphorylated at threonine 231 was used to produce a standard curve. Levels of p-tau₂₃₁ in our patients were calculated from the standard curve and expressed as CSF p-tau₂₃₁ in picograms per milliliter.

Levels of p-tau₁₈₁ were measured using a sandwich enzyme-linked immunosorbent assay method (prototype version of Innostest Phospho-Tau [181p]; Innogenetics), using a combination of monoclonal antibody HT7 (which recognizes amino acids 159-163 in normal tau and p-tau) and biotinylated monoclonal antibody AT270 (which recognizes p-tau containing the phosphorylated threonine 181 residue).³ A synthetic phosphopeptide was used for standardization.

Levels of p-tau₁₉₉ were measured using a previously reported sandwich enzyme-linked immunosorbent assay method (Mitsubishi Chemical Corp, Shinagawa, Japan), using a combination of monoclonal antibody HT7 and polyclonal antibody anti-PS199 (specific for tau phosphorylated at serine 199).⁵

STATISTICAL ANALYSES

Differences among groups regarding age were assessed using the Mann-Whitney (M-W) test and regarding sex distribution using the χ^2 test.

Distributions of p-tau values differed statistically significantly from normal as revealed by the Kolmogorov-Smirnov test. Differences in mean CSF levels of the 3 p-tau subtypes among all groups were assessed using the Kruskal-Wallis test. Pairwise comparisons between patients with AD and the other groups were performed using the M-W test. Correlations between p-tau subtypes were assessed using the Spearman rank correlation.

Cutoff values for p-tau proteins were determined such that 85% of the patients with AD were correctly identified according to the recommendations of the consensus conference¹² of 85% sensitivity for an "excellent" biomarker. To determine differences in diagnostic accuracy among markers, the specificity levels that correspond to the 85% sensitivity cutoff level for each marker in the comparison groups were compared between all possible pairs of markers using the McNemar test.

To develop predictive cutoff values that optimized the combined use of the different p-tau markers, we used classification tree analysis with SYSTAT 7.0 (SPSS Inc, Chicago, Ill). Classification tree analysis uses recursive partitioning to consider all possible binary splits of the data in pursuit of optimal classification.¹⁸ The analysis considered all 3 p-tau markers simultaneously to create cutoff values that maximize separation among groups, resulting in a decision tree. The number of branches depends on the separation that has been achieved at the first split. To avoid overfitting of the data, generation of further branches was interrupted if sensitivity or specificity declined below 80%. A similar technique has been presented in earlier CSF studies in AD.¹⁹

To show sensitivity and specificity levels over the entire range of cutoff levels, we determined receiver operating characteristic curves.




Figure 1. Levels of cerebrospinal fluid (CSF) phosphorylated tau protein (p-tau)₂₃₁ (A), p-tau₁₈₁ (B), and p-tau₁₉₉ (C) in patients and controls. Dashed lines represent the cutoff level when sensitivity was set at 85% or higher. Asterisk indicates differences from AD at $P < .001$; dagger, differences from AD at $P < .05$; AD, Alzheimer disease; ONDs, other neurologic disorders; DLB, dementia with Lewy bodies; FTD, frontotemporal dementia; and VaD, vascular dementia.

RESULTS

CSF p-TAU LEVELS

As illustrated in **Figure 1** and Table 1, levels of all p-tau subtypes were significantly increased in patients with AD compared with the other groups studied. Because the controls were significantly younger than the patients with AD, we investigated correlations between p-tau subtypes and age. In patients with AD, there was no correlation between the 3 p-tau subtypes and age ($\rho = -0.050$ to 0.002 ; $P = .61-.93$). In controls, p-tau₁₈₁ ($\rho = 0.61$; $P = .002$) and p-tau₂₃₁ ($\rho = 0.41$; $P = .05$) correlated with age,

Table 2. Correlations Between p-Tau Epitopes*

Group	p-Tau ₂₃₁ × p-Tau ₁₈₁	p-Tau ₂₃₁ × p-Tau ₁₉₉	p-Tau ₁₈₁ × p-Tau ₁₉₉
AD	0.807†	0.726†	0.734†
DLB	0.832†	0.800†	0.871†
FTD	0.571‡	0.193	0.360
VaD	0.857§	0.775§	0.847§
ONDs	0.433§	0.377	0.568‡
Control	0.632†	-0.069	0.384

Abbreviations: AD, Alzheimer disease; DLB, dementia with Lewy bodies; FTD, frontotemporal dementia; ONDs, other neurologic disorders; p-tau, phosphorylated tau protein; VaD, vascular dementia.

*Values are given as Spearman rho coefficients.

† $P < .001$.

‡ $P < .01$.

§ $P < .05$.

but p-tau₁₉₉ did not ($\rho = 0.05$; $P = .80$). Therefore, we repeated the analyses in a group of 23 patients with AD and 23 controls who were matched for age ($M-W_1 = 208$; $P = .21$) and sex ($\chi^2_1 < 0.001$; $P > .99$). Differences between the AD group and the control group remained unchanged and statistically significant for all 3 p-tau subtypes (p-tau₂₃₁: $M-W_1 = 32$; $P < .001$; p-tau₁₈₁: $M-W_1 = 52.5$; $P < .001$; p-tau₁₉₉: $M-W_1 = 46$; $P < .001$). Consequently, we included all of the patients with AD in our analyses. Correlations among the 3 p-tau subtypes are given in **Table 2**.

SENSITIVITY AND SPECIFICITY OF THE SINGLE MARKERS

Specificity levels of single markers using p-tau proteins when sensitivity was set at 85% or higher, as recommended by a consensus report, are given in **Table 3**.¹² In the case of p-tau₁₉₉, several individuals in each comparison group had a value equal to the 85% sensitivity cutoff value. Therefore, for p-tau₁₉₉, a lower and an upper limit is given for the specificity, corresponding to 2 alternatives: (1) all patients with this specific value were allocated to the AD group and (2) all patients with this specific value were allocated to the comparison group. In the differentiation of patients with AD from those in the non-AD group, p-tau₂₃₁ (85%) and p-tau₁₈₁ (81%) reached specificity levels of 75% or higher, but p-tau₁₉₉ did not (61%-72%). All 3 p-tau proteins showed excellent specificity levels when patients with AD were compared with those with ONDs and controls.

Comparing AD to FTD, p-tau₂₃₁ and p-tau₁₈₁ showed good discriminative power, with specificity levels of 92% and 79%, respectively. For p-tau₁₉₉, specificity ranged from 42% to 54% in the discrimination of AD from FTD. When AD was compared with DLB, p-tau₁₈₁ showed a specificity of 68%, whereas for p-tau₂₃₁ the specificity was 64%. For p-tau₁₉₉, specificity ranged between 50% and 64%. Values for AD vs VaD are not separately given owing to the small sample size of patients with VaD ($n = 7$).

We tested for differences in diagnostic accuracy among p-tau proteins (Table 3). Discriminative power between AD and the combined non-AD groups was significantly higher for p-tau₂₃₁ (McNemar test $P = .004$ to $P < .001$) and p-tau₁₈₁ (McNemar test $P = .10$ to $P < .001$) compared with p-tau₁₉₉. Diagnostic accuracy was also

significantly higher using p-tau₂₃₁ and p-tau₁₈₁ compared with p-tau₁₉₉ in differentiating AD from FTD (p-tau₂₃₁ vs p-tau₁₉₉: McNemar test $P = .004$ to $P < .001$; p-tau₁₈₁ vs p-tau₁₉₉: McNemar test $P = .07$ to $P = .01$). There was no statistically significant difference in diagnostic accuracy among p-tau proteins for discrimination between patients with AD vs ONDs and DLB. Receiver operating characteristic curves for pairwise comparisons of marker levels between patients with AD and comparison groups are shown in **Figure 2**.

COMBINATION OF THE MARKERS

We used classification tree analysis to investigate whether a combination of p-tau markers would improve group discrimination (**Table 4**). For AD vs the combined non-AD groups, AD vs controls, AD vs ONDs, and AD vs FTD, we found that p-tau₂₃₁ accounted for maximal group discrimination, whereas the other p-tau subtypes added no additional discriminatory power. Similarly, p-tau₁₈₁ alone maximized group separation between patients with AD and those with DLB and between the AD and VaD groups (data not shown). For discrimination between AD and non-AD dementias, a combination of p-tau₂₃₁ and p-tau₁₈₁ resulted in a slight increase in sensitivity from 86% to 94% at the cost of decreased specificity from 75% to 66%, resulting in only a slight increase in correct classification accuracy (from 83% to 85%) (**Figure 3**).

EFFECT OF AGE, SEX, MINI-MENTAL STATE EXAMINATION SCORE, AND CENTER ON CSF p-TAU LEVELS IN THE AD GROUP

Levels of CSF p-tau₂₃₁ ($\rho = -0.18$; $P = .06$) and p-tau₁₈₁ ($\rho = 0.002$; $P = .99$) did not correlate with the Mini-Mental State Examination score in patients with AD. Only for p-tau₁₉₉ did we find a correlation with the Mini-Mental State Examination score ($\rho = -0.25$; $P = .01$). There was no significant effect of sex (p-tau₂₃₁: $M-W_{107} = 1302$; $P = .55$; p-tau₁₈₁: $M-W_{107} = 1375$; $P = .89$; p-tau₁₉₉: $M-W_{107} = 1203$; $P = .22$) or age (p-tau₂₃₁: $\rho = 0.002$; $P = .98$; p-tau₁₈₁: $\rho = -0.05$; $P = .61$; p-tau₁₉₉: $\rho = -0.013$; $P = .89$) on levels of p-tau proteins. Levels of p-tau did not differ significantly between participating centers (p-tau₂₃₁: $M-W_{107} = 903$; $P = .74$; p-tau₁₈₁: $M-W_{107} = 822$; $P = .34$; p-tau₁₉₉: $M-W_{107} = 813$; $P = .31$).

COMMENT

In the present study, we investigated the diagnostic performance of 3 different pathophysiologically important CSF tau phosphorylation epitopes (p-tau₂₃₁, p-tau₁₈₁, and p-tau₁₉₉)²⁰⁻²² to discriminate patients with AD from those with other clinically important causes of dementia, those with ONDs, and nondemented controls. All 3 phosphorylation sites were studied using 3 recently developed and reported immunoassays.^{1,3,5} So far, independent early studies of the single phosphorylation sites have indicated statistically significant discriminative power between the AD and non-AD study groups. Subsequently, CSF p-tau proteins in general were suggested as promising biological marker candidates for AD. In none of these pilot stud-

Table 3. Specificity of Single Markers With Sensitivity Set at 85% or Higher and Differences in Diagnostic Accuracy Between the Single Markers*

AD Group	CSF Marker			P Value‡		
	p-Tau ₂₃₁	p-Tau ₁₈₁	p-Tau ₁₉₉ †	p-Tau ₂₃₁ vs p-Tau ₁₈₁	p-Tau ₂₃₁ vs p-Tau ₁₉₉	p-Tau ₁₈₁ vs p-Tau ₁₉₉
Non-AD (n = 98)	83 (85)	79 (81)	60-71 (61-72)	.39	<.01 to <.001	.10 to <.001
Control (n = 23)	22 (96)	21 (91)	17-21 (74-91)	.99	.99	.99
ONDs (n = 22)	21 (95)	19 (86)	19-20 (86-91)	.50	.99	.99
DLB (n = 22)	14 (64)	15 (68)	11-14 (50-64)	.99	.99	.99
FTD (n = 24)	22 (92)	19 (79)	10-13 (42-54)	.38	<.01 to <.001	.07 to <.05

Abbreviations: AD, Alzheimer disease; CSF, cerebrospinal fluid; DLB, dementia with Lewy bodies; FTD, frontotemporal dementia; ONDs, other neurologic disorders; p-tau, phosphorylated tau protein.

*Data are given as number (percentage) of patients. The cutoff levels for the different p-tau assays were 250 pg/mL for p-tau₂₃₁, 14.3 pmol/L for p-tau₁₈₁, and 1.1 fmol/mL for p-tau₁₉₉.

†Since several patients in each group had identical p-tau₁₉₉ values, 2 specificity figures are given for this assay.

‡Calculated using the McNemar test.




Figure 2. Receiver operating characteristic curves for cerebrospinal fluid phosphorylated tau protein (p-tau)₂₃₁, CSF p-tau₁₈₁, and CSF p-tau₁₉₉ when patients with Alzheimer disease were compared with the combined non-Alzheimer disease group (A), the control group (B), patients with other neurologic disorders (C), patients with dementia with Lewy bodies (D), and patients with frontotemporal dementia (E). Diagonal lines indicate an area of 50%, indicating no difference in marker levels between groups.

ies, however, were all 3 p-tau protein assays applied in the same set of subjects and patients to look at differential diagnostic assay accuracy. With the advent of international large-scale multicenter trials on putative biomarkers and neuroimaging in AD, the important issue has not yet been addressed of how p-tau assays perform in general and whether there are potential relevant differences in assay performance, particularly in diagnostic sensitivity and specificity. Moreover, it is not yet known

whether a combination of different p-tau epitopes or assays might improve diagnostic accuracy.

Our group showed that concentrations of all 3 p-tau proteins were equally significantly increased in patients with AD compared with the other groups studied. This finding is in strong agreement with all previously reported results.^{1,3,5,8-10}

In a next step, discriminative power of the individual p-tau proteins was studied. To interpret the clinical

Table 4. Sensitivity, Specificity, and Correctly Allocated Cases (CACs) for Group Comparisons Derived From Tree Analysis*

AD Group vs	Marker Step I	Sensitivity	Specificity	CACs	Marker Step II	Sensitivity	Specificity	CACs
Non-AD	p-Tau ₂₃₁	0.87	0.85	0.86	p-Tau ₁₈₁	0.95	0.77	0.86
Controls	p-Tau ₂₃₁	0.98	0.91	0.97				
ONDs	p-Tau ₂₃₁	0.96	0.91	0.95				
DLB/FTD/VaD	p-Tau ₂₃₁	0.86	0.75	0.83	p-Tau ₁₈₁	0.94	0.66	0.85
DLB	p-Tau ₁₈₁	0.94	0.64	0.88				
FTD	p-Tau ₂₃₁	0.88	0.92	0.89				

Abbreviations: AD, Alzheimer disease; DLB, dementia with Lewy bodies; FTD, frontotemporal dementia; Non-AD, all demented and nondemented non-AD groups studied; ONDs, other neurologic disorders; p-tau, phosphorylated tau protein; VaD, vascular dementia.

*The analysis considered all 3 p-tau markers simultaneously to create cutoff values that maximize separation between groups, resulting in a decision tree. The number of branches depends on the separation that has been achieved at the first split. To avoid overfitting of the data, generation of further branches was interrupted if sensitivity or specificity fell below 80% (eg, see AD vs non-AD and AD vs DLB/FTD/VaD).




Figure 3. Group discrimination for Alzheimer disease (AD) vs non-AD using phosphorylated tau protein (p-tau)₂₃₁ and p-tau₁₈₁ derived from tree analysis. The vertical and horizontal lines indicate the cutoffs for cerebrospinal fluid p-tau₂₃₁ and p-tau₁₈₁, respectively, derived from tree analysis.

cal significance of the determined diagnostic accuracy, we followed the recommendations of a consensus report¹² for useful (ideal) biomarkers of AD, determining specificity levels after the sensitivity level is 85% or higher. According to a consensus report on molecular and biochemical markers of AD, a useful (ideal) biomarker should yield a specificity level of at least 75% to 85%. In the differentiation of AD from non-AD, p-tau₂₃₁ and p-tau₁₈₁, but not p-tau₁₉₉, reached the recommended specificity level. All 3 p-tau proteins, however, showed excellent specificity levels when patients with AD were compared with nondemented controls.

For the clinically relevant differential diagnosis of AD, it is essential to have a marker or a set of markers that discriminates AD from other clinically relevant dementias. Therefore, we included patients with FTD, DLB, and VaD as well. In the discrimination between AD and FTD, p-tau₂₃₁ and p-tau₁₈₁ fulfilled the proposed biomarker criteria for sensitivity and specificity. Comparison of the markers revealed that p-tau₂₃₁ and p-tau₁₈₁ discriminated better than p-tau₁₉₉ between the AD group and the combined non-AD group and between AD and FTD.

The high level of discrimination between AD and FTD may originate in distinct differences in the bio-

chemical and molecular signatures of tau-related pathophysiological changes between the 2 diseases.²³ There was an increase in the p-tau protein level in some patients with DLB and VaD. Concomitant AD-type neuropathological changes in the brain, including neurofibrillary tangles, has been described for many patients with VaD²⁴ and DLB²⁵ who are clinically indistinguishable from those with “pure” VaD and DLB, respectively. In a clinical setting, it has to be assumed that patients with VaD and DLB are heterogeneous regarding underlying AD characteristic neuropathological changes in the brain, resulting in an increase in the p-tau protein level in at least some patients with VaD and DLB.

Using marker combinations did not add discriminative power compared with applying single markers. This might be a consequence of the high intercorrelation of the markers and the accurate discrimination between groups applied as single markers. Group separation was maximized between AD and FTD using p-tau₂₃₁ and between AD and DLB using p-tau₁₈₁.

In addition, we considered the effect of potentially confounding factors on CSF p-tau levels to assess the clinical applicability of p-tau proteins. Mini-Mental State Examination score accounted for approximately 5% of the variance in p-tau levels, being significant only for p-tau₁₉₉. This effect was not reported in previous studies,^{5,8} and it should be followed in independent samples. There was no effect of age and sex on levels of p-tau. In addition, different diagnostic centers did not affect variance of p-tau levels. These findings indicate that p-tau proteins may be valuable markers for the clinical diagnosis of AD irrespective of age, sex, and diagnostic center.

To our knowledge, this is the first comparative study applying 3 developed and published immunoassays detecting different p-tau epitopes on the same set of subjects and patients to compare the individual and combined diagnostic accuracy. The results of this study indicate that all 3 p-tau assays perform nearly equally well in discriminating patients with AD from nondemented controls. Both p-tau₂₃₁ and p-tau₁₈₁ fulfill the proposed criteria for useful biomarkers in the differentiation of AD and non-AD and particularly of AD and FTD.

Although there is no doubt that tau phosphorylation differs in AD, it is hard to speculate why. There have been few studies of phosphoserine 199 and phosphothreonine 181 in the human brain. Except for one study,²²

all we really know about these sites is that they are phosphorylated in advanced AD neuropathological changes. The antibodies used in the 181 and 199 assays have not been investigated to the same extent as the antibodies to phosphothreonine 231. It is well established that phosphorylation of threonine 231 is a very early event in AD, occurring before the formation of paired helical filaments in neurons of the hippocampus.²¹ According to Augustinack and colleagues,²² phosphorylation at threonine 181 and serine 199 occurs later, and these are only found to any appreciable extent in intracellular tangles. Reactivity to TG3, an antibody that recognizes phosphothreonine 231, is found in pretangles, intracellular tangles, and extracellular tangles and so is present at all stages of the disease. Augustinack and colleagues also suggest that several kinases can phosphorylate 199 and 231, but only extracellular regulated protein kinase 2 phosphorylates 181. We suggest further investigations of the temporal sequence of phosphorylations of the 3 sites, using the same antibodies as used in the CSF assays. Moreover, other potentially pathophysiologically relevant p-tau epitopes, such as serine 396 and serine 404, need to be further explored in their ability to differentiate between relevant dementia disorders.²⁶

Another relevant issue is to distinguish patients with mild cognitive impairment (MCI) from controls and particularly to predict AD in MCI. We showed that CSF p-tau₂₃₁ levels are elevated in patients with MCI compared with controls.²⁷ In this longitudinal study, high p-tau₂₃₁ levels at baseline correlated with the rate of cognitive decline in Mini-Mental State Examination scores in patients with MCI. A subgroup of patients with MCI converted to AD. In agreement with the analysis of rates of cognitive decline, increased levels of p-tau₂₃₁ correlated with conversion to AD. De Leon and colleagues²⁸ showed a longitudinal increase in p-tau₂₃₁ levels in patients with MCI. Elevated levels of CSF p-tau₁₈₁ in patients with MCI compared with controls have also been shown.²⁹ Future studies are warranted to further explore CSF p-tau proteins in MCI and to compare their diagnostic and prognostic value.

This study was conducted in an academic clinical setting. Diagnoses were performed by experienced dementia experts according to National Institute of Neurological and Communicative Disorders and Stroke-Alzheimer's Disease and Related Disorders Association criteria,¹³ with an estimated positive predictive value of 89% to 100%.³⁰ Part of our sample is enrolled in an ongoing neuropathological program designed to provide autopsy-confirmed diagnoses. In addition to autopsy-confirmed determination of assay performance, population-based studies are warranted to establish CSF p-tau proteins as potential biomarkers for routine diagnostic use. These studies are currently under way in international large-scale multicenter approaches. One large network, the National Institute on Aging Initiative on Neuroimaging in Alzheimer's Disease, will potentially start to evaluate neuroimaging this year, as well as an array of potential biomarkers in a 5-year longitudinal approach in 650 individuals (patients with MCI, patients with AD, and controls). In their recent proceedings, a subconsortium, the Biological Marker Working Group, has determined that measurement of CSF p-tau levels is a "fea-

sible core marker" within the National Institute on Aging initiative.³¹

Submitted for publication September 11, 2002; final revision received June 24, 2003; accepted July 3, 2003.

From the Dementia Research Section and Memory Clinic, Alzheimer Memorial Center and Geriatric Psychiatry Branch, Department of Psychiatry, Ludwig-Maximilian University, Munich, Germany (Drs Hampel, Buerger, Teipel, Goernitz, and Möller); Applied NeuroSolutions Inc, Vernon Hills, Ill (Drs Zinkowski, DeBernardis, and Kerkman and Ms McCulloch); Karolinska Institute, Neurotec, Division of Geriatric Medicine, Huddinge University Hospital, Stockholm, Sweden (Dr Andreason); the Department of Clinical Neuroscience, Unit of Neurochemistry, University of Göteborg, Sahlgren's University Hospital, Mölndal, Sweden (Drs Sjoegren and Blennow); Mitsubishi Kagaku Institute of Life Sciences, Tokyo, Japan (Dr Ishiguro); Mitsubishi Kagaku Medical, Tokyo (Dr Ohno); Innogenetics NV, Gent, Belgium (Drs Vanmechelen and Vanderstichele); and the Department of Pathology, Albert Einstein College of Medicine, Bronx, NY (Dr Davies). Drs Hampel and Davies are consultants for Applied NeuroSolutions Inc. Drs Zinkowski, DeBernardis, Kerkman, and Davies have stock and employee stock options and Ms McCulloch has employee stock options from Applied NeuroSolutions Inc.

This study was supported by grants from the Volkswagen-Foundation, Hannover, Germany (Dr Hampel); the Hirnliga e.V., Nürmbrecht, Germany (Drs Hampel and Buerger); a grant from the Medical Faculty, Ludwig-Maximilian University (Drs Buerger, Teipel, and Hampel); grants 11560 and 14002 from the Swedish Medical Research Council, Stockholm (Dr Blennow); and a grant from the Stiftelsen för Gamla Tjänarinnor, Stockholm (Dr Blennow).

We thank Felician Jancu, Bea Riemenschneider, Jenny Wagner, and Oliver Pogarell, MD, for clinical support; Thomas Nolde, PhD, and Heike Gluba for technical assistance; and Arun Lawrence Warren Bokde, PhD, and Jens Prüssner, PhD, for helpful discussion of the manuscript.

This study was presented in part at the 8th International Conference on Alzheimer's Disease and Related Disorders 2002, Alzheimer's Association; July 23, 2002; Stockholm, Sweden.

Corresponding authors: Katharina Buerger, MD, and Harald Hampel, MD, Dementia Research Section and Memory Clinic, Alzheimer Memorial Center and Geriatric Psychiatry Branch, Department of Psychiatry, Ludwig-Maximilian University, Nussbaumstrasse 7, 80336 Munich, Germany (e-mail: katharina.buerger@psy.med.uni-muenchen.de; hampel@psy.med.uni-muenchen.de).

Reprints: Raymond Zinkowski, PhD, Applied NeuroSolutions Inc, 50 Lakeview Pkwy, Vernon Hills, IL 60061 (e-mail: zinkowski@moleculargeriatrics.com).

REFERENCES

1. Kohnken R, Buerger K, Zinkowski R, Miller C, Kerkman D, DeBernardis J, Shen J, Moeller HJ, Davies P, Hampel H. Detection of tau phosphorylated at threonine 231 in cerebrospinal fluid of Alzheimer's disease patients. *Neurosci Lett*. 2000; 287:187-190.
2. Buerger K, Zinkowski R, Teipel SJ, Arai H, DeBernardis J, Kerkman D, McCulloch C, Padberg F, Faltraco F, Goernitz A, Tapiola T, Rapoport SI, Pirttila T, Moel-

- ler HJ, Hampel H. Differentiation of geriatric major depression from Alzheimer's disease with CSF tau protein phosphorylated at threonine 231. *Am J Psychiatry*. 2003;160:376-379.
3. Vanmechelen E, Vanderstichele H, Davidsson P, Van Kerschaver E, Van Der Perre B, Sjogren M, Andreasen N, Blennow K. Quantification of tau phosphorylated at threonine 181 in human cerebrospinal fluid: a sandwich ELISA with a synthetic phosphopeptide for standardization. *Neurosci Lett*. 2000;285:49-52.
 4. Schoenknecht P, Pantel J, Hunt A, Volkman M, Buerger K, Hampel H, Schroeder J. Levels of total tau and tau protein phosphorylated at threonine 181 in patients with incipient and manifest Alzheimer's disease. *Neurosci Lett*. 2003;339:172-174.
 5. Itoh N, Arai H, Urakami K, Ishiguro K, Ohno H, Hampel H, Buerger K, Wiltfang J, Otto M, Kretschmar H, Moeller HJ, Imagawa M, Kohno H, Nakashima K, Kuzuhara S, Sasaki H, Imahori K. Large-scale, multicenter study of cerebrospinal fluid tau protein phosphorylated at serine 199 for the antemortem diagnosis of Alzheimer's disease. *Ann Neurol*. 2001;50:150-156.
 6. Blennow K, Vanmechelen E, Hampel H. CSF total tau, Ab42 and phosphorylated tau protein as biomarkers for Alzheimer's disease. *Mol Neurobiol*. 2001;24:87-97.
 7. Blennow K, Hampel H. CSF markers for incipient Alzheimer's disease. *Lancet Neurol*. 2003;2:605-613.
 8. Buerger K, Zinkowski R, Teipel SJ, Tapiola T, Arai H, Blennow K, Andreasen N, Hofmann-Kiefer K, DeBernardis J, Kerkman D, McCulloch C, Kohnken R, Padberg F, Pirttila T, Schapiro MB, Rapoport SI, Moeller HJ, Davies P, Hampel H. Differential diagnosis of Alzheimer's disease with cerebrospinal fluid levels of tau protein phosphorylated at threonine 231. *Arch Neurol*. 2002;59:1267-1272.
 9. Sjogren M, Davidsson P, Tullberg M, Minthon L, Wallin A, Wikkelso C, Granerus AK, Vanderstichele H, Vanmechelen E, Blennow K. Both total and phosphorylated tau are increased in Alzheimer's disease. *J Neurol Neurosurg Psychiatry*. 2001;70:624-630.
 10. Vanmechelen E, Van Kerschaver E, Blennow K, De Deyn PP, Galasko D, Parnett L, Sindic CJM, Arai H, Riemenschneider M, Hampel H, Pottel H, Valgaeren A, Hulstaert F, Vanderstichele H. CSF-phospho-tau (181P) as a promising marker for discriminating Alzheimer's disease from dementia with Lewy bodies. In: Iqbal K, Sisodia S, Winblad B, eds. *Alzheimer's Disease: Advances in Etiology, Pathogenesis and Therapeutics*. New York, NY: John Wiley & Sons Inc; 2001:285-293.
 11. Hampel H, Buerger K, Kohnken R, Teipel SJ, Zinkowski R, Moeller HJ, Rapoport SI, Davies P. Tracking of Alzheimer's disease progression with CSF tau protein phosphorylated at threonine 231. *Ann Neurol*. 2001;49:545-546.
 12. Consensus report of the Working Group on: "Molecular and Biochemical Markers of Alzheimer's Disease": the Ronald and Nancy Reagan Research Institute of the Alzheimer's Association and the National Institute on Aging Working Group. *Neurobiol Aging*. 1998;19:109-116.
 13. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of the Department of Health and Human Services Task Force on Alzheimer's disease. *Neurology*. 1984;34:939-944.
 14. Lund and Manchester Groups. Clinical and neuropathological criteria for frontotemporal dementia. *J Neurol Neurosurg Psychiatry*. 1994;57:416-418.
 15. McKeith IG, Galasko D, Kosaka K, Perry EK, Dickson DW, Hansen LA, Salmon DP, Lowe J, Mirra SS, Byrne EJ, Lennox G, Quinn NP, Edwardson JA, Ince PG, Bergeron C, Burns A, Miller BL, Lovestone S, Collerton D, Jansen EN, Ballard C, de Vos RA, Wilcock GK, Jellinger KA, Perry RH. Consensus guidelines for the clinical and pathologic diagnosis of dementia with Lewy bodies (DLB). *Neurology*. 1996;47:1113-1124.
 16. Roman GC, Tatemichi TK, Erkinjuntti T, Cummings JL, Masdeu JC, Garcia JH, Amaducci L, Orgogozo JM, Brun A, Hofman A, et al. Vascular dementia: diagnostic criteria for research studies: report of the NINDS-AIREN International Workshop. *Neurology*. 1993;43:250-260.
 17. Morris JC, Heyman A, Mohs RC, Moody DM, O'Brien MD, Yamaguchi T, Grafman J, Drayer BP, Bennett DA, Fisher M, Ogata J, Kokmen E, Bermejo F, Wolf PA, Gorelick PB, Bick KL, Pajean AK, Bell MA, DeCarli C, Culebras A, Korszyn AD, Bogousslavsky J, Hartmann A, Scheinberg P. The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). *Neurology*. 1989;39:1159-1165.
 18. Breiman L, Friedman JH, Olshen RA, Stone CJ. *Classification and Regression Trees*. Belmont, Calif: Wadsworth International Group; 1984.
 19. Galasko D, Chang L, Motter R, Clark CM, Kaye J, Knopman D, Thomas R, Kholodenko D, Schenk D, Lieberburg I, Miller B, Green R, Basherad R, Kertiles L, Boss MA, Seubert P. High cerebrospinal fluid tau and low amyloid-beta-42 levels in the clinical diagnosis of Alzheimer disease and relation to apolipoprotein E genotype. *Arch Neurol*. 1998;55:937-945.
 20. Goedert M, Jakes R, Crowther RA, Cohen P, Vanmechelen E, Vandermeeren M, Cras P. Epitope mapping of monoclonal antibodies to the paired helical filaments of Alzheimer's disease: identification of phosphorylation sites in tau protein. *Biochem J*. 1994;301(pt 3):871-877.
 21. Vincent I, Zheng JH, Dickson DW, Kress Y, Davies P. Mitotic phosphoepitopes precede paired helical filaments in Alzheimer's disease. *Neurobiol Aging*. 1998;19:287-296.
 22. Augustinack JC, Schneider A, Mandelkow EM, Hyman BT. Specific tau phosphorylation sites correlate with severity of neuronal cytopathology in Alzheimer's disease. *Acta Neuropathol*. 2002;103:26-35.
 23. Delacourte A. Biochemical and molecular characterization of neurofibrillary degeneration in frontotemporal dementias. *Dement Geriatr Cogn Disord*. 1999;10(suppl 1):75-79.
 24. Kosunen O, Soininen H, Paljarvi L, Heinonen O, Talasniemi S, Riekkinen PJ Sr. Diagnostic accuracy of Alzheimer's disease: a neuropathological study. *Acta Neuropathol*. 1996;91:185-193.
 25. Gomez-Isla T, Growdon WB, McNamara M, Newell K, Gomez-Tortosa E, Hedley-Whyte ET, Hyman BT. Clinicopathologic correlates in temporal cortex in dementia with Lewy bodies. *Neurology*. 1999;53:2003-2009.
 26. Hu YY, He SS, Wang X, Grundke-Iqbal I, Iqbal K, Wang J. Levels of nonphosphorylated and phosphorylated tau in cerebrospinal fluid of Alzheimer's disease patients: an ultrasensitive bienzyme-substrate-recycle enzyme-linked immunosorbent assay. *Am J Pathol*. 2002;160:1269-1278.
 27. Buerger K, Teipel SJ, Zinkowski R, Blennow K, Arai H, Engel R, Hofmann-Kiefer K, McCulloch C, Ptok U, Heun R, Andreasen N, DeBernardis J, Kerkman D, Moeller H, Davies P, Hampel H. CSF tau protein phosphorylated at threonine 231 correlates with cognitive decline in MCI subjects. *Neurology*. 2002;59:627-629.
 28. De Leon MJ, Segal S, Tarshish CY, DeSanti S, Zinkowski R, Mehta PD, Convit A, Caraos C, Rusinek H, Tsui W, Saint Louis LA, DeBernardis J, Kerkman D, Qadri F, Gary A, Lesbre P, Wisniewski T, Poirier J, Davies P. Longitudinal cerebrospinal fluid tau load increases in mild cognitive impairment. *Neurosci Lett*. 2002;333:183-186.
 29. Andreasen N, Vanmechelen E, Vanderstichele H, Davidsson P, Blennow K. Cerebrospinal fluid levels of total-tau, phospho-tau and A beta 42 predicts development of Alzheimer's disease in patients with mild cognitive impairment. *Acta Neurol Scand Suppl*. 2003;179:47-51.
 30. Nagy Z, Esiri MM, Hindley NJ, Joachim C, Morris JH, King EM, McDonald B, Litchfield S, Barnetson L, Jobst KA, Smith AD. Accuracy of clinical operational diagnostic criteria for Alzheimer's disease in relation to different pathological diagnostic protocols. *Dement Geriatr Cogn Disord*. 1998;9:219-226.
 31. Frank RA, Galasko D, Hampel H, Hardy J, de Leon MJ, Mehta PD, Rogers J, Siemers E, Trojanowski JQ. Biological markers for therapeutic trials in Alzheimer's disease: proceedings of a working group: NIA initiative on neuroimaging in Alzheimer's disease. *Neurobiol Aging*. 2003;24:521-536.