

A Controlled Study of Cortical Gray Matter and Ventricular Changes in Alcoholic Men Over a 5-Year Interval

Adolf Pfefferbaum, MD; Edith V. Sullivan, PhD; Margaret J. Rosenbloom, MA; Daniel H. Mathalon, PhD, MD; Kelvin O. Lim, MD

Background: We report on structural brain changes during a 5-year period in healthy control and alcoholic men.

Methods: Alcoholic patients (n = 16), from an initial group of 58 who underwent brain magnetic resonance imaging scanning while in treatment, were rescanned with the same acquisition sequence approximately 5 years later. Control subjects (n = 28) spanning the same age range also were scanned twice at a comparable interval. Changes in brain volume were corrected for error due to differences in head placement between scans and expressed as slopes (cubic centimeters per year), percentage of change over baseline for the control subjects, and standardized change for the alcoholic patients. The alcoholic patients varied considerably in the percentage of time that symptoms of alcohol dependence were present and in the amount of alcohol consumed during follow-up.

Results: The cortical gray matter diminished in volume over time in the control subjects, most promi-

nently in the prefrontal cortex, while the lateral and third ventricles enlarged. The alcoholic patients showed similar age-related changes with a greater rate of gray matter volume loss than the control subjects in the anterior superior temporal lobe. The amount of alcohol consumed during follow-up predicted the rate of cortical gray matter volume loss, as well as sulcal expansion. The rate of ventricular enlargement in alcoholic patients who maintained virtual sobriety was comparable to that in the control subjects.

Conclusions: During a 5-year period, brain volume shrinkage is exaggerated in the prefrontal cortex in normal aging with additional loss in the anterior superior temporal cortex in alcoholism. The association of cortical gray matter volume reduction with alcohol consumption over time suggests that continued alcohol abuse results in progressive brain tissue volume shrinkage.

Arch Gen Psychiatry. 1998;55:905-912

From the Neuropsychiatry Program, Center for Health Sciences, SRI International, Menlo Park, Calif (Dr Pfefferbaum); the Department of Psychiatry and Behavioral Sciences, Stanford University School of Medicine, Stanford, Calif (Drs Pfefferbaum, Sullivan, Mathalon, and Lim and Ms Rosenbloom); and the Psychiatry Service, Veterans Affairs Palo Alto Health Care System, Palo Alto, Calif (Ms Rosenbloom and Dr Lim).

IN VIVO NEUROIMAGING has shown that long-term consumers of large amounts of alcohol have increased cerebrospinal fluid (CSF) spaces and reduced cortical brain volume affecting gray and white matter¹ compared with low alcohol consumers. Alcohol-related volume deficits are present in the frontal lobes,² anterior hippocampus,³ mammillary bodies and cerebellum,^{4,5} and corpus callosum,⁶ particularly in older persons. Although alcohol consumption contributes to these group differences in brain structure, the progression of the disease must also be considered relative to normal aging changes. Studies comparing young and old groups of healthy people^{7,8} or applying regression analysis to samples representing a continuum of ages⁹⁻¹¹ to estimate the rate of change over a given age span provide indirect evidence that the pas-

sage of time contributes to changes in brain structure.

Only longitudinal studies including patients and control subjects can provide direct information about disease progression relative to the normal changes of aging. To date, follow-up studies in alcoholic patients have included periods of treatment detoxification¹²⁻¹⁵ or 3 to 12 months after treatment.¹⁶⁻¹⁹ One study²⁰ retested alcoholic patients after 5 years but did not retest control subjects. The potential interaction between normal aging and long-term excessive alcohol consumption on brain structure can be inferred from cross-sectional analysis showing that the older brain is more vulnerable to the effects of alcohol than the younger brain.²¹ Unlike Alzheimer disease, which progresses inexorably, albeit at variable rates,²² alcoholism typically involves periods of exacerbation, during which ex-

PATIENTS, SUBJECTS, AND METHODS

PATIENTS AND SUBJECTS

Alcoholic patients (group 1) and control subjects (group 2) (**Table 1**) gave written informed consent for participation in the study.

Alcoholic Patients

The patients, recruited from a Veterans Affairs inpatient alcohol rehabilitation program, had previously participated in cross-sectional²¹ (n = 49) and longitudinal¹⁷ (n = 58) neuroimaging studies. At baseline, they met the *Research Diagnostic Criteria*²⁶ (RDC) for alcoholism but not for drug use disorder within the past year. They had no history of hospitalization for schizophrenia, major affective disorders, or medical or neurologic conditions affecting the central nervous system; seizure disorder unrelated to alcohol withdrawal; documented head injury with neurologic sequelae; or use of phenytoin or corticosteroids during the past month. The baseline scans were obtained 4 weeks (mean \pm SD, 28.7 \pm 5.5 days) after admission; 39 patients returned for a short-term follow-up study 2 to 12 months after discharge,¹⁷ and some participated in annual follow-up testing. Of the 16 patients in the present report, 15 were in the initial cross-sectional report, and 14 were also in the short-term follow-up report.

Efforts to recruit patients for a 5-year follow-up included an offer of overnight accommodation at the hospital and payment of \$200 to complete 2 days of testing. Twenty-two alcoholic patients were no longer in active treatment in the Veterans Affairs program or did not respond to mail or telephone contact. Of the remaining 36 patients, 6 had died, and 5 had moved out of state, leaving a sample of 25 targeted for testing; 19 underwent scanning, but the data from 3 could not be used because technical errors in the baseline or follow-up scan precluded a valid assessment of change over time. Therefore, group 1 included 16 patients. The follow-up sample was demographically and clinically comparable with the patients lost to follow-up but showed slightly greater brain volume abnormalities; the difference was significant for cortical gray matter (**Table 2**).

Control Subjects

To determine the amount of change occurring during a comparable period in healthy control subjects, we recalled men whose ages matched those of group 1 patients from a group of healthy community members who had served in earlier

studies of healthy aging⁹ and provided age norms for the initial cross-sectional studies of alcoholism.²¹ This sample of control subjects had been screened originally for psychiatric disorders by use of the Schedule for Affective Disorders and Schizophrenia–Lifetime version²⁷ and for medical or neurologic conditions affecting the central nervous system, including head injury with documented loss of consciousness exceeding 30 minutes, by physical examination, medical history, and standard blood tests. Subjects were excluded if they had ever met the RDC²⁶ for any psychiatric disorder, if they had experienced a drug use disorder during the year before entry into the study, or if they had ever consumed more than 54 g of ethanol per day (equivalent of 4 “drinks” containing an average of 13.6 g of ethanol) for a period exceeding 1 month as assessed through a semistructured interview used to measure alcoholic consumption history variables.^{28,29} Subjects 50 years and older were excluded if they scored 24 or less (maximum, 30) on the Mini-Mental State Examination.³⁰ Group 2 included 28 subjects.

Follow-up

Group 1 and 2 participants underwent a physical examination, completed a questionnaire to identify illness during the intervening period, underwent standard medical laboratory tests, and participated in the Structured Clinical Interview for *DSM-III-R* (SCID)³¹ and a semistructured interview to quantify alcohol consumption^{28,29} since baseline. At baseline, the interviewer had already estimated alcohol consumption to that point, starting from the age at which the participant first drank on a regular basis (at least 1 drink per month) and eliciting the quantity (how many drinks per day) and the frequency (how many drinks on average during a month) of drinking over a series of “drinking stages,” differentiating between normal and maximum quantities and their frequencies. The types of alcoholic beverage (wine, beer, or spirit) were converted into “drink” equivalents, and each was given a value of 13.6 g of absolute alcohol. For follow-up assessment, the interviewer first reviewed the patient’s or subject’s alcohol use history, oriented the participant to the point in time at which the chronology was being resumed, and then elicited information about subsequent epochs of sobriety and drinking, using the same format as the original questionnaire.

MAGNETIC RESONANCE IMAGING

Acquisition and Analysis

The same magnetic resonance imaging protocol^{9,32,33} on a 1.5-T scanner (Signa, General Electric, Milwaukee, Wis)

posure to alcohol is intensified, alternating with periods of sobriety.^{23,24} Thus the study of brain changes during the course of alcoholism must account not only for the age of the patient and the passage of time, but also for the exposure to alcohol. Alcoholic patients who continue regular heavy drinking should show greater change over time than those who maintain relative sobriety. Indeed, several recent studies^{17,18,20} have assessed the effects of such naturally occurring variations in outcome within alcoholic groups, yet the effects of extended so-

briety after periods of heavy drinking in restoring the brain to its “normal” aging course are seldom considered.

The present report gives data from a 5-year follow-up study of alcoholic patients, ranging in age from 31 to 67 years, originally tested after completing a 30-day treatment program. Some continued heavy drinking, while a few substantially reduced their alcohol consumption. Community men, initially recruited to provide age norms for the cross-sectional study, also were retested after a comparable interval.

was used at baseline and at follow-up: axial spin echo, 5-mm thick, 2.5-mm skip; field of view, 24 cm; 256 × 256 matrix; echo times, 20, 80 milliseconds; cardiac cycle-gated effective repetition times, more than 2400 milliseconds; 256 phase encodes; and oblique plane perpendicular to sagittal plane crossing through anterior and posterior commissures.

Images were processed without knowledge of the participant's identity, age, diagnosis, or neuroradiologist's report. The most inferior slice above the level of the orbits, where anterior horns of the lateral ventricles could be seen bilaterally, was identified as the index slice. Index slices for baseline and follow-up scans were individually reviewed to ensure comparability across the 2 scans. Seven consecutive slices, including the index slice and six superior slices, sampled approximately half the total volume of the brain. Each slice from the magnetic resonance image, divided into an inner 55% and an outer 45%, was segmented into CSF, gray matter, and white matter compartments by using a semiautomated image analysis technique.³² This analysis yielded 3 global cortical measures (gray matter, white matter, and sulcal CSF) based on the outer 45%, a measure of lateral ventricles (CSF in the inner 55% of all slices on which they appeared), and the third ventricle measured on the index slice or the slice below it, wherever it appeared larger.

The images were further divided according to anatomical landmarks and a priori geometric rules into 6 standardized cortical regions of interest (ROIs), encompassing the outer 45% of each image and corresponding roughly to lobar anatomy: prefrontal, frontal, anterior superior temporal, posterior superior temporal, anterior superior parietal, and posterior parietal-occipital (**Figure 1**).

Volume Change

Regional volumes were expressed as cubic centimeters. The volume difference between scans provides the first estimate of change over time. The simple difference between scans represents the true biological change between scans plus measurement error due to differences in the position of a person's head in the scanner between imaging sessions.³⁴ We used changes in head size (the sum of the volume of CSF and tissue on all 7 slices analyzed for this study) as an estimate of measurement error, because such change would not normally be expected in adults. To correct for measurement error, we regressed observed changes in CSF and gray and white matter volume for each region against the diagnostic group and differences in total head size between scans, assuming a common slope but different intercepts (if the groups differ in rate of change) for each group.^{17,34} We then added

each person's residual score for each region and tissue type measure to the group intercept for that measure to calculate the adjusted change.

Because patients and subjects had varying interscan intervals, adjusted change scores were divided by the interscan interval, expressing the change as a slope or rate of change per year. To control for absolute baseline differences in structure size between persons and groups and differences between the sizes of regions, adjusted change values were expressed as the percentage of change ([adjusted change/baseline] × 100). An additional procedure standardizing each ROI against change in group 2 was used to compare change across regions within group 1 because the ROIs were of fundamentally different sizes. Finally, slopes were multiplied by a uniform 5-year interval for all participants, and the resulting estimate of 5-year volume change was added to the corresponding baseline measure to yield corrected follow-up volumes. These absolute baseline and adjusted follow-up measures are used in **Figure 2** and **Figure 3** to provide a direct representation of the nature and scale of change occurring during the follow-up interval.

STATISTICAL ANALYSIS

The effects of normal aging were assessed in group 2 by using the percentage of change per year. One-sample *t* tests were performed to detect differences from zero in the 3 global cortical and 2 ventricular brain measures. To determine whether gray matter changed at a greater rate in any 1 of the 6 cortical ROIs than the others, a 1-way repeated-measures analysis of variance (ANOVA) for cortical gray matter was performed and followed up with *t* test comparisons of each ROI against the others if an interaction was found.

To determine whether brain changes progressed at different rates in the 2 groups, the rates of change (cubic centimeters per year) for the 3 global cortical and 2 ventricular brain measures were compared by using *t* tests. To determine whether gray matter loss progressed at a greater rate at any 1 of the 6 cortical ROIs in group 1 than in group 2, a 2-way repeated-measures ANOVA for the 6 cortical ROIs was performed, using standardized scores. Follow-up *t* tests comparing individual ROIs were performed for significant interactions.

To determine whether age at study entry, drinking behavior, or symptom severity during the follow-up period predicted outcome within group 1, these predictors were correlated with the rate of change (slopes). In addition to assessing the effect of drinking behavior as a continuous variable among patients in group 1, we performed an exploratory analysis comparing drinking-defined subgroups with each other and with the control group.

RESULTS

CLINICAL OUTCOME

The *DSM-III-R* diagnoses, SCID-based estimate of the amount of time patients had problems with alcohol, reports of the amount of alcohol consumed during the follow-up interval, and medical records indicated a wide range of outcomes (**Table 3** and **Table 4**). In group 1, 8 of 16 patients received additional inpatient treatment

at the Veterans Affairs Palo Alto Health Care System, Palo Alto, Calif, for alcohol dependence or other psychiatric (depression, posttraumatic stress disorder, anxiety disorders) and/or medical conditions. Four patients in group 1 maintained virtual sobriety during the follow-up period, and an additional 3 were within the control range of drinking. Cumulative but not necessarily continuous sober time during follow-up ranged from 5½ months to the entire period. Although the patients in group 1 differed widely among themselves, they also differed in ag-

gregate from group 2 in the amount of alcohol consumed during follow-up ($t_{42} = 2.46, P < .05$). Laboratory tests, physical examinations, and medical histories for group 2 indicated normal health status for age without serious untreated medical illness, although the SCID detected the emergence of Axis I diagnoses in 2 subjects (Table 3).

EFFECTS OF NORMAL AGING ON BRAIN VOLUMES

The effects of normal aging were assessed by using the percentage of change per year as the dependent variable. One-sample t tests for group 2 alone revealed significant differences from the presumed population mean of 0 for global cortical gray matter ($t_{27} = -3.37, P = .002$, mean = -0.008), third ventricle ($t_{27} = 4.62, P < .001$, mean = 0.08), and lateral ventricles ($t_{27} = 7.71, P < .001$, mean = 0.04) but not for the cortical sulci ($t_{27} = 0.95, NS$). Whether the overall cortical gray matter volume change varied by region was assessed by using a 1-way repeated-measures ANOVA, which yielded a significant region effect ($F_{5,135} = 4.39, P = .001$) (Figure 4, left). Follow-up 1-sample t tests revealed significant divergence from 0 for the prefrontal ($t_{27} = -5.16, P < .001$) and posterior parietal-occipital ($t_{27} = -2.46, P = .02$) gray matter volumes only. For 6 comparisons, with $\alpha = .05$, 2-tailed, the conservative Bonferroni significance level was $P = .008$.

Table 1. Demographic Characteristic of Participants at Study Entry*

	Group 1 (n = 16)	Group 2 (n = 28)
Age, y	45.2 ± 9.92 (26-61)	51.04 ± 13.79 (21-68)
Education, y†	13.78 ± 3.48 (8.5-19)	16.89 ± 2.54 (12-22)
Lifetime alcohol consumption, kg‡	1267 ± 809 (245-2824)	67 ± 78 (0-284)
NART-IQ ²⁵	109.25 ± 10.1 (95-125)	112.43 ± 6.5 (99-125)

*Data are given as mean ± SD (range). Group 1 was alcoholic patients; group 2, control subjects. NART indicates National Adult Reading Test.²⁵

† $P < .005$, 2-tailed t test.

‡ $P < .001$, 2-tailed t test.

GROUP DIFFERENCES IN RATE OF CHANGE IN REGIONAL BRAIN VOLUMES

Group differences in the rate of change (cubic centimeters per year) for the global cortical and ventricular ROIs approached significance only for the lateral ventricles ($t_{42} = 1.99, P = .053$).

To assess whether the cortical gray matter regions were differentially vulnerable to the passage of time in group 1, we standardized the rates of change for each ROI against the control values set to have a mean ± SD of 0 ± 1 . Thus, deviations from 0 within group 1 indicated disease-related changes beyond those observed with normal aging. The 2-group ANOVA for these standardized scores yielded a significant interaction ($F_{5,210} = 2.88, P = .02$) (Figure 4, right). The follow-up analyses, using t tests to compare rates of change (cubic centimeters per year), showed that only the anterior superior temporal gray matter volume decreased significantly faster in group 1 than in group 2 ($t_{42} = -2.88, P = .006$).

Figures 2 and 3 plot the absolute brain volumes underlying the derived and standardized change measures




Figure 1. Axial magnetic resonance images segmented into gray matter (dark gray), white matter (white), and cerebrospinal fluid (black). The curved white lines mark the division of each section into the outer 45% for cortical measures and inner 55% for ventricular measures. The horizontal white lines mark 3 coronal planes used to delineate 4 quadrants for defining the cortical regions of interest. These planes pass through the most anterior extreme of the genu of the corpus callosum, the most posterior extreme of the splenium of the corpus callosum, and midway between them. The 6 cortical regional measures are defined by summing quadrants across the slices as follows: a, prefrontal; b, frontal; c, anterior superior temporal; d, posterior superior temporal; e, anterior superior parietal; and f, posterior parietal-occipital.

Table 2. Mean ± SD Baseline Values for Alcoholic Patients Included in the Study (Followed Up) and Those Who Could Not Be Found (Not Followed Up)

Variable	Not Followed Up (n = 43)	Followed Up (n = 15*)	2-Tailed t_{56}	P
Age, y	45 ± 11	45 ± 10	0.17	<.93
National Adult Reading Test	105.5 ± 7.1	109.2 ± 10.4	1.51	<.14
Education, y	13 ± 2	14 ± 3.5	1.06	<.29
Lifetime alcohol consumption, kg	1281.4 ± 840.7	1189.0 ± 891.3	0.36	<.72
Age at onset, y	24.3 ± 10	28.5 ± 14.6	1.15	<.25
z Score				
Sulcal cerebrospinal fluid	0.84 ± 1.56	1.78 ± 1.91	1.91	<.06
Cortical gray matter	-0.42 ± 1.07	-1.31 ± 1.52	2.48	<.02
Cortical white matter	-0.64 ± 1.24	-0.82 ± 1.25	0.49	<.62
Lateral ventricles	0.749 ± 1.3	1.10 ± 1.83	0.82	<.42
Third ventricle	0.757 ± 1.74	1.24 ± 1.87	0.91	<.37

*Baseline z-score values from 1 alcoholic patient were not available for this analysis.

described. Two-group (groups 1 and 2) repeated-measure (baseline and follow-up) ANOVAs for these raw data yielded essentially the same results.

DRINKING BEHAVIOR AND BRAIN CHANGE OVER TIME

Estimates of total alcohol consumption, the mean number of drinks per month when drinking, and the amount of time during past 5 years a patient had problems with alcohol dependence were correlated with rate of change (slope) in global measures (cortical gray matter and sulci, lateral and third ventricles) and regional cortical gray matter measures. A greater total alcohol consumption was significantly associated with greater decreases in total cortical gray matter (Spearman $\rho = -0.52, P = .04$) (**Figure 5**), particularly in the frontal region (Spearman $\rho = -0.57, P = .03$). Heavier drinking (number of drinks per month

when drinking) was significantly associated with greater increases in cortical sulcal fluid (Spearman $\rho = 0.53, P = .04$) and greater decreases in the total cortical gray matter (Spearman $\rho = -0.49, P = .06$), particularly in the frontal region (Spearman $\rho = -0.51, P = .05$). The amount of time patients in group 1 experienced alcohol dependence symptoms was associated with a change in the total cortical gray matter (Spearman $\rho = -0.53, P = .04$), particularly in the prefrontal (Spearman $\rho = -0.51, P = .05$) and frontal (Spearman $\rho = -0.47, P = .07$) gray matter.

In an exploratory analysis, the patients in group 1 were divided into those who maintained virtual sobriety during the follow-up period ($n = 4$, consumed <5 kg of alcohol) and those who resumed drinking ($n = 12$, consumed >5 kg of alcohol) (Table 4). The rate of change per year in the global ROIs in the sober and drinking subgroups of group 1 and group 2 was compared by using ANOVA. The group effect was significant for the lateral ventricles ($F_{2,43} = 5.425, P < .01$); the change in the sober subgroup was not different from group 2, while the sober subgroup and group 2 differed from the drinking subgroup (**Figure 6**).

AGE AT BASELINE AND BRAIN CHANGE OVER TIME

Group 1 showed no relationships between age at baseline and the rate of change over time, with the exception of the anterior superior temporal gray matter, in which younger alcoholic patients showed more negative slopes ($r = 0.50, P = .02$). Among group 2 subjects, the age at baseline tended to predict the rate of change in the overall cortical gray matter volume during the 5 years ($r = 0.36, P = .057$) with a significant effect only




Figure 2. Estimated cerebrospinal fluid (CSF) volumes for the lateral ventricles (left) and cortical sulci (right) for 16 alcoholic patients (group 1) and 28 control subjects (group 2) at baseline and follow-up. Follow-up values have been adjusted for estimated measurement error. Group differences were significant for sulcal CSF ($F_{1,42} = 5.41, P < .05$) and approached significance for the lateral ventricles ($F_{1,42} = 3.22, P = .08$). Time effects ($F_{1,42} = 73.1, P < .001$) and the group by time interaction ($F_{1,42} = 5.2, P < .05$) were significant only for the ventricles.




Figure 3. Estimated gray matter volumes for 6 cortical regions for 16 alcoholic patients (group 1) and 18 control subjects (group 2) at baseline and at follow-up. Follow-up values have been adjusted for estimated measurement error. Group effects are significant for all but the posterior parietal-occipital region. Time effects are significant for all but the anterior parietal region. The group by time interaction is significant only for the anterior superior temporal region ($P = .006$).

at the prefrontal region ($r = 0.49$, $P = .009$). The direction of the association was that younger control subjects showed more negative slopes (ie, a greater loss of gray matter).

COMMENT

This naturalistic study reports on the change in the volume of cortical gray matter, cortical sulci, and ventricular CSF occurring during a 5-year period in men who met the RDC criteria for alcohol dependence at study entry and men specifically recruited and screened to serve as control subjects who consumed low amounts of alcohol. This study is unique in the length of its follow-up, the quantitative assessment of brain volume change with

a correction for measurement error, and the availability of follow-up data for control subjects so that changes in alcoholic patients can be studied relative to ongoing normal aging changes. The study is limited in that behavior during the follow-up period was assessed retrospectively, and only 25% of the original sample of alcoholic patients was available for retesting. Nevertheless, the follow-up sample was demographically and clinically comparable with the patients lost to follow-up, although the alcoholic patients described herein had slightly greater brain volume abnormalities than did the alcoholic patients who were lost to follow-up.

This longitudinal analysis of healthy men confirms the widely held belief, based on cross-sectional studies (reviewed by Raz³⁵), that the prefrontal cortex undergoes greater normal age-related gray matter loss than other cortical regions. Raz et al,³⁶ using a cross-sectional design, observed a gradient of age-related tissue volume decline greatest in the prefrontal gray matter, less in the primary motor and sensory cortex, and still less in the phylogenetically older brain regions, such as the limbic structures. As a later maturing and particularly plastic and malleable brain region, the prefrontal cortex may be especially susceptible to occult untoward events that accrue over a lifetime. The observations of magnetic resonance images are consistent with cognitive studies that commonly report age-related declines in tests assessing prefrontal cortical functions, including cognitive flexibility, working memory, and recall tasks requiring strategic search processes. The low correlation between ventricular enlargement and prefrontal volume loss ($r = -0.17$, $P = .37$) suggests that although the cortical and subcortical brain regions are susceptible to aging, these age-related effects proceed independently yet at similar rates. Indeed, a cortical-ventricular system independence has been noted in cross-sectional studies of aging.³⁷

Among the patients in group 1, alcohol consumption and health status during the follow-up period varied widely (Tables 3 and 4). Perhaps because of this variability, we found modest evidence, confined to the anterior superior temporal cortex, for accelerated brain deterioration over time in group 1 as a whole relative to group 2. However, among the patients in group 1, the amount

Table 3. Interval Between Scans and Diagnostic Status at Follow-up*

	Group 1 (n = 16)	Group 2 (n = 28)
Interval between scans, Mean ± SD (range), mo	64.6 ± 4.3 (59-74)†	61.04 ± 1.36 (55-62)
Current <i>DSM-III-R</i> diagnosis, No. of participants		
Alcohol dependent	5	0
Alcohol dependent (partial remission)	3	0
Alcohol dependent (full remission)	8	0
Major depression	2	1
Posttraumatic stress disorder	2	0
Obsessive-compulsive disorder	1	0
Social phobia	1	1
No diagnosis	0	26
Time during past 5 y had problems with alcohol, No. of participants		
Not at all	3	28
Rarely	3	0
Substantial minority of time	2	0
Half the time	2	0
Substantial majority of time	4	0
Almost all the time	2	0

*Group 1 was alcoholic patients; group 2, control subjects.
† $P < .005$, 2-tailed t test.

Table 4. Amount of Alcohol Consumed During Follow-up*

Amount, kg (No. of "Drinks")	Group 1		Group 2	
	No. of Patients per Group†	Mean Total Alcohol Consumption, kg	No. of Subjects per Group†	Mean Total Alcohol Consumption, kg
0-5 (0-368)	4	1.8	14	1.8
6-30 (441-2206)	1	16.4	8	14.4
31-60 (2279-4412)	2	55.3	6	49.8
61-100 (4485-7353)	5	79.7
101-150 (7426-11 029)	2	130.5
>150 (>11 029)	2	216 and 970‡
No. of drinks/mo when drinking, mean ± SD (range)	...	369 ± 437 (0-1782)
Total volume consumed, mean ± SD (range), kg	...	124 ± 233§ (0-970)	...	15.7 ± 19.71 ± (0-59)

*Group 1 was alcoholic patients; group 2, control subjects. Ellipses indicate not applicable.
†Denotes subgroup according to the number of drinks.
‡The amounts for each patient are listed separately.
§ $P = .02$, 2-tailed t test.




Figure 4. Left, Adjusted change in cortical gray matter volumes for 6 regions of interest for 28 control subjects (group 2). Change is expressed as a percentage of the baseline volume to standardize for differences in absolute size across each region and as the percentage of change per year to standardize for differences in follow-up intervals. Right, Change score for alcoholic patients (group 1) standardized to the change measured in group 2. P indicates prefrontal; F, frontal; AST, anterior superior temporal; PST, posterior superior temporal; AP, anterior superior parietal; and PPO, posterior parietal-occipital.




Figure 5. The relationship between cortical gray matter rate of change and the amount of alcohol consumed during the follow-up period (left) (Spearman $\rho = -0.52$, $P = .04$) and between the cortical gray matter rate of change and the estimated amount of time during past 5 years that alcoholic patients (group 1) met DSM-III-R criteria for alcohol dependence during the follow-up period (right) (Spearman $\rho = -0.53$, $P = .04$). One alcoholic patient who reported 950 kg of alcohol consumption is omitted from the left panel so as not to distort scaling. The darker circle represents two patients with overlapping values.

of alcohol consumed predicted the amount of cortical gray matter loss, particularly in the prefrontal and frontal brain regions. From the early neuropathological observations of Courville³⁸ through more recent neuropathological³⁹ and in vivo neuroradiological studies,^{2,19,40} the frontal lobes seem particularly affected in persons with chronic alcoholism. While our study provides an in vivo neuropathological basis for the fairly consistent observation of frontal executive dysfunction in detoxified alcoholic patients,⁴¹⁻⁴⁵ the small sample and the limited quantitative information about psychiatric and medical comorbid conditions during the follow-up period preclude a rigorous analysis of the relative contribution of these additional variables to the observed outcome. Similarly, the results of exploratory group analyses, which suggest that with essential sobriety over a 5-year period, the rates of change in ventricular volume become comparable to those seen in control subjects who consume low amounts of alcohol, call for replication in a larger sample.

Accepted for publication June 22, 1998.

This research was supported by grants AA05965, AA10723, AG11427, MH30854 from the National Institutes of Health, Bethesda, Md, and the clinical resources of the Veterans Affairs Palo Alto Health Care System, Palo Alto, Calif.

A preliminary report of these data was presented as a poster at the annual meeting of the Research Society on Alcoholism, San Francisco, Calif, July 20, 1997.

We are grateful to the following people who fulfilled important roles in recalling and testing the alcoholic patients and control subjects: Anajali Deshmukh, MD, Jennifer




Figure 6. Lateral ventricular volume rate of change during the follow-up interval for subgroups of alcoholic patients (group 1) and control subjects (group 2).

Jackson, BA, Leslie O'Neill, MA, and Amy Weimer, BA. In addition, Robert Zipursky, MD, as involved in initial patient recruitment, screening, and diagnosis, and Chung Nim Ha, BA, Dorothy Talbert Richardson, BA, and Stacy DeMent, BA, were involved in scanning and testing patients and control subjects at study entry. Brian Matsumoto, MA, assisted in image analysis and Kenneth Chow, MA, in data processing.

Reprints: Adolf Pfefferbaum, MD, Center for Health Sciences, SRI International, 333 Ravenswood Ave, Menlo Park, CA 94025 (e-mail: dolf@synapse.sri.com).

REFERENCES

1. Rosenbloom M, Pfefferbaum A, Sullivan E. Structural brain alterations associated with alcoholism. *Alcohol Health Res World*. 1996;19:266-272.
2. Pfefferbaum A, Sullivan EV, Mathalon DH, Lim KO. Frontal lobe volume loss ob-

- served with magnetic resonance imaging in older chronic alcoholics. *Alcohol Clin Exp Res*. 1997;21:521-529.
3. Sullivan EV, Marsh L, Mathalon DH, Lim KO, Pfefferbaum A. Anterior hippocampal volume deficits in nonamnesic, aging chronic alcoholics. *Alcohol Clin Exp Res*. 1995;19:110-122.
 4. Davila MD, Shear PK, Lane B, Sullivan EV, Pfefferbaum A. Mammillary body and cerebellar shrinkage in chronic alcoholics: an MRI and neuropsychological study. *Neuropsychology*. 1994;8:433-444.
 5. Shear PK, Sullivan EV, Lane BJ, Pfefferbaum A. Mammillary body and cerebellar shrinkage in chronic alcoholics with and without amnesia [abstract]. *J Int Neuropsychol Soc*. 1996;2:34-35.
 6. Pfefferbaum A, Lim KO, Desmond J, Sullivan EV. Thinning of the corpus callosum in older alcoholic men: a magnetic resonance imaging study. *Alcohol Clin Exp Res*. 1996;20:752-757.
 7. Murphy DGM, DeCarli C, Schapiro MB, Rapoport SI, Horwitz B. Age-related differences in volumes of subcortical nuclei, brain matter, and cerebrospinal fluid in healthy men as measured with magnetic resonance imaging. *Arch Neurol*. 1993;49:839-845.
 8. Strassburger TL, Lee HC, Daly EM, Szczepanik J, Krasuski JS, Mentis MJ, Salerno JA, DeCarli C, Schapiro MB, Alexander GE. Interactive effects of age and hypertension on volumes of brain structures. *Stroke*. 1997;28:1410-1417.
 9. Pfefferbaum A, Mathalon DH, Sullivan EV, Rawles JM, Zipursky RB, Lim KO. A quantitative magnetic resonance imaging study of changes in brain morphology from infancy to late adulthood. *Arch Neurol*. 1994;51:874-887.
 10. Jernigan TL, Archibald SL, Berhow MT, Sowell ER, Foster DS, Hesselink JR. Cerebral structure on MRI. 1: localization of age-related changes. *Biol Psychiatry*. 1991;29:55-67.
 11. Yue NC, Arnold AM, Longstreth WT, Elster AD, Jungreis CA, OLeary DH, Poirier VC, Bryan RN. Sulcal, ventricular, and white matter changes at MR imaging in the aging brain: data from the Cardiovascular Health Study. *Radiology*. 1997;202:33-39.
 12. Zipursky RB, Lim KO, Pfefferbaum A. MRI study of brain changes with short term abstinence from alcohol. *Alcohol Clin Exp Res*. 1989;13:664-666.
 13. Schroth G, Naegele T, Klose U, Mann K, Petersen D. Reversible brain shrinkage in abstinent alcoholics, measured by MRI. *Neuroradiology*. 1988;30:385-389.
 14. Schroth VG, Remmes U, Schupmann A. Brain shrinking in chronic alcoholism: CT follow-up study in 65 patients [in German]. *Fortschr Roentgenst*. 1985;142:363-369.
 15. Claus D, Wille HJ, Neundörfer B, Gmelin E. Is the enlargement of cerebrospinal fluid spaces reversible in abstinent alcoholics because of rehydration [in German]? *Klin Wochenschr*. 1987;65:185-193.
 16. Carlen PL, Wilkinson DA, Wortzman G, Holgate R. Partially reversible cerebral atrophy and functional improvement in recently abstinent alcoholics. *Can J Neurol Sci*. 1984;11:441-446.
 17. Pfefferbaum A, Sullivan EV, Mathalon DH, Shear PK, Rosenbloom MJ, Lim KO. Longitudinal changes in magnetic resonance imaging brain volumes in abstinent and relapsed alcoholics. *Alcohol Clin Exp Res*. 1995;19:1177-1191.
 18. Shear PK, Jernigan TL, Butters N. Volumetric magnetic resonance imaging quantification of longitudinal brain changes in abstinent alcoholics. *Alcohol Clin Exp Res*. 1994;18:172-176.
 19. Ron MA, Acker RW, Shaw GK, Lishman WA. Computerized tomography of the brain in chronic alcoholism: a survey and follow-up study. *Brain*. 1982;105:497-514.
 20. Muuronen A, Bergman H, Hindmarsh T, Telakivi T. Influence of improved drinking habits on brain atrophy and cognitive performance in alcoholic patients: a 5-year follow-up study. *Alcohol Clin Exp Res*. 1989;13:137-141.
 21. Pfefferbaum A, Lim KO, Zipursky RB, Mathalon DH, Lane B, Ha CN, Rosenbloom MJ, Sullivan EV. Brain gray and white matter volume loss accelerates with aging in chronic alcoholics: a quantitative MRI study. *Alcohol Clin Exp Res*. 1992;16:1078-1089.
 22. Stern RG, Mohs RC, Davidson M, Schmeidler J, Silverman J, Kramer-Ginsberg E, Searcey T, Bierer L, Davis KL. A longitudinal study of Alzheimer's disease: measurement, rate, and predictors of cognitive deterioration. *Am J Psychiatry*. 1994;151:390-396.
 23. Schuckit M, Smith TL, Anthenelli R, Irwin M. Clinical course of alcoholism in 636 male inpatients. *Am J Psychiatry*. 1993;150:786-792.
 24. Vaillant GE. A long-term follow-up of male alcohol abuse. *Arch Gen Psychiatry*. 1996;53:243-249.
 25. Nelson HE. *The National Adult Reading Test (NART)*. Windsor, Ontario: Nelson Publishing Co; 1982.
 26. Spitzer RL, Endicott J, Robins E. *Research Diagnostic Criteria (RDC)*. New York, NY: Biometrics Research, New York State Psychiatric Institute; 1975.
 27. Endicott J, Spitzer RL. A diagnostic interview: the Schedule for Affective Disorders and Schizophrenia. *Arch Gen Psychiatry*. 1978;35:837-844.
 28. Skinner HA. *Development and Validation of a Lifetime Alcohol Consumption Assessment Procedure*. Toronto, Ontario: Addiction Research Foundation; 1982.
 29. Skinner HA, Sheu WJ. Reliability of alcohol use indices: the lifetime drinking history and the MAST. *J Stud Alcohol*. 1982;43:1157-1170.
 30. Folstein MF, Folstein SE, McHugh PR. Mini-mental state: a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12:189-198.
 31. Spitzer RL, Williams JBW, Gibbon M, First MB. The structured clinical interview for DSM-III-R (SCID), I: history, rationale, and description. *Arch Gen Psychiatry*. 1992;49:624-629.
 32. Lim KO, Pfefferbaum A. Segmentation of MR brain images into cerebrospinal fluid spaces, white and gray matter. *J Comput Assist Tomogr*. 1989;13:588-593.
 33. Zipursky RB, Lim KO, Sullivan EV, Brown BW, Pfefferbaum A. Widespread cerebral gray matter volume deficits in schizophrenia. *Arch Gen Psychiatry*. 1992;49:195-205.
 34. Shear PK, Sullivan EV, Mathalon DH, Lim KO, Davis LF, Yesavage JA, Tinklenberg JR, Pfefferbaum A. Longitudinal volumetric computed tomographic analysis of regional brain changes in normal aging and Alzheimer's disease. *Arch Neurol*. 1995;52:392-404.
 35. Raz N. Neuroanatomy of the aging brain observed in vivo: a review of structural MRI findings. In: Bigler ED, ed. *Handbook of Human Brain Functioning: Neuroimaging*. Vol 2. New York, NY: Plenum Press; 1996:153-182.
 36. Raz N, Gunning FM, Head D, Dupuis JH, McQuain J, Briggs SD, Loken WJ, Thornton AE, Acker JD. Selective aging of the human cerebral cortex observed in vivo: differential vulnerability of the prefrontal gray matter. *Cereb Cortex*. 1997;7:268-282.
 37. Pfefferbaum A, Sullivan EV, Jernigan TL, Zipursky RB, Rosenbloom MJ, Yesavage JA, Tinklenberg JR. A quantitative analysis of CT and cognitive measures in normal aging and Alzheimer's disease. *Psychiatry Res Neuroimaging*. 1990;35:115-136.
 38. Courville CB. *Effects of Alcohol on the Nervous System in Man*. Los Angeles, Calif: San Lucas Press; 1955.
 39. Harper CG, Kril JJ. Neuropathology of alcoholism. *Alcohol Alcohol*. 1990;25:207-216.
 40. Nicolas JM, Estruch R, Salamero M, Orteu N, Fernandez-Sola J, Sacanella E, Urbano-Marquez A. Brain impairment in well-nourished chronic alcoholics is related to ethanol intake. *Ann Neurol*. 1997;41:590-598.
 41. Oscar-Berman M, Hutner N. Frontal lobe changes after chronic alcohol ingestion. In: Hunt WA, Nixon SJ, eds. *Alcohol-Induced Brain Damage, NIAAA Research Monographs 22*. Rockville, Md: Government Printing Office; 1993:121-156.
 42. Parsons OA, Butters N, Nathan PE, eds. *Neuropsychology of Alcoholism: Implications for Diagnosis and Treatment*. New York, NY: Guilford Press; 1987.
 43. Tarter RE. An analysis of cognitive deficits in chronic alcoholics. *J Nerv Ment Dis*. 1973;157:138-147.
 44. Nixon SJ, Parsons OA. Alcohol-related efficiency deficits using an ecologically valid test. *Alcohol Clin Exp Res*. 1991;15:601-606.
 45. Sullivan EV, Mathalon DH, Zipursky RB, Kersteen-Tucker Z, Knight RT, Pfefferbaum A. Factors of the Wisconsin Card Sorting Test as a test of dorsolateral prefrontal cortical function in schizophrenia and alcoholism. *Psychiatry Res*. 1993;46:175-199.