

# Common Genetic Vulnerability for Nicotine and Alcohol Dependence in Men

William R. True, PhD, MPH; Hong Xian, PhD; Jeffrey F. Scherrer, MA; Pamela A. F. Madden, PhD; Kathleen K. Bucholz, PhD; Andrew C. Heath, DPhil; Seth A. Eisen, MD, MSc; Michael J. Lyons, PhD; Jack Goldberg, PhD; Ming Tsuang, MD, PhD, DSc

**Background:** Nicotine and alcohol dependence often occur together. We examined data from male twin pairs to determine whether there are genetic or environmental influences common to nicotine and alcohol dependence, and, if so, to estimate the magnitude and correlation of these influences.

**Methods:** Subjects were 3356 male-male twin-pair members of the Vietnam Era Twin Registry who participated in a 1992 telephone administration of the Diagnostic Interview Schedule Version 3 Revised. Genetic model fitting was performed to estimate the magnitude and correlation of genetic and environmental contributions to lifetime nicotine and alcohol dependence.

**Results:** The heritability of nicotine dependence was

60.3% (95% confidence interval [CI], 55.4%-65.2%); that of alcohol dependence, 55.1% (95% CI, 49.7%-60.5%). The best-fitting model for the co-occurrence of lifetime nicotine and alcohol dependence included a substantial genetic correlation between both disorders ( $r = 0.68$ ; 95% CI, 0.61-0.74) and a modest unique environmental correlation ( $r = 0.23$ ; 95% CI, 0.14-0.32).

**Conclusions:** These data suggest a common genetic vulnerability to nicotine and alcohol dependence in men. This common genetic influence may partially explain the clinical and epidemiological observations that alcoholics are often dependent smokers.

*Arch Gen Psychiatry.* 1999;56:655-661

**T**HE RELATIONSHIP between smoking and alcoholism has long interested clinicians and researchers. Smoking is nearly universal in treatment-seeking alcoholic samples<sup>1</sup> and has been reported to be as high as 85% in alcoholics identified in a general medical clinic.<sup>2</sup> The average number of cigarettes consumed is also significantly higher among alcoholics compared with nonalcoholic smokers,<sup>1</sup> with rates of heavy smoking (eg, >25 cigarettes per day) among adult alcoholics as high as 90%.<sup>3,4</sup>

Tobacco use and heavy alcohol consumption represent a public health concern. Independent and combined use of these substances contribute to several types of cancer,<sup>5</sup> and synergistic risks are particularly high for head and neck cancers.<sup>6,7</sup> Smoking is also associated with increased risk for alcoholism relapse and continued problems with other substance abuse.<sup>8,9</sup>

Genetic contributions to smoking initiation, persistent smoking, and alcohol dependence have been established in many twin studies. Heritability estimates for

smoking initiation in male twins range from 46% to 84%,<sup>10-13</sup> and estimates of the genetic contribution to long-term persistent smoking range from 58% to 74%.<sup>11,13</sup> Evidence from twin studies performed in Scandinavia, Australia, and the United States indicates that genes account for 50% to 60% of the variance in risk for alcohol dependence in women<sup>14,15</sup> and men,<sup>14,16-18</sup> including data from our study sample, members of the Vietnam Era Twin (VET) Registry.<sup>17</sup> Although we have previously estimated the genetic contribution to alcohol dependence in the VET Registry,<sup>17</sup> we have not addressed the heritability of nicotine dependence or its relationship with alcohol dependence.

A close relationship between smoking and the excessive use of alcohol has been recognized for many years,<sup>19,20</sup> but the underlying mechanisms have received little investigation. Studies in rodents have found that strains of mice bred for alcohol sensitivity have also been observed to be sensitive to nicotine.<sup>21-23</sup> However, there has been little investigation of potential common genetic influences on nicotine and alcohol dependence in humans. Swan

*The affiliations of the authors appear in the acknowledgment section at the end of the article.*

## SUBJECTS AND METHODS

### SAMPLE

The VET Registry consists of 7375 male-male twin pairs born from January 1, 1939, through December 31, 1955, of whom both siblings served on active military duty during the Vietnam era (1965-1975). The characteristics of the VET Registry have been reported elsewhere.<sup>26-28</sup> In 1992, approximately 5000 twin pairs of the VET Registry were invited to participate in a computer-assisted telephone administration of the Diagnostic Interview Schedule Version 3 Revised (DIS-3R),<sup>29</sup> which allowed derivation of psychiatric diagnoses according to *DSM-III-R* criteria.<sup>30</sup> Interviewers contacted twins and began interviewing after verbal informed consent was obtained, a method approved by the institutional review boards at participating universities.

Lifetime nicotine and alcohol dependence diagnoses were obtained by applying standard DIS-3R computer algorithms to the data. To assess reliability of the nicotine and alcohol dependence diagnoses, a subset of 146 individual twins were reinterviewed by a second interviewer. The mean time between interviews was 466 days (SD,  $\pm 50.5$  days). Reliability was assessed using the  $\kappa$  statistic. According to the guidelines for rating reliability of Fleiss,<sup>31</sup> the test-retest reliability of the nicotine dependence diagnosis was excellent ( $\kappa = 0.76$ ), and the test-retest reliability for alcohol dependence was good ( $\kappa = 0.61$ ).<sup>32</sup>

The DIS-3R diagnosis of alcohol dependence in VET Registry members has been reported to have good criterion validity, with 96% of patients treated for alcoholism in a Veterans Affairs hospital also receiving a diagnosis of alcoholism using the DIS-3R algorithm.<sup>32</sup> Although similar data are not available for nicotine dependence, the content validity of the DIS-3R diagnosis of nicotine dependence appears good. For example,

among nicotine-dependent veterans, 43.1% reported failed cessation attempts, and 41.8% were heavy smokers ( $>30$  cigarettes per day) and reported a mean of 3.75 nicotine withdrawal symptoms. In comparison, 19.6% of non-nicotine-dependent smokers were unable to quit, and 7.1% were heavy smokers and reported a mean of 0.59 withdrawal symptoms.

Eligibility criteria for the present analyses were the following: (1) both members of the twin pair were identified from Department of Defense computer files<sup>33</sup>; (2) both members of the pair completed all tobacco and alcohol use questions from the 1992 interview; and (3) zygosity could be definitively assigned. The final analyses were conducted using a sample of 3356 twin pairs (1864 were monozygotic [MZ]; 1492, dizygotic [DZ]).

The mean age of respondents at interview in 1992 was 44.6 years (SD,  $\pm 2.8$  years; range, 36-55 years). Most (90.4%) were non-Hispanic white; 4.9%, African American; 2.7%, Hispanic; 1.3%, native American or Alaskan native; and 0.7%, other. High school graduates accounted for 33.3%; college graduates, 38.6%. Most (92.6%) were employed full-time; 1.8% were employed part-time; and 5.6% were not employed at the time of interview.

### STATISTICAL ANALYSIS

Three sources of influences accounting for individual differences are additive genetic effects (denoted A for statistical analysis), shared environmental effects (denoted C for statistical analysis), and unique environmental effects (denoted E for statistical analysis).<sup>34</sup> Additive genetic influences are correlated 100% between members of an MZ twin pair and 50% between members of a DZ twin pair. Shared environmental influences are experiences that twins have in common such as exposure to parental smoking or drinking, attending the same school or church, or living in the same neighborhood. Shared environmental influences are assumed to contribute to similarity equally in MZ and DZ twin pairs. Finally, unique environmental influences are

et al<sup>24</sup> reported a genetic correlation between smoking and alcohol consumption of 0.47 in male World War II US veteran twins. In a cohort of female twins, Prescott and Kendler<sup>25</sup> found family environmental experiences modestly contributed to nicotine and alcohol dependence, with only very small genetic influences common to both substances. These contrasting studies suggest further research is needed to elucidate the genetic and environmental influences common and specific to nicotine and alcohol dependence.

From the perspective of prevention of smoking- and alcohol-related diseases, the co-occurrence of nicotine and alcohol dependence is an important priority for study. To address this issue, we investigated whether there are common genetic and/or family environmental contributions to lifetime history of nicotine and alcohol dependence in 3356 male-male twin pairs from the VET Registry.

## RESULTS

Among those who met criteria for nicotine and alcohol dependence, the mean ages of onset for regular smok-

ing and alcohol dependence were 17.1 years (SD,  $\pm 3.1$  years) and 20.4 years (SD,  $\pm 4.7$  years), respectively. The lifetime prevalence of nicotine and alcohol dependence was 47.8% and 35.2%, respectively. Lifetime prevalence of nicotine dependence for MZ twins was 46.2%; for DZ twins, 49.7%. Lifetime prevalence of alcohol dependence for MZ twins was 34.8%; for DZ twins, 35.8%. Among those with nicotine dependence, 50.9% met criteria for alcohol dependence compared with those without nicotine dependence, among whom 20.8% had a diagnosis of alcohol dependence. Among persons with a lifetime diagnosis of nicotine dependence, 62.6% were current smokers at interview. Characteristics of smoking and drinking behaviors were similar for MZ and DZ twin pairs (**Table 1**).

For MZ twin pairs, the tetrachoric correlation for nicotine dependence was 0.61; for DZ twin pairs, 0.31. For alcohol dependence, the tetrachoric correlation for MZ twin pairs was 0.55; for DZ twin pairs, 0.29. In each case, the MZ twin pair correlation was approximately twice that of the DZ twin pair correlation, which suggested that both phenotypes were influenced by additive genetic effects.

nonshared experiences that contribute to differences within MZ and DZ twin pairs. Unique environmental influences are uncorrelated between twin pairs and include measurement error. The greater similarity for a phenotype among MZ twins compared with DZ twins, as shown by a higher correlation coefficient, suggests genetic influences, as found in our analysis.

Therefore, we used model fitting to resolve competing hypotheses that nicotine and alcohol dependence are influenced by genes and environment, with environmental influences across a pair being uncorrelated (AE); environmentally determined with some environmental elements resulting from experiences shared equally by both members of a twin pair (CE); or influenced by genes, shared environment, and unique environment (ACE).

We began by computing twin-pair contingency tables separately for nicotine and alcohol dependence. Tetrachoric correlations calculated for each disorder by zygosity were computed under the assumption of a multifactorial threshold model in which multiple unmeasured genetic and environmental risk factors determine an underlying continuous liability for development of nicotine or alcohol dependence. The liability threshold model assumes that there is a single normally distributed dimension of liability with abrupt thresholds. Above a given threshold, the disorder is expressed, and below the threshold, the disorder does not develop.

Univariate genetic path models were fit to the contingency tables using maximum likelihood.<sup>34</sup> Univariate models estimated the proportion of variance attributable to additive genetic, shared environmental, and unique environmental effects, including error. Submodels were fit that deleted genetic (CE) or shared environmental parameters (AE). The goodness of fit of full models (ACE) and submodels (AE or CE) was determined using likelihood ratio  $\chi^2$ , with degrees of freedom for the  $\chi^2$  difference between the full model and submodel computed as degrees of freedom of the submodel minus degrees of freedom of the full model.

As shown in **Table 2**, a full model that allowed for additive genetic, shared, and unique environmental influences on liability to nicotine dependence (ACE) and a reduced model that did not allow for shared environmental influences (AE) produced good fits to the nicotine dependence data ( $\chi^2_3 = 6.79$  [ $P = .08$ ] and  $\chi^2_4 = 6.80$  [ $P = .15$ ], respectively). The AE model was chosen as the best-fitting univariate model for nicotine dependence, since it gave a more parsimonious fit to the data and was not significantly worse fitting than the full model ( $\Delta\chi^2_1 = 0.01$  [ $P > .10$ ]). A model that did not allow for additive genetic contributions (CE) gave a significantly poorer fit to the data compared with the full model and was rejected at  $P < .001$ .

For alcohol dependence, the full and reduced models produced good fits to the data ( $\chi^2_3 = 4.20$  [ $P = .24$ ] and  $\chi^2_4 = 4.38$  [ $P = .36$ ], respectively). The more parsimonious reduced model was not significantly worse fitting than the full model ( $\Delta\chi^2_1 = 0.18$  [ $P > .10$ ]) and was chosen as best fitting the data. A model that did not allow for additive genetic contributions resulted in a poor fit ( $P < .001$ ).

A bivariate genetic model<sup>34</sup> was fitted to estimate the genetic and environmental correlations between nicotine and alcohol dependence. For bivariate modeling, a  $4 \times 4$  correlation matrix was calculated for nicotine and alcohol dependence data of MZ and DZ twin pairs. This resulted in 2 within-diagnosis cross-twin correlations, 2 within-twin cross-diagnosis correlations, and 2 cross-twin cross-diagnosis correlations.

Bivariate analyses compared the fit of the full model (ACE) for nicotine and alcohol dependence with that of reduced models that removed 1 or more genetic (A) or environmental (C and E) parameters. A  $\chi^2$  difference statistic again determined the best fitting model. If 2 or more reduced models were observed to give an adequate fit for the data, the model with the lowest Akaike information criterion<sup>35</sup> was accepted as the best fitting model. MX software<sup>36</sup> was used for the genetic modeling, and the PRELIS 2 program<sup>37</sup> was used to compute tetrachoric correlations and asymptotic covariance matrices.

#### DATA WEIGHTING TO ACCOUNT FOR RESPONSE BIAS

Since nonresponse may introduce bias in our genetic model fitting, we computed weights to account for systematic nonresponse to the 1992 interview. Nonresponders were twins who responded to a 1987 survey but not to the 1992 interview. Responders were twins who participated in both data collection efforts. Nonresponders in 1992 were more likely than responders to be persistent smokers (64.9% vs 57.1%), to have ever smoked (74.1% vs 68.7%), to report drinking more than 3 times per week (39.3% vs 31.8%), and to report heavy drinking ( $\geq 6$  drinks per drinking occasion) (25.4% vs 20.3%). A response propensity analysis approach<sup>38</sup> was used to generate sampling weights to account for response bias effects. We calculated tetrachoric correlations by zygosity for nicotine and alcohol dependence to compare weighted and unweighted correlations. Since no differences were observed, all analyses were reported using unweighted data.

Under the full univariate models, genes accounted for 59.6% (95% confidence interval [CI], 41.5%-65.4%) and 51.5% (95% CI, 32.1%-60.5%) of the variance in risk for nicotine and alcohol dependence, respectively. Under the full models, shared family environmental influences did not significantly account for variance in risk for nicotine (1.0% [95% CI, 0.0%-16.5%]) or alcohol dependence (3.5% [95% CI, 0.0%-20.0%]). The remainder of variance for both disorders was due to unique environmental factors that included measurement error.

Bivariate analyses determined the degree to which additive genetic and environmental contributions to nicotine and alcohol dependence were correlated. The bivariate model-fitting results are portrayed in **Table 3**. A model that allowed for genetic and unique environmental influences to nicotine and alcohol dependence but did not allow for shared environmental influences produced the most parsimonious fit to the data ( $\chi^2_8 = 12.73$  [ $P = .12$ ]; Akaike information criterion, -3.28).

For the best-fitting bivariate model, we calculated the variance due to genetic and environmental influences and the genetic and environmental correlations for

**Table 1. Characteristics of Smoking and Alcohol Use in Nicotine- and Alcohol-Dependent Members of the Vietnam Era Twin Registry by Zygosity\***

Characteristics	Subjects	
	MZ	DZ
<b>Nicotine-Dependent Subjects</b>		
	<b>(n = 1722)</b>	<b>(n = 1484)</b>
Age smoked regularly, y	17.41 ± 3.15	17.47 ± 3.31
No. of cigarettes per day when smoking the most in lifetime	32.11 ± 16.12	32.40 ± 15.61
No. of lifetime nicotine dependence symptoms	4.19 ± 1.07	4.21 ± 1.09
Percentage who ever tried to quit or cut down smoking	96.2 ± 19.1	95.5 ± 20.8
If ever tried, percent who reported that they could not quit or cut down	43.0 ± 49.5	43.1 ± 49.5
If unable to quit, percent who tried to cut down several times	83.9 ± 36.8	82.5 ± 38.0
<b>Alcohol-Dependent Subjects</b>		
	<b>(n = 1296)</b>	<b>(n = 1068)</b>
Age first had 1 drink 1 time per mo for ≥6 mo or more (ie, began drinking regularly), y	17.75 ± 2.88	17.66 ± 2.75
Age first had ≥20 drinks in a day, y	22.12 ± 5.18	21.85 ± 5.04
No. of lifetime alcohol dependence symptoms	4.93 ± 1.74	4.96 ± 1.78
No. of drinks on typical drinking day during period of heaviest period of consumption in lifetime	10.78 ± 9.12	11.21 ± 9.50
No. of days per month drinking during heaviest period of consumption in lifetime	15.59 ± 10.18	15.97 ± 10.28
Percentage who ever tried to quit or cut down on drinking	80.5 ± 39.7	78.4 ± 41.2
If ever tried, percent unable to quit or cut down on drinking	17.9 ± 38.4	18.4 ± 38.8
If unable to quit, percent who were unable to quit or cut down more than once	61.5 ± 48.8	68.8 ± 46.5

\*Data are given as mean ± SD. MZ indicates monozygotic; DZ, dizygotic. The numbers reported for each zygosity include complete twin pairs and individuals from incomplete pairs. Numbers vary from the total number of pairs (N = 3356 pairs; 1864 MZ; 1492 DZ) analyzed in the genetic model fitting since only complete twin pairs, which included nonaffected twins of each zygosity, were included in the model-fitting analyses.

**Table 2. Univariate Genetic Model-Fitting Results for Full and Reduced Models of Nicotine and Alcohol Dependence for 3356 Male-Male Twin Pairs\***

Phenotype	Model	Goodness of Fit		Parsimony AIC
		$\chi^2_{df}$	P	
Nicotine dependence	ACE	6.786 <sub>3</sub>	.08	0.786
	<b>AE</b>	<b>6.800<sub>4</sub></b>	<b>.15</b>	<b>-1.200</b>
	CE	49.273 <sub>4</sub>	<.001	41.273
Alcohol dependence	ACE	4.203 <sub>3</sub>	.24	-1.797
	<b>AE</b>	<b>4.375<sub>4</sub></b>	<b>.36</b>	<b>-3.625</b>
	CE	31.56 <sub>4</sub>	<.001	23.558

\*A indicates additive genetic factors; C, shared environmental factors; E, unique environmental factors; and AIC, Akaike information criterion. The model with the lowest AIC was accepted as the best-fitting model (indicated in boldface).

lifetime nicotine and alcohol dependence. The best-fitting bivariate model (**Figure**) included a substantial genetic correlation between both disorders ( $r = 0.68$ ; 95% CI, 0.61-0.74) and a modest unique environmental correlation ( $r = 0.23$ ; 95% CI, 0.14-0.32). Under this model, genes accounted for 60.3% (95% CI, 55.4%-65.2%) and 55.1% (95% CI, 49.7%-60.5%) of the variance in risk for lifetime nicotine and alcohol dependence, respectively. Unique environmental influences, which included measurement error, accounted for 39.7% (95% CI, 34.8%-44.6%) and 44.9% (95% CI, 39.5%-50.3%) of the variance in risk for lifetime nicotine and alcohol dependence, respectively.

Under this model, we computed the amount of genetic variance in risk for alcohol dependence that is common to nicotine dependence, where  $r_A$  is the genetic correlation coefficient and  $h_a$  is the total genetic variance in risk for alcohol dependence. Thus,  $[r_A \times (h_a)^{1/2}]^2 = [0.68 \times (55.1\%)^{1/2}]^2 = 25.5\%$  of the total variance in risk for al-

cohol dependence overlaps with the genetic risk for nicotine dependence. Similarly, we found 2.4% of the variance in risk for alcohol dependence overlapped with the unique environmental influences to nicotine dependence.

#### COMMENT

The co-occurrence of lifetime nicotine and alcohol dependence among middle-aged male twins was best explained by a model that included a substantial genetic correlation ( $r = 0.68$ ) between both disorders, allowed for no shared family environmental influences, and evidenced considerable additive genetic contributions to nicotine and alcohol dependence (60.3% and 55.1%, respectively). Unique environmental influences were modestly correlated for both disorders ( $r = 0.23$ ). Of the total variance in risk for alcohol dependence, 25.5% was common with the genetic influence to nicotine dependence, but only 2.4% of the total variance in risk for alcohol dependence was common with the unique environmental influences to nicotine dependence.

Heritability estimates of 55% obtained in this sample for alcohol dependence are within the range of 39% to 60% previously reported in other twin studies.<sup>14-17</sup> We are not aware of previous reports of the heritability of DSM-III-R nicotine dependence, but heritability estimates of 58% to 74% have been reported for smoking persistence,<sup>11,13</sup> which may be related to the heritability of nicotine dependence estimated in our study. We recognize that the heritability estimates obtained may be limited by the reliability of each diagnosis. The estimate of the common genetic contribution to alcohol and nicotine dependence may vary with the reliability of the diagnoses.

Our finding that the liability for alcohol dependence is partially explained by additive genetic influ-

**Table 3. Genetic Model Fitting Comparisons for Full and Reduced Models of Additive Genetic (A), Shared Environmental (C), and Unique Environmental (E) Influences to Comorbid Lifetime Nicotine Dependence and Lifetime Alcohol Dependence for 3356 Male-Male Twin Pairs**


Model		Phenotypic Correlations*			Goodness of Fit		Parsimony, AIC
Nicotine	Alcohol	$r_A$	$r_C$	$r_E$	$\chi^2_{df}$	P	
ACE	ACE	0.62	1.0	0.25	10.03 <sub>5</sub>	.07	0.03
ACE	ACE	†	1.0	0.34	33.90 <sub>6</sub>	<.001	21.90
ACE	ACE	‡	0.22	0.21	22.31 <sub>6</sub>	<.01	10.31
ACE	ACE	0.68	†	0.23	12.72 <sub>6</sub>	.05	0.72
ACE	ACE	0.62	‡	0.25	10.03 <sub>6</sub>	.12	-1.97
ACE	ACE	0.76	1.0	†	36.06 <sub>6</sub>	<.001	24.06
<b>A&amp;E</b>	<b>A&amp;E</b>	<b>0.68</b>	<b>NA</b>	<b>0.23</b>	<b>12.73<sub>8</sub></b>	<b>.12</b>	<b>-3.28</b>
A&E	A&E	†	NA	0.59	325.92 <sub>9</sub>	<.001	307.92
A&E	A&E	‡	NA	0.05	88.24 <sub>9</sub>	<.001	70.24

\* $r_A$  indicates genetic correlation between lifetime nicotine dependence and lifetime alcohol dependence;  $r_C$ , shared family environmental correlation between lifetime nicotine dependence and lifetime alcohol dependence;  $r_E$ , unique environmental correlation between lifetime nicotine dependence and lifetime alcohol dependence; AIC, Akaike information criterion; and NA, not applicable. The model with the lowest AIC was accepted as the best-fitting model (indicated in boldface).

†Indicates correlation fixed to 0.

‡Indicates correlation fixed to 1.

§Indicates parameter C fixed to 0.



General bivariate model of risk for lifetime nicotine and alcohol dependence for 3356 male-male twin pairs. The magnitude of genetic ( $r_A$ ) and unique environmental correlations ( $r_E$ ) (95% confidence intervals [CIs]) between lifetime nicotine and alcohol dependence and the variance (95% CI) in risk for lifetime nicotine and alcohol dependence that is due to genes (A) and unique environmental (E) effects (the latter includes measurement error) are seen.

ences in common with nicotine dependence is consistent with reports from other researchers that alcoholics are often nicotine dependent,<sup>39,40</sup> heavy smokers,<sup>2,41</sup> and less successful at smoking cessation compared with non-alcoholics.<sup>41,42</sup>

Limitations of this study should be considered when interpreting our results. First is potentially decreased generalizability due to the high rates of nicotine and alcohol dependence among VET Registry respondents, perhaps associated with the ubiquitous exposure to tobacco and alcohol in the military. The prevalence of lifetime nicotine (47.8%) and alcohol dependence (35.2%) among VET Registry twins is greater than that reported from other epidemiologic samples. We are aware of only 1 other study of the prevalence of DSM-III-R nicotine dependence, which found that 20.0% of young adult (median age, 26 years) members of a Michigan health maintenance organization met lifetime DSM-III-R criteria for nicotine dependence.<sup>43</sup> The high prevalence of nicotine dependence in

our sample is consistent with the high rates of lifetime and current smoking in other veteran cohorts.<sup>44</sup>

The lifetime prevalence of 35.2% for alcohol dependence in the VET Registry is greater than the 20.1% reported by the National Comorbidity Study.<sup>45</sup> Reanalyses of National Comorbidity Study data, limited to men aged 34 through 55 years, yielded a lifetime prevalence of 21.1%. The prevalence of alcohol dependence in the VET Registry is close to the 39.2% lifetime prevalence of alcohol abuse and dependence reported by veterans in the National Vietnam Veterans Readjustment Study.<sup>46</sup> Our prevalence data may not generalize to nonveteran populations, including populations of persons who are too ill to be eligible for military service. However, the variances due to genetic influences on nicotine (60.3%) and alcohol dependence (55.1%) found in our analyses are similar to results from other twin panels.<sup>11,12,15,16</sup>

Second, our results should not be applied to women, since there may be a sex difference in the role of genes in the tobacco-alcohol relationship.<sup>25,47</sup> No nationally distributed female twin registry has been constructed for comparison with VET Registry data.

A basic assumption of twin studies is that MZ twin brothers are not treated differently than DZ twin brothers. This equal-environments assumption asserts that family environmental influences for MZ twins do not differ from those for DZ twins in their importance in the etiology of psychiatric disorders. Our analyses do not include a test of the equal-environments assumption; however, the equal-environments assumption has been shown to hold true for nicotine and alcohol dependence in other large twin populations.<sup>48</sup>

There are several strengths of our study. Our data were derived from a structured interview, and diagnoses of nicotine and alcohol dependence were derived according to standardized DSM-III-R criteria. Estimates of reliability of the nicotine and alcohol dependence diagnoses were good to excellent. Variance component estimates derived under best-fitting models were precise,

yielding narrow 95% CIs. Data were collected from a large, nationally distributed, nonclinical population, which increased generalizability.

The combined use of tobacco and alcohol is a serious public health concern. Tobacco use alone is responsible for more than 400 000 deaths per year in the United States.<sup>49</sup> Hurt and colleagues<sup>50</sup> found that tobacco-related conditions were associated with 50.9% of deaths among addicted persons and were twice those expected for the general population. Alcohol-related conditions were associated with 36.0% of deaths and were more than 4 times those expected in the general population. Their study emphasized the importance of the nicotine-alcohol relationship as a serious risk factor for increased mortality among substance-dependent persons. Our observation that these 2 dependencies have substantial common genetic causes suggests that lifetime nicotine and alcohol dependence comorbidity may be heritable, and for some nicotine-dependent alcoholics, smoking may be a trait coinherited with alcoholism. In such cases, smoking behavior would not be purely coincidental and may be construed as part of the larger risk for becoming dually addicted.

Although recent studies suggest smoking cessation is associated with continued abstinence from alcohol<sup>18,42</sup> and that smoking cessation efforts do not undermine alcoholism treatment,<sup>51</sup> the clinical research has recognized the lack of evidence for a mechanism driving these observations. Our results do not resolve the biological mechanisms of the nicotine-alcohol dependence relationship; however, possible mechanisms suggested by others include cross-tolerance to nicotine and alcohol,<sup>52,53</sup> common neural reward pathways,<sup>54,55</sup> and cued responding.<sup>39,56</sup> Our analyses identified a common genetic contribution to these dependencies that may underlie and influence some of the mechanisms of the nicotine-alcohol relationship.

## CONCLUSIONS

Others have already reported that smoking initiation and alcohol consumption have familial risk factors in common that are in part due to genetic influences, especially in young adults.<sup>57</sup> Our data suggest that among tobacco and alcohol users, genetic influences also contribute to the risk for dual dependence. Adolescents beginning to experiment with cigarettes and alcohol have little understanding of the powerful role that genes play in determining the risk for becoming an addicted smoker and alcohol dependent. It may be a prudent practice to stress the common genetic risk for dual addiction to nicotine and alcohol in efforts to prevent smoking and teenage drinking.

We have confirmed other reports of the high heritability of alcohol dependence, and we have identified a large genetic contribution to nicotine dependence. Most important, we have identified a genetic correlation between nicotine and alcohol dependence and genetic variance in risk for alcoholism that overlaps risk for nicotine dependence.

Accepted for publication February 23, 1999.

From the School of Public Health, St Louis University Health Sciences Center (Dr True and Mr Scherrer), Re-

search Service (Drs True, Xian, and Eisen and Mr Scherrer) and Medical Service (Dr Eisen), St Louis Veterans Affairs Medical Center, and the Departments of Psychiatry (Drs Madden, Bucholz, and Heath), Psychology and Genetics (Dr Heath), and Internal Medicine, Division of General Medical Sciences (Drs Xian and Eisen), Washington University School of Medicine, St Louis, Mo; the Department of Psychiatry, Harvard Medical School, Massachusetts Mental Health Center (Drs Lyons and Tsuang), the Department of Psychology, Boston University (Dr Lyons), and the Harvard Institute of Psychiatric Epidemiology and Genetics (Dr Tsuang), Boston, Mass; the Department of Veterans Affairs, Health Services Research and Development, Cooperative Studies in Health Services, Hines, Ill (Dr Goldberg); and the Epidemiology Program, School of Public Health, University of Illinois, Chicago (Dr Goldberg).

This study was supported by the Department of Veterans Affairs Health Services Research and Development Service and the Cooperative Studies Program, Washington, DC, Study 992. Partial support was provided by grants DA4604 and DA00272 from the National Institute on Drug Abuse (NIDA), Bethesda, Md; grants AA10339, AA11822, and AA07728 from the National Institute on Alcohol Abuse and Alcoholism (NIAAA), Great Lakes Veterans Affairs Health Services Research and Development Program, Ann Arbor, Mich, Locally Initiated Project 41-065; grants MH-37685 and MH-31302 from the National Institute of Mental Health (NIMH), Rockville, Md; and training grant DAO72261-01 from the NIDA (Washington University, St Louis). The National Comorbidity Study is funded by grants R01 MH/DA46376 and R01 MH49098 from the NIMH, and a supplement to R01 MH/DA46376 from the NIDA.

We acknowledge the work of the Midwest Center for Health Services and Policy Research and the Cooperative Studies Program, VET Registry (Director, W. G. Henderson, PhD; Epidemiologist, Dr Goldberg; Registry Programmer, K. Bukowski; and Coordinator, M. E. Vitek); the VET Registry advisory committee (A. G. Bearn, MD [past]; G. Chase, ScD [past]; T. Colton, ScD; W. E. Nance, MD, PhD; R. S. Paffenbarger, Jr, MD, DrPH; M. M. Weissman, PhD; and R. R. Williams, MD); Department of Veterans Affairs Chief Research and Development Officer John R. Feussner, MD, Health Services Research and Development Service (Deputy Director, S. Meehan, MBA, PhD), and Program Manager, C. Welch III, PhD; Ross Brownson, PhD, St Louis University School of Public Health, for his contribution to an earlier version of the manuscript; G. Carey, PhD, L. Eaves, PhD, DSc, and I. I. Gottesman, PhD, for their contributions to the early development of the VET Registry; the Department of Defense, National Personnel Records Center of the National Archives and Records Administration, the Internal Revenue Service, National Opinion Research Center, National Research Council of the National Academy of Sciences, and the Institute for Survey Research, Temple University, for invaluable support in the conduct of this study; and most important, the continued cooperation and participation of the members of the VET Registry. Without their contribution, this research would not have been possible.

Reprints: William R. True, PhD, MPH, Veterans Affairs Medical Center—St Louis Research Service, 151-JC, 915 N Grand Blvd, St Louis, MO 63106 (e-mail: true@slu.edu).

1. Maletzky BM, Klotter J. Smoking and alcoholism. *Am J Psychiatry*. 1974;131:445-447.
2. Cyr MG, Wartman SA. The effectiveness of routine screening questions in the detection of alcoholism. *JAMA*. 1988;259:51-54.
3. Kozlowski LT. Rehabilitating a genetic perspective in the study of tobacco and alcohol use. *Br J Addict*. 1991;86:517-520.
4. Shiffman S, Balabanis M. Associations between alcohol and tobacco. In: Fertig JB, Allen JP, eds. *Alcohol and Tobacco: From Basic Science to Clinical Practice*. Bethesda, Md: National Institute on Alcohol Abuse and Alcoholism; 1995:17-36. National Institute on Alcohol Abuse and Alcoholism research monograph 30. National Institutes of Health publication 95-3931.
5. Zacny JP. Behavioral aspects of alcohol-tobacco interactions. *Recent Dev Alcohol*. 1990;8:205-219.
6. Johnson KA, Jennison KM. The drinking-smoking syndrome and social context. *Int J Addict*. 1992;27:749-792.
7. Olsen J, Sabreo S, Fasting U. Interaction of alcohol and tobacco as risk factors in cancer of the laryngeal region. *J Epidemiol Community Health*. 1985;39:165-168.
8. Sees KL, Clark HW. When to begin smoking cessation in substance abusers. *J Subst Abuse Treat*. 1993;10:189-195.
9. Sobell LC, Sobell MB. Alcohol abuse and smoking: dual recoveries. *Alcohol Health Res World*. 1996;20:124-127.
10. Carmelli D, Swan GE, Robinette D, Fabsitz R. Genetic influence on smoking: a study of male twins. *N Engl J Med*. 1992;327:829-833.
11. Heath AC, Madden PAF. Genetic influences on smoking behavior. In: Turner JR, Cardon LR, Hewitt JK, eds. *Behavior Genetic Approaches in Behavioral Medicine*. New York, NY: Plenum Publishing Corp; 1995:45-66.
12. Heath AC, Martin NG. Genetic models for the natural history of smoking: evidence for a genetic influence on smoking persistence. *Addict Behav*. 1993;18:19-34.
13. True WR, Heath AC, Scherrer JF, Waterman B, Goldberg J, Lin N, Eisen SA, Lyons MJ, Tsuang MT. Genetic and environmental contributions to smoking. *Addiction*. 1997;92:1277-1287.
14. Heath AC. Genetic influences on alcoholism risk: a review of adoption and twin studies. *Alcohol Health Res World*. 1995;19:166-171.
15. Kendler KS, Prescott CA, Neale MC, Pedersen NL. Temperance board registration for alcohol abuse in a national sample of Swedish male twins, born 1902 to 1949. *Arch Gen Psychiatry*. 1997;54:178-184.
16. Heath AC, Bucholz KK, Madden PAF, Dinwiddie SH, Slutske WS, Bierut LJ, Statham DJ, Dunne MP, Whitfield JB, Martin NG. Genetic and environmental contributions to alcohol dependence risk in a national twin sample: consistency of findings in men and women. *Psychol Med*. 1997;27:1381-1396.
17. True WR, Heath AC, Bucholz K, Slutske W, Romeis JC, Scherrer JF, Lin N, Eisen SA, Goldberg J, Lyons MJ, Tsuang MT. Models of treatment seeking for alcoholism: the role of genes and environment. *Alcohol Clin Exp Res*. 1996;20:1577-1581.
18. Heath AC, Slutske WS, Madden PAF. Gender differences in the genetic contribution to alcoholism risk and to alcohol consumption patterns. In: Wilsnack RW, Wilsnack SC, eds. *Gender and Alcohol*. Rutgers, NJ: Rutgers University Press; 1997:114-149.
19. Battjes RJ. Smoking as an issue in alcohol and drug abuse treatment. *Addict Behav*. 1988;13:225-230.
20. Istvan J, Matarazzo JD. Tobacco, alcohol and caffeine use: a review of their interrelationships. *Psychol Bull*. 1984;95:301-326.
21. De Fiebre CM, Medhurst LJ, Collins AC. Nicotine response and nicotinic receptors in long-sleep and short-sleep mice. *Alcohol*. 1987;4:493-501.
22. De Fiebre CM, Collins AC. Classical genetic analyses of responses to nicotine and ethanol in crosses derived from long- and short-sleep mice. *J Pharmacol Exp Ther*. 1992;261:173-180.
23. Collins AC. Interactions of ethanol and nicotine at the receptor level. *Recent Dev Alcohol*. 1990;8:221-231.
24. Swan GE, Carmelli D, Cardon LR. The consumption of tobacco, alcohol, and coffee in Caucasian male twins: a multivariate analysis. *J Subst Abuse*. 1996;8:19-31.
25. Prescott CA, Kendler KS. Genetic and environmental influences on alcohol and tobacco dependence among women. In: Fertig JB, Allen JP, eds. *Alcohol and Tobacco: From Basic Science to Clinical Practice*. Bethesda, Md: National Institute on Alcohol Abuse and Alcoholism; 1995:59-87. National Institute on Alcohol Abuse and Alcoholism research monograph 30. National Institutes of Health publication 95-3931.
26. Eisen S, True W, Goldberg J, Henderson W, Robinette CD. The Vietnam Era Twin (VET) Registry: method of construction. *Acta Genet Med Gemellol (Roma)*. 1987;36:61-66.
27. Eisen S, Neuman R, Goldberg J, Rice J, True W. Determining zygosity in the Vietnam Era Twin Registry: an approach using questionnaires. *Clin Genet*. 1989;35:423-432.
28. Henderson WG, Eisen S, Goldberg J, True WR, Barnes JE, Vitek ME. The Vietnam Era Twin Registry: a resource for medical research. *Public Health Rep*. 1990;105:368-373.
29. Robins L, Helzer J, Cottler L, Goldring E. *NIMH Diagnostic Interview Schedule Version III Revised (DIS-III-R)*. St Louis, Mo: Dept of Psychiatry, Washington University Medical School; 1988.
30. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Third Edition Revised*. Washington, DC: American Psychiatric Association; 1987.
31. Fleiss JL. *Statistical Methods for Rates and Proportions*. 2nd ed. New York, NY: John Wiley & Sons Inc; 1981.
32. Slutske WS, True WR, Scherrer JF, Goldberg J, Bucholz KK, Heath AC, Henderson WG, Eisen SA, Lyons MJ, Tsuang MT. Long-term reliability and validity of alcoholism diagnoses and symptoms in a large national telephone interview survey. *Alcohol Clin Exp Res*. 1998;22:553-558.
33. Goldberg J, True W, Eisen S, Henderson W, Robinette CD. The Vietnam Era Twin (VET) Registry: ascertainment bias. *Acta Genet Med Gemellol (Roma)*. 1987;36:67-78.
34. Neale MC, Cardon LR. *Methodology for Genetic Studies of Twins and Families*. Norwell, Mass: Kluwer Academic Publishers; 1992.
35. Akaike H. Factor analysis and AIC. *Psychometrics*. 1987;52:317-333.
36. Neale MC. *Statistical Modeling With Mx*. Richmond: Dept of Human Genetics, Medical College of Virginia, Virginia Commonwealth University; 1991.
37. Joreskog KG, Sorbom D. PRELIS: a program for multivariate data screening and data summarization. In: *A Preprocessor for LISREL*. 2nd ed. Mooresville, Ind: Scientific Software; 1988.
38. Rosenbaum PR, Rubin DB. The central role of propensity score in observational studies for causal effects. *Biometrika*. 1983;70:41-55.
39. Abrams DB, Rohsenow DJ, Niaura RS, Pedraza M, Longabaugh R, Beattie MC, Binkoff JA, Noel NE, Monti PM. Smoking and treatment outcome for alcoholics: effects on coping skills, urge to drink, and drinking rates. *Behav Ther*. 1992;23:283-297.
40. Henningfield JE, Clayton R, Pollin W. Involvement of tobacco in alcoholism and illicit drug use. *Br J Addict*. 1990;85:279-292.
41. DiFranza JR, Guarrera MP. Alcoholism and smoking. *J Stud Alcohol*. 1990;51:130-135.
42. Bobo JK, Gilchrist LD, Schilling RF II, Noach B, Schinck SP. Cigarette smoking cessation attempts by recovering alcoholics. *Addict Behav*. 1987;12:209-215.
43. Breslau N, Kilbey MM, Andreski P. Nicotine dependence, major depression, and anxiety in young adults. *Arch Gen Psychiatry*. 1991;48:1069-1074.
44. Klevens RM, Giovino GA, Peddicord JP, Nelson DE, Mowery P, Grummer-Strawn L. The association between veteran status and cigarette-smoking behaviors. *Am J Prev Med*. 1995;11:245-250.
45. Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen HU, Kendler KS. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: results from the National Comorbidity Survey. *Arch Gen Psychiatry*. 1994;51:8-19.
46. Kulka RA, Schlenger WE, Fairbank JA, Hough RL, Jordan BK, Marmar CR, Weiss DS. *The National Vietnam Veterans Readjustment Study: Tables of Findings and Technical Appendices*. New York, NY: Brunner/Mazel Inc; 1990.
47. Madden PAF, Heath AC, Martin NG. Smoking and intoxication after alcohol challenge in women and men: genetic influences. *Alcohol Clin Exp Res*. 1997;21:1732-1741.
48. Kendler KS, Gardner CO Jr. Twin studies of adult psychiatric and substance dependence disorders: are they biased by differences in the environmental experiences of monozygotic and dizygotic twins in childhood and adolescence. *Psychol Med*. 1998;28:625-633.
49. United States Department of Health and Human Services. *Smoking and Health in the Americas: Executive Summary*. Atlanta, Ga: US Dept of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 1992. Dept of Health and Human Services publication (CDC) 92-8421.
50. Hurt RD, Offord KP, Croghan IT, Gomez-Dahl L, Kotke TE, Morse RM, Melton LJ. Mortality following inpatient addictions treatment: role of tobacco use in a community-based cohort. *JAMA*. 1996;275:1097-1103.
51. Hurt RD, Eberman KM, Croghan IT, Offord KP, Kavis LJ Jr, Morse RM, Palmen MA, Bruce BK. Nicotine dependence treatment during inpatient treatment for other addictions: a prospective intervention trial. *Alcohol Clin Exp Res*. 1994;18:867-872.
52. Burch JB, de Fiebre CM, Marks MJ, Collins AC. Chronic ethanol or nicotine treatment results in partial cross-tolerance between these agents. *Psychopharmacology*. 1988;95:452-458.
53. de Fiebre CM, Collins AC. A comparison of the development of tolerance to ethanol and cross-tolerance to nicotine after chronic ethanol treatment in long- and short-sleep mice. *J Pharmacol Exp Ther*. 1993;266:1398-1406.
54. Imperato A, Mulas A, Di Chiara G. Nicotine preferentially stimulates dopamine release in the limbic system of freely moving rats. *Eur J Pharmacol*. 1986;132:337-338.
55. Imperato A, Di Chiara G. Preferential stimulation of dopamine release in the nucleus accumbens of freely moving rats by ethanol. *J Pharmacol Exp Ther*. 1986;239:219-239.
56. Niaura RS, Rohsenow DJ, Binkoff JA, Monti PM, Pedraza M, Abrams DB. Relevance of cue reactivity to understanding alcohol and smoking relapse. *J Abnorm Psychol*. 1988;97:133-152.
57. Koopmans J. *The Genetics of Health-Related Behavior: A Study in Adolescent Twins and Their Parents* [dissertation]. Amsterdam, the Netherlands: Vrije Universiteit; 1997.