

Neurobehavioral Deficits at Adolescence in Children at Risk for Schizophrenia

The Jerusalem Infant Development Study

Sydney L. Hans, PhD; Joseph Marcus, MD; Keith H. Nuechterlein, PhD; Robert F. Asarnow, PhD; Benedict Styr, MD; Judith G. Auerbach, PhD

Background: The Jerusalem Infant Development Study is a prospective investigation comparing offspring of schizophrenic parents with offspring of parents who have no mental disorder or have nonschizophrenic mental disorders. During infancy and school age, a subgroup of offspring of schizophrenic parents showed global neurobehavioral deficits that were hypothesized to be indicators of vulnerability to schizophrenia. The purposes of the present investigation were to determine if neurobehavioral deficits were present in the offspring of schizophrenics at adolescence, to examine their stability over time, and to explore their relation to concurrent mental adjustment.

Methods: Sixty-five Israeli adolescents were assessed on a battery of neurologic and neuropsychological assessments. They were also administered psychiatric interviews from which best-estimate *DSM-III-R* diagnoses and scores of global adjustment were derived.

Results: Adolescents with poor neurobehavioral functioning were identified from composites of motor and cog-

nitive-attentional variables. A disproportionate number of offspring of schizophrenic parents (42%; 10/24), and especially male offspring of schizophrenic parents (73%; 8/11), showed poor neurobehavioral functioning relative to offspring of nonschizophrenic parents (22%; 9/41). Adolescent offspring of schizophrenics with poor neurobehavioral functioning had been poorly functioning at earlier ages and had poor psychiatric adjustment at adolescence. All 4 offspring of schizophrenics receiving schizophrenia spectrum diagnoses by adolescence showed a pattern of poor neurobehavioral functioning across developmental periods.

Conclusions: Results are consistent with the hypothesis that individuals at genetic risk for schizophrenia may display lifelong neurobehavioral signs that are indicators of vulnerability to schizophrenia and that are associated with psychiatric adjustment generally and schizophrenic spectrum disorder specifically.

Arch Gen Psychiatry. 1999;56:741-748

From the Department of Psychiatry, The University of Chicago, Chicago, Ill (Drs Hans and Marcus); the Neuropsychiatric Institute, the University of California at Los Angeles (Drs Nuechterlein and Asarnow); Eitanim Psychiatric Hospital, Jerusalem, Israel (Dr Styr); and the Department of Behavioral Sciences, Ben-Gurion University, Beer Sheva, Israel (Dr Auerbach).

SCHIZOPHRENIC patients show neurobehavioral deficits in a variety of motor, visual-motor, attentional, and cognitive tasks.¹⁻⁶ While some neurobehavioral signs may simply accompany symptoms of schizophrenic illness, family studies suggest that some may also be indicators of genetic vulnerability to schizophrenia. First-degree nonpsychotic relatives of schizophrenic patients are more likely than individuals with no schizophrenic relatives to have abnormalities in smooth-pursuit eye movements,⁷⁻⁹ grip-induced muscle tension,¹⁰ perceptual motor speed,^{11,12} sustained attention,^{13,14} and mental flexibility.¹⁵⁻¹⁷ Offspring of schizophrenics show deficits on the Continuous Performance Test,¹⁸⁻²¹ the Span of Apprehension Test,²² eye-tracking tasks,^{23,24} the visual backward masking procedure,²⁵ and fine motor coordination tasks.²⁶⁻²⁹ Anomalous pat-

terns of neurobehavioral development have been observed in offspring of schizophrenics as early as the first days of life.³⁰⁻³⁹ Debate continues about which specific neurobehavioral signs show the greatest sensitivity and specificity to schizophrenia and whether specific or general deficits are better indicators of vulnerability to schizophrenia.³⁵⁻⁴⁰

Although studies of first-degree relatives of schizophrenics suggest that neurobehavioral signs are genetic vulnerability indicators, this view would be strengthened by evidence that neurobehavioral deficits are stable over time and are associated with psychopathology, particularly with disorders in the schizophrenia spectrum. The longitudinal high-risk studies of offspring of schizophrenics are uniquely well suited to examine these issues.

To date, high-risk studies have reported stability of neurobehavioral signs from infancy to middle childhood²⁹ and

SUBJECTS AND METHODS

ORIGINAL STUDY SAMPLE: INFANCY AND SCHOOL AGE

The original JIDS sample was recruited from 1973 through 1977 by identifying pregnant women from Jerusalem's maternal and child care centers and mental health clinics.³² Research Diagnostic Criteria diagnoses³⁴ were made for all biological mothers and fathers based on a Hebrew-language version of the Current and Past Psychopathology Scales⁵⁵ and/or psychiatric and social work records for individuals who had received treatment. Based on these diagnoses, families were assigned to 3 groups: schizophrenic parent, parent with non-schizophrenic mental disorder, and parent with no mental illness.

A follow-up of the JIDS children was conducted when the children were a mean age of 10.3 years. Siblings of the original children who had not been part of the infant cohort were added to the school-age sample if they were between the ages of 8 and 13 years.

At the school-age follow-up, more than 80% of biological parents were interviewed using the Schedule for Affective Disorders and Schizophrenia–Lifetime Version.⁵⁶ Hospital and clinic records were assembled for all parents receiving mental health treatment. If parents were deceased or unavailable, spouses provided mental health history updates. Based on all available information, revised lifetime parent diagnoses were made using *DSM-III-R*³⁷ criteria and, where necessary, changes made in group assignment. All but 7 parents were available for research psychiatric interviewing at the infant and/or school-age assessments, and those 7 (3 schizophrenics, 1 mentally ill spouse of a schizophrenic, 2 nonschizophrenics with mental disorders) had sufficiently complete psychiatric or social worker records to make group assignment.

ADOLESCENT FOLLOW-UP

Beginning in 1992, a follow-up of the JIDS sample was conducted when the offspring were between the ages of 14 and

21 years (mean, 17.56; SD, 1.75). Sixty-five adolescents participated in the follow-up: 29 females, 36 males; 24 from the schizophrenic group, 25 from the other mental illness group, and 16 from the no mental illness group (**Table 1**). The parent diagnosis groups did not differ significantly in terms of mean age of adolescents (17.8 years for schizophrenia, 17.5 years for other mental illness, and 17.3 years for no mental illness) or proportion of males (46% [11/24] for schizophrenia, 56% [14/25] for other mental illness, and 69% [11/16] for no mental illness). Seven families (8 offspring) chose not to participate in the adolescent follow-up and 1 youth was excluded from participation because of a head injury that impacted neuropsychological functioning.

During the adolescent follow-up, parents were rediagnosed only if additional treatment records were available. These records required some revised diagnoses and eliminated some previously “questionable” diagnoses, but did not necessitate changes in group assignment.

Written informed consent was obtained from youth participants older than 18 years and from parents of those younger than 18 years.

Offspring Neurobehavioral Assessments

Adolescents were administered a neurobehavioral battery during two 2½-hour sessions at the Hebrew University, Jerusalem, Israel. Examiners were a psychiatrist (for neurologic examination) and master's degree-level psychologists (for all other instruments) trained to reliability and unaware of parents' diagnoses. The battery included assessments used in the National Institute of Mental Health Israeli High-Risk Study^{28,58} and the school-age JIDS, with additional tests found in other studies to be sensitive to schizophrenic diathesis.^{38,59,60} The battery included neurologic tasks scored on 4-point clinical scales (indicating no, mild, moderate, or severe impairment)⁶¹⁻⁶³ and averaged across multiple presentations of items as well as standard administered neuropsychological tests and procedures.^{58-60,64-81} For analyses in this article, 20 summary variables were selected from these instruments (**Table 2**).

from childhood through adolescence.^{27,41} They have also shown correlations between childhood measures of neurobehavioral functioning, particularly attentional functioning, and later indicators of general adjustment^{19,41-48} and schizophrenia spectrum disorders.⁴⁹⁻⁵¹ The anomalous neurodevelopmental pattern of infancy termed *pandysmaturation* has also been related to adult schizophrenia spectrum disorder.^{52,53}

The present article reports on the development of adolescents at risk for schizophrenia from the Jerusalem Infant Development Study (JIDS). Within this longitudinal sample, more global neurobehavioral signs were previously identified in infant and school-aged offspring of schizophrenic parents than in children whose parents had no mental disorder or nonschizophrenic disorders. Assuming that neurobehavioral signs are indicators of genetic vulnerability to schizophrenia, in this article we will examine hypotheses that (1) during adolescence neurobehavioral signs will be more preva-

lent in adolescent offspring of schizophrenics than in other young people, (2) adolescent offspring of schizophrenics with neurobehavioral signs will have shown signs at earlier developmental stages, and (3) neurobehavioral signs will be associated with psychiatric adjustment at adolescence, as well as with adolescent-onset schizophrenia spectrum disorder.

RESULTS

DIAGNOSTIC GROUP DIFFERENCES

Forty-two percent (n = 10) of the offspring of schizophrenic parents showed poor neurobehavioral functioning compared with 22% (n = 9; 5 from the other mental illness and 4 from the no mental illness group) of the offspring of nonschizophrenic parents.

The logistic regression model predicting poor neurobehavioral functioning was significant ($\chi^2_3 = 8.05$,

Offspring Psychiatric Assessment

During home visits, each adolescent and 1 parent were administered 2 diagnostic interviews: the Schedule for Affective Disorders and Schizophrenia for School-Age Children—Epidemiologic Version (K-SADS-E)⁸² and the Semi-Structured Kiddie Interview for Personality Syndromes (K-SKIPS),^{83,84} which, modeled after the Structured Clinical Interview for DSM-III-R Disorders,⁸⁵ was designed to assess personality syndromes falling within the schizophrenia spectrum. Parental information was used primarily to corroborate or add to the information provided by the adolescent. Where available, additional information from school records and mental health providers was obtained. All available information was used to make best-estimate DSM-III-R diagnoses⁸⁷ and Children's Global Assessment Scale ratings.⁸⁶ The interviewers included a senior psychiatrist and 2 other experienced clinicians who had received specialized training on administration of the K-SADS-E and K-SKIPS. Twenty randomly selected cases were blindly reviewed and independently diagnosed by 2 psychiatrists. Interrater agreement was high, although this random sample did not contain cases within the schizophrenia spectrum. All 7 cases diagnosed within the schizophrenia spectrum were independently reviewed by one of the senior investigators (J.M.), who agreed that all belonged within the spectrum. Disagreement on the specific schizophrenia spectrum diagnosis occurred for 1 case and was resolved by further discussion among all authors.

DATA ANALYSIS

In the infancy and school-age phases of this project, global measures of neurobehavioral functioning derived from 2-dimensional data structures were more effective than specific items or tests at discriminating a subgroup of poorly functioning offspring of schizophrenic parents from other children.²⁹ To produce a 2-dimensional data structure, the 20 adolescent neurobehavioral variables were subjected to a 2-factor principal components analysis with varimax rotation. Three of the variables (abnormal arm movements, tendon reflexes, and eye movements) had loadings less than 0.35. The principal components analysis was recomputed

without these 3 variables. The rotated components in this analysis explained 20.3% and 18.9% of the total variance. Variables with loadings of at least 0.35 on the first component were primarily cognitive-attentional; variables with loadings of at least 0.35 on the second component were primarily motoric (Table 2). The 17 variables were standardized, and cognitive-attentional and motor component scores were computed by averaging items with unit weights. Bender scores were entered into both the motor and cognitive-attentional averages. All variables were scaled before averaging so that positive scores indicated more problematic behavior.

To divide the sample into subjects with good and poor neurobehavioral functioning, the contours of an Epanechnikov kernel⁸⁷⁻⁸⁹ were superimposed on a plot of subjects' 2 principal components scores (**Figure**). This nonparametric procedure identifies the most concentrated region of data points in an empirical bivariate distribution. Because we wanted an empirical basis for identifying a region of normality, we used only the subjects with mentally healthy parents as a basis for computing the kernel and set program defaults so that approximately 68% of subjects fell within the kernel. Subjects from all 3 groups whose data points fell within the kernel, as well as any whose data points fell outside the boundary of the kernel but in the direction of better than average performance, were considered to be functioning well. All others outside the boundaries of the kernel were considered functioning poorly.

The relation of parent diagnosis to offspring neurobehavioral functioning was examined using contingency tables and logistic regressions predicting to the dichotomous neurobehavioral outcome from dummy variables for parental diagnosis, sex, and age. The stability of good and poor neurobehavioral outcomes over time was examined using contingency tables and κ statistics as a measure of association appropriate for dichotomous variables. The relation of neurobehavioral functioning to offspring psychopathology was examined using an analysis of variance with Children's Global Assessment Scale scores as the dependent variable and neurobehavioral functioning (2 levels) and parental diagnosis (3 levels) as independent variables.

All statistical tests are 2-tailed. For testing primary hypotheses, α levels were set at $P = .05$.

$P < .05$). Male children were nearly 4 times more likely to be poorly functioning (odds ratio, 3.99; 95% confidence interval, 1.12-14.20) than females. Trends suggested that offspring of schizophrenics were more than 3 times as likely to be poorly functioning as offspring of parents with other mental illness (odds ratio, 3.64; 95% confidence interval, 0.93-14.24; $P = .06$) and no mental illness (odds ratio, 3.15; 95% confidence interval, 0.70-14.15; $P = .13$). Since the odds ratios for the 2 comparison groups were similar, they were combined to provide greater statistical power and regressions recomputed. In this analysis, offspring of schizophrenic parents were more than 3 times as likely to be poorly functioning as offspring of nonschizophrenic parents (odds ratio, 3.43; 95% confidence interval, 1.03-11.42; $P < .05$).

Because inclusion of siblings in the sample violates the statistical assumption of independent sampling, we explored whether a few sibships were unduly contributing to the finding of poor functioning in offspring of

schizophrenics. This was not the case. In the 9 schizophrenic sibships, there were only 2 in which both siblings were poorly functioning and 2 in which both were well functioning.

Sex differences moderated the relation of parent diagnosis to poor neurobehavioral functioning. Seventy-three percent of the male offspring of schizophrenics (8 of 11) were poorly functioning, compared with 24% of the male offspring of nonschizophrenic parents (6 of 25; $\chi^2_1 = 7.63$, $P = .006$), 15% of the female offspring of schizophrenics (2 of 13; $\chi^2_1 = 8.06$, $P = .005$), and 19% of the female offspring of nonschizophrenic parents (3 of 16; $\chi^2_1 = 7.87$, $P = .005$).

Because in the JIDS school-age follow-up, pregnancy and birth complications (PBCs) were related to poor motoric functioning in the offspring of schizophrenic parents, we explored the role of PBCs for the 36 adolescent subjects with Research Obstetric Scale⁹⁰ data. Within the group of male offspring of schizophrenics, there was a

Table 1. Sample Size at Infancy, School Age, and Adolescent Assessments (Based on Parental Diagnoses Confirmed at School Age)

Parental Diagnosis	School Age					Adolescence			Total Adolescence Sample
	Infancy		School-aged Children Observed		School-aged Siblings	Adolescents Observed From Infancy	Adolescent Siblings Observed From School Age		
	Families	Infants	Families	From Infancy					
Schizophrenia	17	19	14	15	10	14*	15	9	24
Other mental illness	19	20	17	19	11	14	15	10	25
No mental illness	18	19	11	11	7	9	10	6	16
Total	54	58	42	45	28	37	40	25	65

*Eleven with schizophrenic mother, 2 with schizophrenic father, 1 with 2 schizophrenic parents.

Table 2. Summary of Measures Used in Principal Components Analysis With Loadings >0.35 Indicated

Type of Functioning	Variable	Cognitive- Attentional Component	Motor Functioning Component
Concentration ⁶⁶	Wechsler Intelligence Scale for Children Digit Span scaled score	-0.72	...
Perceptual motor speed ^{67,68}	Total seconds on Reitan Trail-Making Test (A and B)	0.71	...
Responses to signal ^{63,69}	Total errors on opposing action, alternating movement to command, go-no-go tasks	0.64	...
Flexibility of cognitive set ^{60,70}	Wisconsin Card Sorting Test, perseverative errors	0.61	...
Vigilance ⁷¹⁻⁷³	Continuous Performance Test (sensitivity score, A ¹ , during degraded stimulus condition)	-0.59	...
Span of apprehension ⁷⁴⁻⁷⁶	Span of Apprehension Test (number detected, 12-item array)	-0.56	...
Motor synchrony ⁵⁹	Average of synchrony scores from 20, 40, 80, 120, and 200 beats-per-minute trials	-0.45	...
Interference with cognitive set ^{77,78}	Stroop Color-Word Test interference score	-0.39	...
Auditory visual integration ⁷⁹	Birch-Belmont Test total incorrect trials	0.38	...
Fine motor coordination ⁶¹	Average of right- and left-hand scores from finger opposition, diadochokinesis, finger following, and match stick placement tasks	...	0.74
Associated movements ⁶¹	Average of overflow scores during forcible tongue protrusion, walking on tiptoe, walking on heels, and hopping on 1 leg	...	0.71
Serial coordinated movements ⁶²	Average scores from fist-clap-cut test, fist-ring test, and Ozeretski test	...	0.69
Mirror movements ⁶¹	Average of right- and left-hand overflow scores during finger opposition and diadochokinesis tasks	...	0.66
Visual-motor flexibility ^{58,64}	Mirror drawing—distance score	...	0.53
Fine motor speed ⁶⁵	No. of pegs placed on Purdue Pegboard (right and left hands)	...	-0.49
Abnormal postures ⁶¹	Average of ratings of abnormalities in rotation, extension, and abduction of raised arms and finger spooning	...	0.46
Visual-motor-perceptual maturity ^{80,81}	Bender Motor Gestalt Test with Pascal-Suttell scoring	0.39	0.44
Abnormal arm movements ⁶¹	Average of scores from distal choreiform movements, proximal choreiform movements, nonspecific gross movements, nonspecific small movements, and tremors
Tendon reflexes ⁶¹	Average of knee jerk and ankle jerk scores
Abnormal eye movements ⁶¹	Average of scores assessing convergence and fixation on objects, jerks during horizontal following and fixation, and strabismus

strong trend for PBCs to be related to poor motoric functioning (component 2) ($r = 0.64$, $P < .09$, 2-tailed; $n = 8$). The 3 male offspring of schizophrenics with more than 4 PBCs (numbers 45, 58, 61 in the Figure) all showed poor motoric functioning. Exploratory analyses suggested no relation between PBCs and poor motoric behavior in female offspring of schizophrenics or offspring without a schizophrenic parent. Pregnancy and birth complications were also unrelated to poor cognitive-attentional functioning.

STABILITY OF POOR FUNCTIONING FROM EARLIER AGES TO ADOLESCENCE

Offspring of schizophrenic parents showed greater stability in neurobehavioral functioning over age (Cohen $\kappa = 0.92$) than offspring of nonschizophrenic parents

($\kappa = 0.48$). More offspring of schizophrenics (42%; $n = 10$) were poorly functioning at both ages than offspring of mentally healthy parents (12.5%, $n = 2$; $\chi^2_1 = 3.89$, $P < .05$) or than offspring of parents with other mental disorders (12%, $n = 3$; $\chi^2_1 = 5.53$, $P < .05$) (Table 3). In contrast, stable good functioning across 2 age periods occurred for a high proportion of the offspring of schizophrenics (54%; $n = 13$) and the offspring of nonschizophrenics (71%; $n = 29$).

Similar analyses comparing neurobehavioral functioning at infancy and adolescence suggest considerably less stability. The offspring of schizophrenic parents showed only very modest stability from infancy (Cohen $\kappa = 0.32$) and the offspring of nonschizophrenic parents showed no stability (Cohen $\kappa = 0.08$).

Although many children did show shifts in functioning across age periods, fully half of the 40 children tracked from birth showed consistent levels of function-

ing across infancy, school age, and adolescence. Twelve offspring of nonschizophrenics (48%) showed consistently good functioning at all 3 developmental periods, and none showed consistently poor functioning. Four offspring of schizophrenics (27%) showed consistently good functioning, and 6 (40%) showed consistently poor functioning.

RELATION OF NEUROBEHAVIORAL FUNCTIONING TO GLOBAL PSYCHIATRIC ADJUSTMENT


Analysis of variance computed on the Children's Global Assessment Scale scores using parent diagnosis (3 levels) and neurobehavioral functioning at adolescence (2 levels) as the independent variables showed a significant effect for neurobehavioral functioning ($F_{1,56} = 8.61$, $P < .001$) and no effects for parent diagnosis ($F_{2,56} = 2.25$) or the interaction between parent diagnosis and neurobehavioral functioning ($F_{2,56} = 2.25$). Post hoc Fisher least significant difference tests indicated that the offspring of schizophrenic parents with poor neurobehavioral functioning had poorer mean Children's Global Assessment Scale scores ($n = 8$; mean, 49.13; SD, 13.67) than the offspring of schizophrenic parents with good neurobehavioral scores ($n = 13$; mean, 71.08; SD, 12.06; $P < .001$), the offspring of parents with other mental illness who had poor ($n = 5$; mean, 64.80; SD, 14.84; $P < .05$) or good ($n = 20$; mean, 67.80; SD, 13.85; $P < .001$) neurobehavioral scores, and the offspring of parents with no mental illness who had good neurobehavioral scores ($n = 12$; mean, 72.42; SD, 14.47; $P < .001$). None of the other groups differed from one another.

RELATION OF NEUROBEHAVIORAL FUNCTIONING TO SCHIZOPHRENIC SPECTRUM ILLNESS

Although the JIDS offspring were early in the period of risk for schizophrenic breakdown at the time of follow-up, 4 youth with schizophrenic parents received diagnoses in the schizophrenia spectrum: 1 schizophrenia, 1 schizotypal personality disorder, and 2 paranoid personality disorder. All 4 showed a stable pattern of poor neurobehavioral functioning at school age and adolescence. Three also showed poor infant functioning as reported by Marcus et al³²; the fourth was not in the sample during infancy. In an independent analysis of the infant behavior of these children, Fish et al⁵³ identified 2 of these infants as having probable pandysmaturations (developmental problems and growth retardation) and another as having a developmental pattern that was consistent with pandysmaturations but inconclusive because of missing physical growth data. An additional 3 male offspring without schizophrenic parents also received paranoid personality disorder diagnoses, but did not show consistently poor neurobehavioral functioning.

COMMENT

Three types of evidence from the adolescent follow-up of JIDS converge to suggest that neurobehavioral signs



Individual subjects plotted by cognitive-attentional and motor functioning components. Component 1 (on the y-axis) is the mean of concentration, perceptual motor speed, responses to signal, flexibility of cognitive set, vigilance, span of apprehension, motor synchrony, interference with cognitive set, auditory visual integration, and visual-motor-perceptual maturity standard scores. Component 2 (on the x-axis) is the mean of fine motor coordination, associated movements, serial coordinator movements, mirror movement, visual-motor flexibility, fine motor speed, postures, and visual-motor-perceptual maturity standard scores. An Epanechnikov kernel identifies the region of most concentrated data points for offspring of parents with no mental illness that is used to demarcate the offspring with good and poor neurobehavioral functioning. Identification numbers are noted for offsprings of schizophrenics only and are consistent with those reported in the school-age follow-up.

may be markers of vulnerability to schizophrenia: (1) global neurobehavioral signs were more prevalent in offspring of schizophrenic parents than in other young people; (2) neurobehavioral signs were stable over development for a subgroup of offspring of schizophrenic parents, but not other young people; (3) for offspring of schizophrenic parents, neurobehavioral signs were associated with psychopathology and adolescent schizophrenia spectrum disorders.

When the JIDS sample was assessed during the first year of life, more offspring of schizophrenics showed problems in motor and sensorimotor behavior than offspring of parents with no mental disorder or nonschizophrenic mental disorder. During school age (7-13 years) when the JIDS children and their similarly aged siblings were assessed, again a subgroup of offspring of schizophrenics showed poor motor and cognitive functioning. The present article, reporting on the adolescent follow-up of the original JIDS sample and their siblings, again identifies a subgroup of children of schizophrenics with poor neurobehavioral functioning. Overall, 42% ($n = 10$) of the offspring of schizophrenics were poorly functioning at adolescence compared with only 22% ($n = 9$) of the offspring of parents with other mental disorders or no mental disorder. During the school-age and infancy assessments, the proportion of poorly functioning offspring of schizophrenics had been 44% and 68%, respectively, compared with 20% and 23% for the children without schizophrenic parents. After controlling for age and sex in logistic regression analyses, offspring of schizo-

Table 3. Cross Tabulation of Well- and Poor-Functioning Subgroups at School Age and Adolescence Grouped by Parent Diagnosis

		Offspring of					
		Schizophrenic Parents		Parents With Other Mental Illness		Parents With No Mental Illness	
Adolescence	Poor	0	10	2	3	2	2
	Good	13	1	17	3	12	0
		Good	Poor	Good	Poor	Good	Poor
School Age							

phrenics were more than 3 times as likely as other young people to have neurobehavioral signs.

Although we had not originally hypothesized sex differences, adolescents with neurobehavioral signs, particularly those with motoric signs, were predominantly male offspring of schizophrenics. The incidence of poor neurobehavioral functioning was 73% in the males with a schizophrenic parent compared with less than 25% in the female offspring of schizophrenics and male and female offspring of nonschizophrenics. Although the incidence of schizophrenia is similar between males and females, sex differences abound in the study of schizophrenia.⁹¹⁻⁹³ Castle and Murray⁹⁴ have even hypothesized that male schizophrenics typically suffer from a different type of schizophrenia than females, with schizophrenia in males characterized by more neurobehavioral and neuroanatomical signs and more closely associated with PBCs than genetic vulnerability. Data from the present study are consistent with this hypothesis in that at-risk males show poorer neurodevelopmental course prior to illness onset and their poor functioning, at least in the motoric domain, may be associated with a history of PBCs in the small number of subjects for whom PBC data were available. However, data from the present study also suggest that the role of genetics cannot be ignored in considering the course of neurodevelopmental problems in schizophrenia for males, since neurobehavioral signs were only increased in males with PBCs and a schizophrenic parent. We believe the most parsimonious explanation of the data is that males at genetic risk are especially vulnerable to the effects of prenatal hazards.

The global neurobehavioral measures used in this study showed stability across ages. This stability was strong between school age and adolescence, although only modest between infancy and adolescence. As reported in other high-risk studies,^{41,95} the stability was greater for the offspring of schizophrenics than for the children with nonschizophrenic parents and most notably was anchored by a poorly functioning subgroup of offspring of schizophrenics. Fully 42% of the offspring of schizophrenics had been poorly functioning at both school age and adolescence, compared with only 12% of the offspring of nonschizophrenics. Six of the 40 children in the sample observed from birth through adolescence have shown consistently poor neurobehavioral functioning over time. All 6 of these children were the offspring of schizophrenics.

An association was found between the neurobehavioral measures and psychopathology. Other high-risk stud-

ies have found linkages between neurobehavioral signs (especially attentional deviance) and concurrent global psychopathology^{19,41} and later schizophrenia spectrum disorders.^{49,50,52,96} The present study determined that the adolescent offspring of schizophrenics showing neurobehavioral signs were at risk for poorer global psychiatric adjustment as well as specific schizophrenia spectrum disorders.

At the time JIDS began, virtually no empirical attention had been given to biobehavioral measures indicating enhanced vulnerability to schizophrenia. Fish⁹⁶ had described a pattern of anomalous early development she termed "pandysmaturation," which she proposed was "a 'marker' in infancy for the inherited neurointegrative defect in schizophrenia." We interpreted the JIDS infancy findings as suggesting "some kind of genetically determined vulnerability to schizophrenia."³² Today, hosts of studies on first-degree relatives of schizophrenics have searched for behavioral, neurochemical, neuroanatomical, and genetic vulnerability indicators of schizophrenia,^{97,98} and a number of theoretical conceptions of vulnerability indicators have evolved from these (see further Kremen et al⁹⁹). The JIDS adolescent follow-up data, in showing associations between neurobehavioral signs and genetic risk for schizophrenia, stability in neurobehavioral signs over time, and correlation between these signs and psychopathology, continue to support the hypothesis that global neurodevelopmental deficits may be premorbid indicators of genetic vulnerability to schizophrenia. Because the measures we have used at adolescence, school age, and infancy are global, we do not claim that they are specific to schizophrenia. In fact, they occur, although with lower incidence and little stability, in offspring of parents with no mental illness and offspring of parents with other, nonschizophrenic mental disorders.

Researchers studying the brain basis of schizophrenia are suggesting that the types of neuroanatomical anomalies associated with schizophrenia do not result from degenerative processes, but from developmental processes that may begin early in life.^{100,101} The JIDS longitudinal data, combined with the pioneering work of Fish,⁹⁶ the recent studies of Walker et al,^{102,103} and an accumulation of other infancy data from the high-risk field,³³ add further support for the view that schizophrenia is a neurodevelopmental disorder with origins very early in life.

The primary limitation of the present study, and most high-risk studies, is its sample size. Results need to be

interpreted with caution and in relation to findings from other studies. A second limitation is that, because of the long-term longitudinal design, measurement techniques are not always contemporary by the time of follow-up. Even in the most recent follow-up, limitations of budget and technology available in Israel did not allow for state-of-the-art measures such as neuroimaging or brain potentials. The greatest limitation of the present report is that none of the subjects in the sample have yet passed through the period of risk for schizophrenia and conclusions about actual schizophrenic illness must be treated cautiously.

Accepted for publication March 30, 1999.

Collection of adolescent data and preparation of this article were supported by grant R01 MH45208 from the National Institute of Mental Health, Rockville, Md. Collection of infancy data was supported by grants from the US-Israel Binational Science Foundation (grant 598), Jerusalem, Israel; the Chief Scientists' Office of the Israel Ministry of Health, Jerusalem; the Olivetti Foundation, Boston, Mass; the Center for the Study of Human Sciences of the Hebrew University, Jerusalem; the Department of Psychiatry, the University of Chicago, Chicago, Ill; Forest Hospital Foundation, Des Plaines, Ill, and Harry Jacobs, Cleveland, Ohio. Collection and analysis of school-age follow-up data were supported by the W. T. Grant Foundation, New York, NY; the Scottish Rite Schizophrenia Research Program, Lexington, Mass; the Sturman Center of Human Development, Hebrew University; and by a gift from Sarah Cowan, Cleveland.

Data at the adolescent follow-up were collected by Miriam Barasch, MSW, Nomi Ban, MA, Batya Aloni, Slava Feinstein, MD, Sharon Arnon, MA, Nurit Kaveh, MA, Nili Mor, MA, and Gil Amihai, MA; the database was managed by Linda Henson. Statistical consultation was provided by Leland Wilkinson, PhD, Department of Statistics, Northwestern University, Evanston, Ill.

Reprints: Sydney L. Hans, PhD, Department of Psychiatry, MC3077, The University of Chicago, 5841 S Maryland Ave, Chicago, IL 60637.

REFERENCES

- Cox SM, Ludwig AM. Neurological soft signs and psychopathology: incidence in diagnostic groups. *Can J Psychiatry*. 1979;24:668-673.
- Manschreck TC, Ames D. Neurologic features and psychopathology. *Biol Psychiatry*. 1984;19:703-719.
- Heinreichs DW, Buchanan RW. The significance and meaning of neurological signs in schizophrenia. *Am J Psychiatry*. 1988;145:11-18.
- Nasrallah HA, Weinberger DR, eds. *The Handbook of Schizophrenia: The Neurology of Schizophrenia*. Vol 1. Amsterdam, the Netherlands: Elsevier Science Publishers; 1986.
- Steinhauer SR, Gruzeller JH, Zubin J, eds. *Handbook of Schizophrenia: Neuropsychology, Psychophysiology and Information Processing*. Vol 5. Amsterdam, the Netherlands: Elsevier Science Publishers; 1991.
- Seidman LJ, Cassens GP, Kremen WS, Pepple JR. Neuropsychology of schizophrenia. In: White RF, ed. *Clinical Syndromes in Adult Neuropsychology: The Practitioner's Handbook*. New York, NY: Elsevier Science Inc; 1992:381-449.
- Clementz BA, Sweeney JA, Hirt M, Haas G. Pursuit gain and saccadic intrusions in first-degree relatives of probands with schizophrenia. *J Abnorm Psychol*. 1992;99:327-335.
- Holzman PS, Proctor LR, Levy DL, Yasillo NJ, Meltzer HY, Hurst SW. Eye-tracking dysfunctions in schizophrenic patients and their relatives. *Arch Gen Psychiatry*. 1974;31:143-151.
- Keefe RS, Silverman JM, Mohs RC, Siever LJ, Harvey PD, Friedman L, Roitman SEL, DuPre RL, Smith CJ, Schmeidler J, Davis KL. Eye tracking, attention, and schizotypal symptoms in nonpsychotic relatives of patients with schizophrenia. *Arch Gen Psychiatry*. 1997;54:169-176.
- Rosen AJ, Lockhart JJ, Gants ES. Maintenance of grip-induced muscle tension: a behavioral marker of schizophrenia. *J Abnorm Psychol*. 1991;100:583-593.
- Keefe RSE, Silverman JM, Lees Roitman SE, Harvey PD, Duncan MA, Alray D, Siever LJ, Davis KL, Mohs RC. Performance of nonpsychotic relatives of schizophrenic patients on cognitive tests. *Psychiatry Res*. 1994;53:1-12.
- Pogue-Geile MF, Garrett AH, Brunke JJ, Hall JK. Neuropsychological impairments are increased in siblings of schizophrenic patients [abstract]. *Schizophr Res*. 1991;4:390.
- Wood RL, Cook M. Attentional deficit in the siblings of schizophrenics. *Psychol Med*. 1979;9:465-467.
- Steinhauer SR, Zubin J, Condray R, Shaw DB, Peters JL, Van Kammen DP. Electrophysiological and behavioral signs of attentional disturbance in schizophrenics and their siblings. In: Tamminga CA, Schulz SC, eds. *Schizophrenia Research: Advances in Neuropsychiatry and Psychopharmacology*. Vol 1. New York, NY: Raven Press; 1991:169-178.
- Faraone SV, Seidman LJ, Kremen WS, Pepple JR, Lyons MJ, Tsuang MT. Neuropsychological functioning among the nonpsychotic relatives of schizophrenic patients: a diagnostic efficiency analysis. *J Abnorm Psychol*. 1995;104:286-304.
- Pogue-Geile MF. Siblings of schizophrenic probands: presence of neuropsychological impairments [abstract]. *Schizophr Res*. 1990;3:62.
- Franke P, Maier W, Hain C, Klingler T. Wisconsin Card Sorting Test: an indicator of vulnerability to schizophrenia? *Schizophr Res*. 1992;6:243-249.
- Nuechterlein KH. Signal detection in vigilance tasks and behavioral attributes among offspring of schizophrenic mothers and among hyperactive children. *J Abnorm Psychol*. 1983;92:4-28.
- Cornblatt B, Erlenmeyer-Kimling L. Early attentional predictors of adolescent behavioral disturbances in children at risk for schizophrenia. In: Watt N, Anthony EJ, Wynn L, Rofe J, eds. *Children at Risk for Schizophrenia: A Longitudinal Perspective*. New York, NY: Cambridge University Press; 1984:198-211.
- Rutschmann J, Cornblatt B, Erlenmeyer-Kimling L. Sustained attention in children at risk for schizophrenia. *Arch Gen Psychiatry*. 1977;34:571-575.
- Rutschmann J, Cornblatt B, Erlenmeyer-Kimling L. Sustained attention in children at risk for schizophrenia: findings with two visual continuous performance tests in a new sample. *J Abnorm Child Psychol*. 1986;14:365-385.
- Asarnow RF, Steffy RA, MacCrimmon DJ, Cleghorn JM. An attentional assessment of foster children at risk for schizophrenia. *J Abnorm Psychol*. 1977;86:267-274.
- Schreiber H, Stolz-Born G, Born J, Rothmeier J, Rothenberger A, Jurgens R, Becker W, Kornhuber HH. Visually-guided saccadic eye movements in adolescents at genetic risk for schizophrenia. *Schizophr Res*. 1997;25:97-109.
- Rosenberg DR, Sweeney JA, Squires-Wheeler E, Keshavan MS, Cornblatt BA, Erlenmeyer-Kimling L. Eye-tracking dysfunction in offspring from the New York High-Risk Project: diagnostic specificity and the role of attention. *Psychiatry Res*. 1997;66:121-130.
- Green MF, Nuechterlein KH, Breitmeyer B. Backward masking performance in unaffected siblings of schizophrenic patients: evidence for a vulnerability indicator. *Arch Gen Psychiatry*. 1997;54:465-472.
- Erlenmeyer-Kimling L, Cornblatt B, Friedman D, Marcuse Y, Rutschmann J, Simmens S. Neurological, electrophysiological and attentional deviations in children at risk for schizophrenia. In: Nasrallah HA, Henn FA, eds. *Schizophrenia as a Brain Disease*. New York, NY: Oxford University Press Inc; 1982:61-98.
- Marcus J, Hans SL, Lewow E, Wilkinson L, Burack CM. Neurological findings in high-risk children: childhood assessment and 5-year followup. *Schizophr Bull*. 1985;11:85-100.
- Marcus J, Hans SL, Mednick S, Schulsinger F, Michelsen N. Neurological dysfunctioning in offspring of schizophrenics in Israel and Denmark: a replication analysis. *Arch Gen Psychiatry*. 1985;42:753-761.
- Marcus J, Hans SL, Auerbach JG, Auerbach AG. Children at risk for schizophrenia: the Jerusalem Infant Development Study, II: neurobehavioral deficits at school age. *Arch Gen Psychiatry*. 1993;50:797-809.
- Fish B, Alpert M. Abnormal states of consciousness and muscle tone in infants born to schizophrenic mothers. *Am J Psychiatry*. 1962;119:439-445.
- Fish B, Dixon WJ. Vestibular hyporeactivity in infants at risk for schizophrenia: its association with critical developmental disorders. *Arch Gen Psychiatry*. 1978;35:963-971.
- Marcus J, Auerbach J, Wilkinson L, Burack CM. Infants at risk for schizophrenia: the Jerusalem Infant Development Study. *Arch Gen Psychiatry*. 1981;38:703-713.
- Hans SL, Marcus J. Neurobehavioral development of infants at risk for schizophrenia: a review. In: Walker EF, ed. *Schizophrenia: A Life-Course Developmental Perspective*. New York, NY: Academic Press Inc; 1991:33-57.
- Schubert EW, Blennow G, McNeil TF. Wakefulness and arousal in neonates born to women with schizophrenia: diminished arousal and its association with neurological deviations. *Schizophr Res*. 1996;22:49-59.
- Blanchard JJ, Neale JM. The neuropsychological signature of schizophrenia: generalized or differential deficit? *Am J Psychiatry*. 1994;151:40-48.
- Braff DL, Heaton R, Kuck J, Cullum M, Moranville J, Grant I, Zisook S. The generalized pattern of neuropsychological deficits in outpatients with chronic schizophrenia with heterogeneous Wisconsin Card Sorting Test results. *Arch Gen Psychiatry*. 1991;48:891-898.
- Flashman LA, Flaum M, Gupta S, Andreasen NC. Soft signs and neuropsychological performance in schizophrenia. *Am J Psychiatry*. 1996;153:526-532.

38. Asarnow RF, Granholm E, Sherman T. Span of apprehension in schizophrenia. In: Steinhauer SR, Gruzeliel JH, Zubin J, eds. *Handbook of Schizophrenia: Neuropsychology, Psychophysiology and Information Processing*. Vol 5. Amsterdam, the Netherlands: Elsevier Science Publishers; 1991:335-370.
39. Nuechterlein K. Vigilance in schizophrenia and related disorders. In: Steinhauer SR, Gruzeliel JH, Zubin J, eds. *Handbook of Schizophrenia: Neuropsychology, Psychophysiology and Information Processing*. Vol 5. Amsterdam, the Netherlands: Elsevier Science Publishers; 1991:397-433.
40. Saykin AJ, Gur RC, Gur RE, Mozley PD, Mozley LH, Resnick SM, Kester B, Stafinjak P. Neuropsychological function in schizophrenia: selective impairment in memory and learning. *Arch Gen Psychiatry*. 1991;48:618-624.
41. Winters L, Cornblatt B, Erlenmeyer-Kimling L. The prediction of psychiatric disorders in late adolescence. In: Walker E, ed. *Schizophrenia: A Life-Course Developmental Perspective*. New York, NY: Academic Press Inc; 1991:123-137.
42. Erlenmeyer-Kimling LA, Cornblatt BA. The New York High-Risk Project: a follow-up report. *Schizophr Bull*. 1987;13:451-461.
43. Cornblatt B, Marcuse Y. Children at high risk for schizophrenia: predictions from childhood to adolescence. In: Erlenmeyer-Kimling L, Miller N, eds. *Life-span Research on the Prediction of Psychopathology*. Hillsdale, NJ: Lawrence Erlbaum Associates; 1986:81-100.
44. Erlenmeyer-Kimling L, Cornblatt B, Golden R. Early indicators of vulnerability to schizophrenia in children at high genetic risk. In: Guze SB, Earsl FJ, Barrett JE, eds. *Childhood Psychopathology and Development*. New York, NY: Raven Press; 1983:247-261.
45. Cornblatt B, Erlenmeyer-Kimling L. Global attentional deviance as a marker of risk for schizophrenia: specificity and predictive validity. *J Abnorm Psychol*. 1985;94:470-486.
46. Cornblatt BA, Lenzenweger M, Dworkin R, Erlenmeyer-Kimling L. Positive and negative schizophrenic symptoms, attention, and information processing. *Schizophr Bull*. 1985;11:397-407.
47. Dworkin RH, Cornblatt BA, Friedmann R, Kaplansky LM, Lewis JA, Rinaldi A, Shiliday C, Erlenmeyer-Kimling L. Childhood precursors of affective vs social deficits in adolescents at risk for schizophrenia. *Schizophr Bull*. 1993;19:563-577.
48. Erlenmeyer-Kimling L, Cornblatt BA, Rock D, Roberts S, Bell M, West A. The New York High-Risk Project: anhedonia, attentional deviance, and psychopathology. *Schizophr Bull*. 1993;19:141-153.
49. Marcus J, Hans SL, Nagler S, Auerbach JG, Mirsky AF, Aubrey A. A review of the NIMH Israeli Kibbutz-City Study and the Jerusalem Infant Development Study. *Schizophr Bull*. 1987;13:425-437.
50. Hans SL, Marcus J. A process model for the development of schizophrenia. *Psychiatry*. 1987;50:361-370.
51. Mirsky AF, Ingraham LJ, Kugelmass S. Neuropsychological assessment of attention and its pathology in the Israeli cohort. *Schizophr Bull*. 1995;21:193-204.
52. Fish B. Infant predictors of the longitudinal course of schizophrenic development. *Schizophr Bull*. 1987;13:395-409.
53. Fish B, Marcus J, Hans SL, Auerbach JG, Perdue S. Infants at risk for schizophrenia: sequelae of a genetic neurointegrative defect: a review and replication analysis of pandsymptomatology in the Jerusalem Infant Development Study. *Arch Gen Psychiatry*. 1992;49:221-235.
54. Spitzer RL, Endicott J, Robins E. *Research Diagnostic Criteria (RDC) for a Selected Group of Functional Disorders*. 2nd ed. New York: Biometrics Research, New York State Psychiatric Institute; 1975.
55. Endicott J, Spitzer RL. Current and Past Psychopathology Scales (CAPPS): rationale, reliability, and validity. *Arch Gen Psychiatry*. 1972;27:678-687.
56. Spitzer RL, Endicott J. *Schedule for Affective Disorders and Schizophrenia—Lifetime Version*. New York: New York State Psychiatric Institute; 1975.
57. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Revised Third Edition*. Washington, DC: American Psychiatric Association; 1987.
58. Lifshitz M, Kugelmass S, Karov M. Perceptual-motor and memory performance of high-risk children. *Schizophr Bull*. 1985;11:74-84.
59. Manschreck TC, Maher BA, Rucklos MW, Vereen DR, Ader DN. Deficient motor synchrony in schizophrenia. *J Abnorm Psychol*. 1981;90:321-328.
60. Grant DA, Berg EA. A behavioral analysis of degree of reinforcement and ease of shifting to new responses in a Weigl-type card-sorting problem. *J Exp Psychol*. 1948;38:404-411.
61. Touwen BC, Prechtl HFR. The neurological examination of the child with minor nervous dysfunction. In: *Clinics in Developmental Medicine No. 8*. London, England: William Heinemann Medical Books Ltd; 1970.
62. Luria AR. *Higher Cortical Functions in Man*. New York, NY: Basic Books; 1966.
63. Luria AR. *The Working Brain: An Introduction to Neuropsychology*. New York, NY: Basic Books; 1973.
64. Wechsler D, Hartogs R. The clinical measurement of anxiety: an experimental approach. *Psychiatr Q*. 1945;19:618-635.
65. Purdue Research Foundation. *Examiner's Manual for the Purdue Pegboard*. Chicago, Ill: Science Research Associates; 1948.
66. Wechsler D. *Manual for the Wechsler Intelligence Scale for Children—Revised*. New York, NY: The Psychological Corp; 1974.
67. Reitan RM. Investigation of the validity of Halstead's measures of biological intelligence. *Arch Neurol Psychiatry*. 1955;73:28-35.
68. Reitan RM, Tarshes EL. Differential effects of lateralized brain lesions on the Trail Making Test. *J Nerv Ment Dis*. 1959;129:257-262.
69. Benson DF, Stuss DT. Motor abilities after frontal leukotomy. *Neurology*. 1982;32:1353-1357.
70. Loong J. *Wisconsin Card Sorting Test—IBM Version*. San Luis Obispo, Calif: Wang Neuropsychological Laboratories; 1991.
71. Rosvold HE, Mirsky AF, Sarason I, Bransome ED Jr, Beck LH. A continuous performance test of brain damage. *J Consult Psychol*. 1956;20:343-350.
72. Nuechterlein KH, Parasuraman R, Jiang Q. Visual sustained attention: image degradation produces rapid sensitivity decrement over time. *Science*. 1983;220:327-329.
73. Nuechterlein KH, Asarnow RF. *Continuous Performance Test (CPT) Program for IBM-Compatible Microcomputers, Version 4, for Degraded Stimulus CPT*. Los Angeles, Calif: Nuechterlein and Asarnow; 1990.
74. Estes WK, Taylor HA. Visual detection in relation to display size and redundancy of critical elements. *Percept Psychophys*. 1966;1:9-16.
75. Asarnow RF, MacCrimmon DF. Span of apprehension deficits during the post-psychotic stages of schizophrenia: a replication and extension. *Arch Gen Psychiatry*. 1981;38:1006-1011.
76. Asarnow RF, Nuechterlein KH. *Span of Apprehension Program for IBM-Compatible Microcomputers, Version 4*. Los Angeles, Calif: Asarnow and Nuechterlein; 1991.
77. Stroop JR. Studies of interference in serial verbal reactions. *J Exp Psychol*. 1935;18:643-662.
78. Ingraham LJ, Chard F, Wood M, Mirsky AF. A Hebrew language version of the Stroop Test. *Percept Mot Skills*. 1988;67:187-192.
79. Birch HG, Belmont L. Auditory-visual integration in brain damaged and normal children. *Dev Med Child Neurol*. 1965;7:135-144.
80. Bender L. *A Visual Motor Test and Its Clinical Use*. New York, NY: American Orthopsychiatric Association; 1938. American Orthopsychiatric Association Research Monograph No. 3.
81. Pascall GR, Suttell BJ. *The Bender-Gestalt Test: Quantification and Validity for Adults*. New York, NY: Grune & Stratton Inc; 1951.
82. Orvaschel H, Puig-Antich J. *Schedule for Affective Disorders and Schizophrenia for School-Age Children—Epidemiologic Version*. Ft Lauderdale, Fla: Nova University, Center for Psychological Study; 1987.
83. Asarnow JR, Talovic S. *Semi-Structured Kiddie Interview for Personality Syndromes (K-SKIPS)*. Los Angeles: University of California, Los Angeles, Dept of Psychiatry; 1986.
84. Asarnow JR, Tompson MC, Goldstein MJ. Childhood-onset schizophrenia. *Schizophr Bull*. 1994;20:599-617.
85. Spitzer RL, Williams JB. *Structured Clinical Interview for DSM-III-R, Personality Disorders*. New York: New York State Psychiatric Institute; 1986.
86. Shaffer D, Gould MS, Brasic J, Ambrosini P, Fisher P, Bird H, Aluwahlia S. A Children's Global Assessment Scale (CGAS). *Arch Gen Psychiatry*. 1983;40:1228-1231.
87. Silverman BW. *Density Estimation for Statistics and Data Analysis*. London, England: Chapman & Hall; 1986.
88. Scott DW. *Multivariate Density Estimation: Theory, Practice, and Visualization*. New York, NY: John Wiley & Sons; 1992.
89. Wilkinson L. *SYSTAT: The System for Statistics*. Evanston, Ill: SYSTAT Inc; 1990.
90. Zax M, Sameroff AJ, Baglioni HM. Birth outcomes in the offspring of mentally disordered women. *Am J Orthopsychiatry*. 1977;47:218-230.
91. Lewine RRJ. Gender and schizophrenia. In: Tsuang MT, Simpson JC, eds. *Handbook of Schizophrenia: Nosology, Epidemiology and Genetics of Schizophrenia*. Vol 3. Amsterdam, the Netherlands: Elsevier Science Publishers; 1988:379-397.
92. Lewis S. Sex and schizophrenia: vive la difference. *Br J Psychiatry*. 1992;161:445-450.
93. Goldstein JM, Tsuang MT. Gender and schizophrenia: an introduction and synthesis of findings. *Schizophr Bull*. 1990;16:179-183.
94. Castle DJ, Murray RM. The neurodevelopmental basis of sex differences in schizophrenia. *Psychol Med*. 1991;21:565-575.
95. Marcus J, Hans S, Byhouwer B, Norem J. Relationships among neurological functioning, intelligence quotients and physical anomalies. *Schizophr Bull*. 1985;11:101-106.
96. Fish B. Neurobiologic antecedents of schizophrenia in children. *Arch Gen Psychiatry*. 1977;34:1297-1313.
97. Kremen WS, Seidman LJ, Peppel JR, Lyons MJ, Tsuang MT, Faraone SV. Neuropsychological risk indicators for schizophrenia: a review of family studies. *Schizophr Bull*. 1994;20:103-119.
98. Moldin SO, Erlenmeyer-Kimling L. Measuring liability to schizophrenia: progress report 1994: editors' introduction. *Schizophr Bull*. 1994;20:25-29.
99. Kremen WS, Tsuang MT, Faraone SV, Lyons MJ. Using vulnerability indicators to compare conceptual models of genetic heterogeneity in schizophrenia. *J Nerv Ment Dis*. 1992;180:141-152.
100. Weinberger DR. Implications of normal brain development for the pathogenesis of schizophrenia. *Arch Gen Psychiatry*. 1987;44:660-669.
101. Murray RM, Lewis SW. Is schizophrenia a neurodevelopmental disorder? *BMJ*. 1987;295:681-682.
102. Walker EF, Savoie T, Davis D. Neuromotor precursors of schizophrenia. *Schizophr Bull*. 1994;20:441-451.
103. Walker EF, Grimes KE, Davis DM. Childhood precursors of schizophrenia: facial expressions of emotion. *Am J Psychiatry*. 1993;150:1654-1660.