

Prevalence and Persistence of Psychiatric Disorders in Youth After Detention

A Prospective Longitudinal Study

Linda A. Teplin, PhD; Leah J. Welty, PhD; Karen M. Abram, PhD; Mina K. Dulcan, MD; Jason J. Washburn, PhD

Context: Psychiatric disorders are prevalent among incarcerated juveniles. Most juveniles eventually return to their communities, where they become the responsibility of the community mental health system. However, no large-scale study has examined psychiatric disorders after youth leave detention.

Objective: To examine changes in the prevalence and persistence of psychiatric disorders during the 5 years after detention, focusing on sex and racial/ethnic differences.

Design: Prospective longitudinal study with up to 5 interviews (1829 youth: 1172 males and 657 females). To ensure representation of key demographic subgroups, the randomly selected sample was stratified by sex, race/ethnicity (African American, non-Hispanic white, and Hispanic), age, and legal status (juvenile or adult court).

Setting: The Northwestern Juvenile Project, sampling youth from the Cook County Juvenile Temporary Detention Center, Chicago, Illinois.

Participants: Detained youth, aged 10 to 18 years at baseline interview.

Main Outcome Measures: At baseline, the Diagnostic Interview Schedule for Children Version 2.3. At fol-

low-up interviews, the Diagnostic Interview Schedule for Children Version IV (Child and Young Adult versions) and the Diagnostic Interview Schedule Version IV (substance use disorders and antisocial personality disorder).

Results: Five years after baseline, more than 45% of males and nearly 30% of females had 1 or more psychiatric disorders with associated impairment. More than 50% of males and more than 40% of females had 1 or more psychiatric disorders without impairment. Substance use disorders were the most common; males, however, had higher rates over time (5 years after baseline, adjusted odds ratio [AOR], 2.61; 95% CI, 1.96-3.47). Non-Hispanic whites and Hispanics also had higher rates of substance use disorders vs African Americans (AOR, 1.96; 95% CI, 1.54-2.49 and AOR, 1.59; 95% CI, 1.24-2.03). Females had higher rates of major depression over time (AOR, 1.59; 95% CI, 1.22-2.08).

Conclusions: Although prevalence rates of most psychiatric disorders declined as youth aged, a substantial proportion of delinquent youth continue to have disorders. There are notable sex and racial/ethnic differences in the prevalence and persistence of psychiatric disorders in this population.

Arch Gen Psychiatry. 2012;69(10):1031-1043

PSYCHIATRIC DISORDERS ARE prevalent among incarcerated juveniles.¹⁻¹⁴ A 2008 literature review¹⁴ concluded that psychiatric disorders are substantially more common in adolescents in detention than among adolescents in the general population. The Northwestern Juvenile Project¹⁵ found that at intake to detention, even after excluding conduct disorder, more than 60% of juvenile detainees met diagnostic criteria with diagnosis-specific impairment for 1 or more psychiatric disorders. Among youth incarcerated for 9 months, Karnik et al¹⁶ found even higher rates: approximately 90% of de-

tainees had a psychiatric disorder other than conduct disorder or oppositional defiant disorder. Using the lower rate,¹⁵ more than 55 000 of the 92 854 youth held in US correctional facilities each day¹⁷ have 1 or more psychiatric disorders.

Many psychiatric disorders are likely to persist as these juveniles become young adults. Risk factors for psychiatric disorders are common among delinquent youth: maltreatment,¹⁸⁻²⁰ dysfunctional families,^{19,21} family substance abuse,¹⁸ and brain injury.²² Because delinquent youth have few protective factors to offset these risks, many are vulnerable to continued psychiatric morbidity as they age.¹⁸

Author Affiliations are listed at the end of this article.

Despite their importance, few longitudinal studies have examined the prevalence and persistence of psychiatric disorders after youth leave detention. Instead, studies²³⁻²⁶ of delinquent youth have focused on the association between psychiatric disorders and criminal recidivism, antisocial behavior, or social functioning. We could find only one longitudinal study of the persistence and prevalence of psychiatric disorders in detained youth. Harrington et al²⁷ found that 2 years after detention, many mental health problems persisted or had worsened. Their sample, however, excluded females, was 80% white, and was too small (N=97) to permit detailed analyses. Moreover, the study was conducted in the United Kingdom, limiting generalizability to juvenile detainees in the United States.

The related literature—longitudinal studies of “high-risk” youth—also provides little information. Youth with histories of detention have been included in studies of high-risk youth: the homeless,^{28,29} youth living in impoverished or high-crime neighborhoods,³⁰⁻³⁴ and the offspring of parents who used substances or had psychiatric disorders.^{29,35-46} Yet, none of these studies distinguished between youth with and without histories of detention (summary tables available on request).

Delinquent youth are also underrepresented in general population longitudinal studies, such as household-based surveys (eg, Great Smoky Mountains Study of Youth,^{47,48} National Comorbidity Surveys,⁴⁹⁻⁵¹ and National Epidemiologic Survey on Alcohol and Related Conditions⁵²) and studies drawn from school-based samples (eg, Oregon Adolescent Depression Project^{53,54} and Pittsburgh Youth Study⁵⁵) and general pediatric clinics.^{56,57} Household surveys typically exclude persons in correctional institutions.^{47,50,58-61} School-based samples omit youth who are truant or miss school because they are detained.^{53-55,62-65} Samples drawn from general pediatric clinics omit youth who do not receive medical care.^{56,57} Even if sampled initially, youth may be lost to follow-up when they are incarcerated because they cannot be found and because studying prisoners requires special procedures and approvals from the Secretary of Health and Human Services.⁶⁶

In sum, to our knowledge, no large-scale longitudinal study has examined the prevalence and persistence of psychiatric disorders after youth leave detention. This omission is critical. Juvenile detainees have a median length of stay of only 2 weeks.⁶⁷ Thus, juvenile detainees become a community public health problem when they are released and may continue to burden society as they age. Epidemiologic studies are the first step to improve prevention and treatment in correctional facilities and in the community.⁶⁸ Data are also needed to address health disparities, a priority of Healthy People 2020⁶⁹ and the Institute of Medicine.⁷⁰ Of the approximately one-half million incarcerated youth and young adults (aged 24 years or younger), nearly two-thirds are African American or Hispanic,^{17,71} compared with one-third in the general population.⁷²

This study presents data from the Northwestern Juvenile Project, the first comprehensive longitudinal study of psychiatric disorders in youth after they leave detention. The sample is large (N=1829), racially/ethnically diverse, and includes males and females. We examined changes in the prevalence and persistence of disorders

during the first 5 years after detention, focusing on sex and racial/ethnic differences.

METHODS

SAMPLE AND PROCEDURES

Baseline Interviews

We recruited a stratified random sample of 1829 youth at intake to the Cook County Juvenile Temporary Detention Center (CCJTDC) in Chicago, Illinois, between November 20, 1995, and June 14, 1998. The CCJTDC is used for pretrial detention and for offenders sentenced for fewer than 30 days. To ensure adequate representation of key subgroups, we stratified our sample by sex, race/ethnicity (African American, non-Hispanic white, Hispanic, or other), age (10-13 years or 14-18 years), and legal status (processed in juvenile or adult court). Final sampling fractions for strata ranged from 0.11 to 0.69.

All detainees who were awaiting the adjudication or disposition of their case were eligible to participate in the study. Among these, 2275 detainees were randomly selected; 4.2% (34 youth and 62 parents or guardians) refused to participate. There were no significant differences in refusal rates by sex, race/ethnicity, or age. Twenty-seven youth left the detention center before an interview could be scheduled; 312 left CCJTDC while we attempted to locate their caretakers for consent. Eleven others were excluded from the sample because they were unable to complete the interview. The final sample size was 1829: 1172 males and 657 females; 1005 African Americans, 296 non-Hispanic whites, 524 Hispanics, and 4 other race/ethnicity; and age range, 10 to 18 years (mean, 14.9 years; median, 15 years) (eTable 1; <http://www.archgenpsychiatry.com>). Face-to-face structured interviews were conducted at the detention center in a private area, most within 2 days of intake.

Follow-up Interviews

Our design included (1) follow-up interviews at 3 and 4½ years after baseline for the entire sample and (2) 2 additional interviews at 3½ and 4 years after baseline for a random subsample of 997 individuals (600 males and 397 females). For each follow-up, we interviewed participants regardless of where they were living: in the community (approximately two-thirds of interviews), at correctional facilities (nearly 30% of interviews), or by telephone if they lived more than 2 hours away (<5% of interviews).

Participants were paid \$25 for the 2- to 3-hour baseline interview and \$50 for each of the 3- to 4-hour follow-up interviews. Most interviewers had graduate degrees in psychology or an associated field and had experience interviewing at-risk youth; one-third were fluent in Spanish. All interviewers were trained for at least 1 month. For each wave, consistency across interviewers was established and maintained via scripted mock interviews after training and midwave. Additional information on our methods is in the eMethods and is also published elsewhere.^{15,73,74}

Procedures to Obtain Assent and Consent at Baseline and Follow-up

For all interviews, participants signed either an assent form (if they were aged <18 years) or a consent form (if they were aged ≥18 years). The Northwestern University Institutional Review Board and the Centers for Disease Control and Prevention Institutional Review Board approved all study proce-

dures and waived parental consent for persons younger than 18 years, consistent with federal regulations regarding research with minimal risk.⁷⁵ We nevertheless attempted to contact parents of minors to obtain their consent and to provide them with information on the study and used an independent participant advocate to represent the minors' interests⁷⁵ (eMethods).

MAIN OUTCOME MEASURES

Baseline

We used the Diagnostic Interview Schedule for Children Version 2.3 (DISC-2.3),^{76,77} the most recent English and Spanish versions then available. This version of the DISC, based on the *DSM-III-R*, assesses the presence of disorders in the past 6 months. The DISC-2.3 is highly structured, contains detailed symptom probes, has acceptable reliability and validity, and requires relatively brief training.^{76,78-81} Because the DISC-2.3 did not include posttraumatic stress disorder (PTSD), we used the module from the DISC-IV when it became available, 13 months after the study began.⁷³ Additional detail on baseline diagnostic decisions can be found elsewhere.^{15,73,74}

Follow-up Interviews

We administered the DISC-IV (Child and Young Adult versions), based on the *DSM-IV*, to assess for schizophrenia, mood disorders, anxiety disorders, attention-deficit/hyperactivity disorder, and disruptive behavior disorders in the past year.^{82,83} Impairment was defined as moderate impairment in at least one area of functioning (criterion A).⁸⁴ To facilitate comparison with studies that do not impose impairment criteria, we present prevalence rates of disorders with and without the impairment criterion; our analyses of changes in disorders over time use disorders with impairment.

To assess substance use disorders and antisocial personality disorder at follow-up, we administered the Diagnostic Interview Schedule Version IV (DIS-IV).^{85,86} We used the DIS-IV to assess substance use disorders because the DISC-IV is not sufficiently detailed for our population. Antisocial personality disorder was assessed for participants 18 years or older (who were no longer eligible for diagnoses of childhood disruptive behavior disorders). Disorders are assessed for the year before the interview. Consistent with the National Comorbidity Survey Replication (NCS-R),⁵⁰ participants who met the criteria for substance use disorder or antisocial personality disorder with "partial recovery" were scored as having the disorder. We did not implement *DSM* exclusionary criteria.

Comparability of Diagnoses Over Time

Our diagnostic measures changed over time for 3 reasons: (1) the release of the DISC-IV (based on *DSM-IV* criteria) mid-study, (2) some participants turned 18 years old and were therefore ineligible for childhood disruptive behavior disorders, and (3) our need to use a more comprehensive measure of substance use disorder (DIS-IV) for the follow-up interviews.

To check that changes in prevalence rates over time were not the result of changes in measurement, we conducted sensitivity analyses. We created a set of adjusted scoring algorithms to maximize comparability among the DISC-2.3, DISC-IV, and DIS-IV criteria while minimizing alterations (Author's Appendix [<http://hdl.handle.net/2166/repo/teplin-0001>]). All analyses were conducted twice, with and without these adjusted criteria. Because there were no substantive differences in findings and to enable comparisons with other

studies, we present the results using the original, unadjusted diagnoses.

STATISTICAL ANALYSIS

Prevalence Rates of Disorder at Specific Times

Because some participants were interviewed more often than others, we used a subset of interviews to summarize prevalence rates at 3 time points: baseline (time 0), time 1, and time 2.

Time 1. Time 1 is the first follow-up interview but excludes interviews that occurred more than 18 months after the interview due date. Using a narrower window would restrict the generalizability of our findings because, in this high-risk and highly mobile population, participants can be difficult to track. The median time between baseline and time 1 was 3.0 years (mean [SD], 3.2 [0.3] years; range, 2.7-4.5 years). For simplicity, we refer to the time 1 interview as occurring approximately 3 years after baseline. As reported in eTable 1, which summarizes sample demographics and retention rates, 90.7% of the participants had a time 1 interview.

Time 2. For each participant, time 2 consists of the earliest follow-up interview that occurred approximately 4.5 years after baseline. As with time 1, we excluded interviews that occurred more than 18 months after this due date. The median time between baseline and the time 2 interview was 4.7 years (mean [SD], 4.9 [0.4] years; range, 4.3-6.0 years). To ensure that prevalence rates reflect temporally distinct cross-sections of the sample, we required at least 16 months between the time 1 and time 2 interviews. We subsequently refer to the time 2 interviews as occurring approximately 5 years after baseline. As reported in eTable 1, 85.3% of participants had a time 2 interview.

To generate prevalence rates that reflect CCJTDC's population, each participant was assigned a sampling weight augmented with a nonresponse adjustment to account for missing data.⁸⁷ Taylor series linearization was used to estimate standard errors.^{88,89}

Changes in Prevalence Over Time

We used all available interviews, with an average of 3.9 interviews per person (range, 1-5 interviews per person); 88.7% of the participants completed all scheduled interviews. Using generalized estimating equations (GEEs),⁹⁰ we fit marginal models examining (1) differences in the prevalence of disorders by sex and race/ethnicity over time and (2) changes in disorders as the youth aged. Disorder was modeled as binomial with a logit link function. We used a robust sandwich estimator with an unstructured correlation matrix; in the few instances in which models failed to converge, we specified an exchangeable correlation structure. All statistically significant ($P < .05$) odds ratios contrasting sex and racial/ethnic differences are noted and were determined using GEEs. Unless otherwise noted, these odds ratios contrast sex and race/ethnicity over time. Only 2 sets of odds ratios—those examining persistence of disorders and those examining prevalence of disorders among participants living in the community—contrast sex and race/ethnicity at single time points (time 1 or time 2). These were estimated using logistic regression.

All GEE models included covariates for sex, race/ethnicity (African American, Hispanic, or non-Hispanic white), and aging (time since baseline). We also included age at baseline (10-18

years) and legal status at detention (processed in juvenile or adult court) because they were stratification characteristics. The 4 participants who identified themselves as *other* race/ethnicity at baseline were excluded. We estimated models with quadratic terms for aging when the linear term was statistically significant. When main effects were significant, we estimated models with the corresponding interaction terms. Only interaction terms or quadratic aging terms that reached statistical significance were included in the final models. For models with significant interactions between sex and aging, we report model-based odds ratios for sex differences at 3 and 5 years after baseline. The GEE models were estimated with sampling weights to account for study design. We examined the sensitivity of our models to attrition by using weights that combined the probability of being sampled with the probability of dropping out.⁹¹ Results (available on request) were substantially similar.

The GEE analyses were conducted on disorders (with the impairment criterion) except for (1) schizophrenia, because there were too few cases; (2) hypomania, because participants with this disorder were significantly more likely to be missing follow-up data than were those without the disorder (33.3% vs 11.0%, $P = .01$); and (3) diagnoses assessed only for juveniles (attention-deficit/hyperactivity disorder, conduct disorder, and oppositional defiant disorder) or adults (antisocial personality disorder), because nearly all participants became 18 years old or older during the follow-up period. Instead, we estimated changes over time in any disruptive behavior disorder, defined as conduct disorder or oppositional defiant disorder (for participants aged <18 years) or antisocial personality disorder (for participants aged ≥ 18 years).

Because detention may alter access to substances, all models describing substance use disorders included linear and quadratic terms for time incarcerated prior to the interviews. When substance use disorder was measured at a follow-up interview, we used number of days incarcerated in the past year because disorders were assessed in the past year. When substance use disorder was measured at the baseline interview (and therefore assessed in the past 6 months), we used time incarcerated in the past 90 days, the best available estimate. All analyses were conducted using commercial software (Stata 11; StataCorp) with its survey routines.⁹²

RESULTS

PREVALENCE

Table 1 and **Table 2** report prevalence rates of disorders—computed with and without the impairment criterion—at baseline, time 1, and time 2 for males and females. At time 2, more than 45% of the males and nearly 30% of the females had any disorder with impairment. More than 50% of the males and more than 40% of the females had 1 or more psychiatric disorders without impairment. Even excluding disruptive behavior disorders, 36.9% of males and 25.3% of females had any disorder.

Prevalence rates of disorder by race/ethnicity for males and females are given in eTable 2 and eTable 3. Among males, 44.3% of African Americans, 49.8% of Hispanics, and 63.9% of non-Hispanic whites had any disorder at time 2. More than one-quarter of African American females and more than one-third of Hispanic and non-Hispanic white females had any disorder.

MOOD DISORDERS

Other than mania, prevalence rates decreased as the participants aged. For every additional year, there was an approximately 10% decrease in the odds of any mood disorder (adjusted odds ratio [AOR], 0.91; 95% CI, 0.85-0.97) and its subcategories: any major mood disorder (AOR, 0.89; 95% CI, 0.82-0.96) and major depression (AOR, 0.91; 95% CI, 0.84-0.98). The prevalence of dysthymia also decreased with age (AOR, 0.57; 95% CI, 0.46-0.70).

Sex Differences

Over time, females had higher rates of any mood disorder (AOR, 1.33; 95% CI, 1.05-1.68) and its subcategories: any major mood disorder (AOR, 1.54; 95% CI, 1.19-2.00) and major depression (AOR, 1.59; 95% CI, 1.22-2.08). **Figure 1** illustrates prevalence rates of major mood disorders over time by sex.

Racial/Ethnic Differences

The only significant racial/ethnic difference was for mania, which was more prevalent among minorities over time (African American vs non-Hispanic white: AOR, 6.92; 95% CI, 1.78-26.88; Hispanic vs non-Hispanic white: AOR, 8.01; 95% CI, 2.05-31.38).

ANXIETY DISORDERS

The prevalence of panic disorder increased slightly overall (AOR, 1.30 per year; 95% CI, 1.03-1.65). **Figure 1** shows changes in prevalence rates over time by sex.

Sex Differences

Females had higher rates of any anxiety disorder (AOR, 1.42; 95% CI, 1.06-1.91). Although rates of PTSD appeared to decrease, especially for females, this trend was not statistically significant (AOR, 0.93; 95% CI, 0.81-1.05).

Racial/Ethnic Differences

Compared with non-Hispanic whites, Hispanics had more than twice the odds of any anxiety disorder (AOR, 2.18; 95% CI, 1.37-3.49) and its subcategory PTSD (AOR, 2.82; 95% CI, 1.66-4.80). Compared with African Americans, Hispanics had 3.82 times the odds of panic disorder (95% CI, 1.54-9.47) and 1.54 times the odds of PTSD (95% CI, 1.01-2.34). In addition, African Americans were more likely than non-Hispanic whites to have PTSD (AOR, 1.84; 95% CI, 1.10-3.07), although non-Hispanic whites were more likely than African Americans to have panic disorder (AOR, 3.56; 95% CI, 1.28-9.94).

DISRUPTIVE BEHAVIOR DISORDERS

The prevalence of any disruptive behavior disorder decreased over time, but the rate of decrease depended on

Table 1. Prevalence of Disorder at Baseline, Time 1, and Time 2 for Males^a

Disorder	Prevalence, % (SE)					
	With Impairment Criteria			Without Impairment Criteria		
	Baseline	Time 1	Time 2	Baseline	Time 1	Time 2
Any disorder ^b	61.8 (4.1)	51.7 (2.8)	46.5 (2.9)	67.0 (4.0)	57.9 (2.7)	51.7 (2.9)
Any disorder, except behavioral ^b	60.2 (4.1)	45.1 (2.7)	36.9 (2.7)	64.4 (4.0)	50.7 (2.8)	42.4 (2.8)
Schizophrenia ^c	...	0.2 (0.1)	0.1 (0.1)	...	0.2 (0.1)	0.1 (0.1)
Any mood disorder	15.8 (1.8)	14.9 (1.9)	8.8 (1.6)	18.8 (1.9)	16.2 (2.0)	9.7 (1.7)
Any major mood disorder	12.7 (1.7)	9.5 (1.6)	6.4 (1.4)	14.3 (1.8)	10.5 (1.6)	7.3 (1.5)
Mania	2.0 (0.7)	0.5 (0.2)	0.7 (0.4)	2.2 (0.8)	0.6 (0.2)	0.7 (0.4)
Major depression	11.0 (1.6)	9.1 (1.5)	6.4 (1.4)	13.0 (1.7)	10.1 (1.6)	7.3 (1.5)
Hypomania ^d	2.1 (0.7)	6.3 (1.4)	2.1 (0.7)	2.1 (0.7)	6.3 (1.4)	2.1 (0.7)
Dysthymia	9.9 (1.5)	1.1 (0.6)	1.0 (0.6)	12.2 (1.7)	1.5 (0.7)	1.0 (0.6)
Any anxiety disorder ^b	10.8 (2.7)	9.8 (1.6)	7.7 (1.6)	16.7 (3.2)	10.8 (1.6)	9.2 (1.7)
Generalized anxiety disorder	3.8 (1.0)	2.6 (0.9)	1.9 (0.9)	7.1 (1.3)	2.6 (0.9)	1.9 (0.9)
Panic disorder	0.1 (0.1)	1.4 (0.5)	0.5 (0.2)	0.3 (0.1)	2.2 (0.6)	1.0 (0.5)
Posttraumatic stress disorder ^b	7.9 (2.3)	7.6 (1.4)	5.4 (1.3)	10.9 (2.6)	8.0 (1.4)	6.4 (1.4)
Attention-deficit/hyperactivity disorder, age <18 y ^e	11.2 (1.5)	6.5 (2.3)	4.2 (2.3)	16.6 (1.8)	7.0 (2.3)	5.6 (2.7)
Any disruptive behavior disorder ^f	29.5 (2.2)	21.9 (2.2)	22.1 (2.4)	41.8 (2.4)	28.7 (2.4)	26.1 (2.5)
Conduct disorder, age <18 y ^e	24.3 (2.0)	20.5 (3.8)	9.3 (3.2)	37.8 (2.4)	29.7 (4.6)	13.9 (4.0)
Oppositional defiant disorder, age <18 y ^e	12.6 (1.6)	15.7 (3.8)	10.0 (3.4)	14.5 (1.7)	17.6 (4.0)	10.6 (3.6)
Antisocial personality disorder, age ≥18 y ^g	NA	20.4 (2.4)	22.2 (2.4)	NA	26.6 (2.7)	26.2 (2.5)
Any substance use disorder	45.7 (2.5)	29.4 (2.4)	28.0 (2.4)	50.7 (2.5)	37.3 (2.6)	32.2 (2.5)
Alcohol use disorder	19.9 (2.0)	15.6 (1.9)	17.1 (2.0)	25.9 (2.2)	18.6 (2.0)	19.0 (2.0)
Drug use disorder	42.3 (2.4)	22.0 (2.2)	18.8 (2.0)	45.4 (2.5)	30.2 (2.4)	23.4 (2.2)

Abbreviation: NA, not applicable.

^aDescriptive statistics are weighted to adjust for sampling design and reflect the demographic characteristics of the Cook County Juvenile Temporary Detention Center. Because some participants were interviewed more often than others, we used a subset of interviews to summarize prevalence rates at 3 time points: baseline (time 0), time 1, and time 2. The sample included 1172, 1054, and 993 males at baseline, time 1, and time 2, respectively.

^bAssessed at baseline for 541 males who were interviewed after the Diagnostic Interview Schedule for Children, Version IV posttraumatic stress disorder module became available.

^cNot assessed at baseline.

^dBecause there are no impairment criteria for a diagnosis of hypomania, prevalence rates of hypomania with and without impairment are the same.

^eAssessed for males younger than 18 years at time 1 (n = 350) and time 2 (n = 96).

^fFor participants younger than 18 years, any disruptive behavior disorder is defined as having conduct disorder or oppositional defiant disorder. For participants aged 18 years or older, it is defined as having antisocial personality disorder.

^gAssessed for males aged 18 years or older at time 1 (n = 704) and time 2 (n = 897); not assessed at baseline because the sample consisted of juveniles.

sex. **Figure 2** illustrates these sex differences over time.

Sex Differences

Males and females did not have significantly different rates of any disruptive behavior disorder at baseline, but prevalence decreased faster among females than among males. Three years after baseline, males had 1.82 times the odds of any disruptive behavior disorder (95% CI, 1.46-2.26) compared with females; 5 years after baseline, males had 2.95 times the odds (95% CI, 2.16-4.02).

Racial/Ethnic Differences

As illustrated in Figure 2, non-Hispanic whites had the highest rates of any disruptive behavior disorder over time (non-Hispanic white vs African American: AOR, 2.34; 95% CI, 1.80-3.04; non-Hispanic white vs Hispanic: AOR, 1.56; 95% CI, 1.16-2.11), followed by Hispanics (Hispanic vs African American: AOR, 1.50; 95% CI, 1.10-2.04).

SUBSTANCE USE DISORDERS

Substance use disorders were the most prevalent disorders in our sample. The prevalence generally decreased over time, but the rate of decrease depended on sex. Figure 2 illustrates sex and racial/ethnic differences over time.

Sex Differences

At baseline, compared with females, males had about one-third greater odds of any substance use disorder (AOR, 1.34; 95% CI, 1.05-1.71) and its subcategory, drug use disorder (AOR, 1.37; 95% CI, 1.07-1.75); rates of alcohol use disorder were not significantly different. By the follow-up interviews, however, the disparities between males and females increased substantially because prevalence rates decreased faster for females than for males. Three years after baseline, compared with females, males had approximately twice the odds of any substance use disorder (AOR, 2.00; 95% CI, 1.64-2.43) and its subcategories, drug use disorder (AOR, 2.01; 95% CI, 1.62-2.49) and alcohol use disorder (AOR, 1.97; 95% CI, 1.53-2.53). Five years after

Table 2. Prevalence of Disorder at Baseline, Time 1, and Time 2 for Females^a

Disorder	Prevalence, % (SE)					
	With Impairment Criteria			Without Impairment Criteria		
	Baseline	Time 1	Time 2	Baseline	Time 1	Time 2
Any disorder ^b	65.3 (2.8)	42.9 (2.3)	29.0 (2.1)	74.4 (2.4)	52.5 (2.3)	40.8 (2.5)
Any disorder, except behavioral ^b	62.9 (3.5)	38.6 (2.3)	25.3 (2.0)	69.7 (2.6)	45.7 (2.3)	32.1 (2.1)
Schizophrenia ^c	...	0.2 (0.2)	0 (...)	...	0.2 (0.2)	0 (...)
Any mood disorder	22.8 (2.1)	17.0 (1.6)	11.9 (1.4)	27.7 (2.1)	19.5 (1.7)	12.5 (1.4)
Any major mood disorder	19.9 (2.1)	13.2 (1.4)	10.4 (1.3)	23.0 (2.1)	15.2 (1.5)	10.8 (1.3)
Mania	1.2 (0.4)	1.6 (0.5)	1.5 (0.5)	1.8 (0.5)	2.3 (0.6)	1.7 (0.5)
Major depression	18.9 (2.1)	12.7 (1.4)	10.2 (1.3)	21.6 (2.1)	14.6 (1.5)	10.4 (1.3)
Hypomania ^d	0.3 (0.2)	4.1 (0.8)	0.8 (0.4)	0.3 (0.2)	4.1 (0.8)	0.8 (0.4)
Dysthymia	12.5 (1.3)	1.5 (0.5)	0.7 (0.3)	15.8 (1.4)	1.9 (0.6)	0.9 (0.4)
Any anxiety disorder ^b	18.9 (3.4)	12.4 (1.5)	8.1 (1.2)	24.1 (3.4)	14.1 (1.5)	8.7 (1.3)
Generalized anxiety disorder	5.1 (0.9)	3.3 (0.8)	2.1 (0.6)	7.3 (1.0)	3.3 (0.8)	2.1 (0.6)
Panic disorder	1.0 (0.4)	2.3 (0.6)	0.9 (0.4)	1.5 (0.4)	3.0 (0.7)	1.1 (0.5)
Posttraumatic stress disorder ^b	14.6 (3.4)	7.9 (1.1)	5.8 (1.0)	17.7 (3.4)	9.3 (1.2)	6.4 (1.0)
Attention-deficit/hyperactivity disorder, age <18 y ^e	16.4 (1.5)	9.7 (2.6)	0 (...)	21.4 (1.6)	11.1 (2.7)	0 (...)
Any disruptive behavior disorder ^f	34.6 (2.1)	16.6 (2.1)	7.3 (1.1)	45.7 (2.1)	24.6 (2.2)	15.4 (2.5)
Conduct disorder, age <18 y ^e	28.5 (2.1)	13.5 (3.0)	...	40.6 (2.1)	20.9 (3.5)	18.8 (8.6)
Oppositional defiant disorder, age <18 y ^e	15.1 (1.4)	9.1 (2.4)	4.0 (3.9)	17.5 (1.5)	12.9 (2.9)	8.1 (5.6)
Antisocial personality disorder, age ≥18 y ^g	NA	15.4 (2.6)	7.2 (1.2)	NA	23.1 (2.6)	15.1 (2.6)
Any substance use disorder	41.7 (2.3)	18.0 (2.1)	13.5 (1.5)	46.8 (2.2)	24.7 (2.1)	20.4 (1.7)
Alcohol use disorder	20.0 (2.1)	7.8 (1.1)	6.0 (1.0)	26.5 (2.1)	10.6 (1.2)	8.5 (1.2)
Drug use disorder	38.4 (2.4)	12.7 (2.0)	9.2 (1.2)	42.6 (2.3)	18.5 (2.1)	15.3 (1.5)

Abbreviation: NA, not applicable.

^aDescriptive statistics are weighted to adjust for sampling design and reflect the demographic characteristics of the Cook County Juvenile Temporary Detention Center. Because some participants were interviewed more often than others, we used a subset of interviews to summarize prevalence rates at 3 time points: baseline (time 0), time 1, and time 2. The sample included 657, 605, and 568 females at baseline, time 1, and time 2, respectively.

^bAssessed at baseline for 374 females who were interviewed after the Diagnostic Interview Schedule for Children, Version IV posttraumatic stress disorder module became available.

^cNot assessed at baseline.

^dBecause there are no impairment criteria for a diagnosis of hypomania, prevalence rates of hypomania with and without impairment are the same.

^eAssessed for females younger than 18 years at time 1 (n = 148) and time 2 (n = 21). We did not estimate prevalence for cells with fewer than 20 participants.

^fFor participants younger than 18 years, any disruptive behavior disorder is defined as having conduct disorder or oppositional defiant disorder. For participants aged 18 years or older, it is defined as having antisocial personality disorder.

^gAssessed for females aged 18 years or older at time 1 (n = 457) and time 2 (n = 547); not assessed at baseline because the sample consisted of juveniles.

baseline, males had more than 2.5 times the odds of these disorders compared with females (any substance use disorder: AOR, 2.61; 95% CI, 1.96-3.47; drug use disorder: AOR, 2.60; 95% CI, 1.88-3.58; and alcohol use disorder: AOR, 2.87; 95% CI, 2.00-4.13). Although the prevalence rates of most disorders decreased for males and females (Figure 2), 3 years after baseline, rates of alcohol disorder were no longer decreasing among males (AOR, 1.03 per year; 95% CI, 0.96-1.11).

Racial/Ethnic Differences

Even after adjusting for time spent in correctional facilities, substance use disorders were more common among non-Hispanic whites and Hispanics than among African Americans (Figure 2). Compared with African Americans, non-Hispanic whites had nearly twice the odds of any substance use disorder and its subcategory, drug use disorder (AOR, 1.96; 95% CI, 1.54-2.49 and AOR, 1.91; 95% CI, 1.50-2.43, respectively), and 1.57 times the odds of alcohol disorder (95% CI, 1.15-2.13). Hispanics had approximately 1.5 times the odds of any

substance use disorder and its subcategory, drug use disorder, compared with African Americans (AOR, 1.59; 95% CI, 1.24-2.03 and AOR, 1.46; 95% CI, 1.12-1.92, respectively).

We also analyzed substance use disorders without impairment criteria, an approach used by recent studies of the general population.^{50,58} Results were substantially similar (available on request).

SUBSTANCE USE DISORDERS AMONG PARTICIPANTS LIVING IN THE COMMUNITY AT TIME 2

Because substance use is restricted in jails and prisons, we examined rates of substance use disorders only among participants who had lived in the community the entire year before time 2 (345 males and 479 females). These prevalence rates, and the demographic differences, were substantially similar to those in the entire sample (eTable 4).




Figure 1. Past-year prevalence of major mood disorders (major depressive disorder [MDD], mania) and anxiety disorders (posttraumatic stress disorder [PTSD], generalized anxiety disorder [GAD], and panic disorder) by sex.

PERSISTENCE

Table 3 reports the persistence of disorders: among participants who had the disorder at baseline, the proportion who still had the disorder at time 1 or time 2. For most disorders, rates of persistence were higher at time 1 than at time 2.

Sex Differences

Regardless of sex, approximately 1 in 5 participants had a mood disorder that persisted to time 2. Substance use disorders were among the most persistent disorders for both males and females but were significantly more likely to persist among males. Any disruptive behavior disorder was also among the most persistent disorders in males and at time 2 was significantly more likely to persist in males than in females.

Racial/Ethnic Differences

Among males, there were no significant racial/ethnic differences in the persistence of disorders (eTable 5). Among females, there were several significant differences (eTable 6). At time 1, any substance use disorder and its subcategory, alcohol use disorder, had approximately 3 times the odds of persisting among non-Hispanic whites and Hispanics than among African Americans. At time 2, drug use disorders also had greater odds of persisting among non-Hispanic whites than among African Americans.

COMMENT

Although prevalence rates of most psychiatric disorders declined over time, a substantial proportion of delinquent youth continued to have disorders as they aged. For some youth, detention may coincide with a period of crisis that subsequently abates. Many youth, however, continue to struggle: 5 years after detention, when participants were ages 14 to 24 years, nearly half of males and nearly 30% of females had 1 or more psychiatric disorders with associated impairment.

Substance use and disruptive behavior disorders continued to be the most common disorders. For many delinquent youth—especially males—externalizing disorders were not limited to adolescence. Five years after baseline, males had 2 to 3 times the odds of having substance use and disruptive behavior disorders compared with females. Furthermore, the disparity between males and females increased over time. Males were also more likely than females to have persistent substance use disorders and disruptive behavior disorder.

The observed sex differences in externalizing disorders are consistent with those in the general population, where males are up to 10 times more likely than females to persist in antisocial behavior from childhood to adulthood.⁹³ Why might males fare worse than females? First, delinquent males are less likely to receive services.⁹⁴ Second, delinquent males may have fewer opportunities to assume age-appropriate social roles (ie, jobs, postsecondary schooling)—turning points that might reduce problem be-




Figure 2. Past-year prevalence of substance use disorders (alcohol disorder [Alc], drug disorder) and any disruptive behavior disorder (DBD) by sex and race/ethnicity.

haviors.⁹⁵ Finally, precocious transition to adult social roles, such as parenthood, may be associated with worse outcomes for males than for females.⁹⁶⁻⁹⁸

As in the general population,⁹⁹ females had higher rates of internalizing disorders than males. The persistence of mood disorders, about 20%, was similar in males and females.

Rates of substance use disorders and disruptive behavior disorders were lower in African Americans than non-Hispanic whites. These findings may reflect underlying racial/ethnic disparities in the legal system^{71,100-102} and the different pathways by which non-Hispanic whites and racial/ethnic minorities enter the system. We found racial/ethnic differences in substance use disorders even after taking into account that African Americans spend more time in correctional facilities, where access to alcohol and drugs is restricted.^{100,101} Our findings add to the growing debate about how the “war on drugs” has affected the disproportionate incarceration of African Americans. Many investigators suggest that disproportionate confinement of racial/ethnic minorities for drug offenses is the result, in part, of disparate enforcement of drug laws in African American communities rather than higher rates of use or dealing.¹⁰³⁻¹⁰⁷ Our findings appear consistent with this view.

It is difficult to compare specific prevalence rates in our sample with those in the general population because differences in instrumentation and sample demographics limit meaningful comparisons. The National Comorbidity Survey Replication (NCS-R) provides the most comparable data to our time 2 interview. Although the NCS-R used different, and often less stringent, impairment criteria and did not assess the same disorders (eg, antisocial personality disorder), it provides *DSM-IV* diagnoses for a similarly aged sample (18-24 years).^{108,109}

The most marked discrepancies between our findings and those of NCS-R were for drug use disorders, regardless of sex and race/ethnicity. For example, we found that approximately 20% of males had a drug use disorder compared with approximately 7% in the NCS-R; nearly 14% of Hispanic females and nearly 25% of Hispanic males had a drug use disorder compared with less than 5% of Hispanics in the NCS-R.⁹⁹

Changes in prevalence over time mirror those in the general population for most disorders. As summarized in the recent literature review by Costello et al,¹¹⁰ many disorders in the general population decrease from adolescence to young adulthood, with the exception of panic disorders and substance use disorders, which increase^{48,111-116}; findings on depression have been equivocal.^{48,111-116} Again, our sample is most notably different from the general population in substance use disorders: our rates decreased over time. Perhaps substance abuse peaks earlier in delinquent youth, coinciding with the general course of delinquent behavior.^{117,118} In contrast, general population youth may experience events that increase the likelihood of substance abuse as they age,^{119,120} such as living in college dormitories, freedom from social controls, and delays in assuming adult responsibilities such as parenting—events less likely to be experienced by delinquent youth.¹²¹

In terms of persistence, the most recent comparable investigation⁴⁸ that was conducted in the United States using a sample of similar age- and *DSM*-based criteria (albeit different measures) found lower rates of persistence of depression and disruptive behavior disorders than in our sample. (Persistence of substance use disorders is not comparable because the definitions differed; Copeland et al⁴⁸ used more liberal impairment criteria and included nicotine.)

Table 3. Persistence of Disorders From Baseline to Time 1 and From Baseline to Time 2, by Sex^a

Disorder	Males			Females			OR (95% CI) ^b	
	No. With Disorder at Baseline	% (SE) Persisting		No. With Disorder at Baseline	% (SE) Persisting		Persistence to Time 1, M vs F	Persistence to Time 2, M vs F
		Time 1	Time 2		Time 1	Time 2		
Any disorder ^c	335	52.1 (5.8)	48.7 (6.1)	233	54.0 (4.2)	34.9 (4.9)
Any mood disorder	163	28.0 (6.1)	18.9 (5.7)	144	30.4 (4.6)	20.9 (4.2)
Any major mood disorder	127	18.8 (5.6)	17.3 (6.1)	124	25.9 (4.7)	19.2 (4.3)
Mania ^d	16	8
Major depression	116	20.0 (6.2)	15.7 (6.3)	118	25.3 (4.7)	17.9 (4.2)
Hypomania ^d	16	2
Dysthymia	98	1.6 (1.0)	0.0 (...)	87	3.8 (2.2)	3.9 (2.2)
Any anxiety disorder ^c	50	6.3 (3.1)	14.8 (9.7)	58	19.3 (6.3)	17.3 (7.1)
Generalized anxiety disorder	34	18.8 (11.3)	0.0 (...)	35	9.4 (5.2)	10.4 (7.0)
Panic disorder ^d	3	7
Posttraumatic stress disorder ^c	37	5.8 (3.6)	4.0 (2.9)	42	4.5 (3.2)	5.4 (3.7)
Any disruptive behavior disorder	388	36.6 (4.7)	31.1 (4.7)	230	30.7 (4.8)	10.5 (2.4)	...	3.83 (1.98-7.43)
Any substance use disorder ^e	517	38.0 (3.8)	34.3 (3.8)	266	30.5 (4.6)	18.2 (2.7)	1.95 (1.09-3.51)	3.68 (2.10-6.44)
Alcohol use disorder ^e	219	30.4 (5.6)	29.3 (5.7)	127	16.2 (3.4)	12.3 (3.3)	2.66 (1.21-5.85)	3.62 (1.49-8.77)
Drug use disorder ^e	482	28.4 (3.7)	23.1 (3.4)	246	22.6 (5.2)	13.6 (2.4)	...	2.96 (1.59-5.51)

Abbreviation: OR, odds ratio.

^aRates of persistence and their associated SEs are weighted to adjust for sampling design and reflect the demographic characteristics of the Cook County Juvenile Temporary Detention Center. Persistence is presented for disorders assessed with impairment criteria, with the exception of hypomania, which has no impairment criteria for diagnosis. We do not present rates of persistence for disorders assessed only for juveniles or adults (attention-deficit/hyperactivity disorder, conduct disorder, oppositional defiant disorder, or antisocial personality disorder).

^bOdds ratios are included only for statistically significant findings.

^cAssessed at baseline for participants who were interviewed after the Diagnostic Interview Schedule for Children, Version IV posttraumatic stress disorder module became available (541 males and 374 females).

^dRates of persistence are not presented for disorders with fewer than 20 cases at baseline.

^eOdds ratios describing sex differences in the persistence of substance use disorders are adjusted for number of days incarcerated in the past year.

LIMITATIONS

Our data are subject to the limitations of self-report. Moreover, it was not feasible to study more than one jurisdiction; because the prevalence of psychiatric disorders may vary across jurisdictions,^{14,122,123} generalizability may be limited. We cannot know whether psychiatric disorders increase the likelihood of arrest and detention or vice versa. Findings might have been marginally different had we been able to use identical measures and time frames at the baseline and follow-up interviews. Rates would likely have been higher had caretakers been available to provide independent reports at baseline.¹⁵ At the follow-up interviews, many participants were not eligible to have caretakers interviewed because the participants were older than 17 years or did not live with a caretaker. Although retention rates were high, participants who missed interviews might be more likely to have had disorders than those who were found. Our findings do not take into account mental health services that might have been provided. Despite these limitations, our findings have implications for future research and mental health policy.

DIRECTIONS FOR FUTURE RESEARCH

1. Retain incarcerated persons in longitudinal studies of psychiatric disorders. Most large-scale longitudinal studies of the general population do not retain persons who become incarcerated by follow-up (eg, the National Epidemiologic Survey on Alcohol and Related Conditions [B.

Grant, PhD, PhD, email communication, August 13, 2010]) or reinterview too few to analyze (the Epidemiologic Catchment Area Study [W. Eaton, PhD, oral communication, August 11, 2010]). Thus, these samples are biased: they systematically exclude persons who, as our study suggests, are likely to have psychiatric disorders and poor outcomes. Excluding incarcerated persons biases prevalence rates, especially for African American males. At any given time, nearly 1 in 9 African American males aged 25 to 34 years are incarcerated.⁷¹ To address health disparities, we must include correctional populations, which have increased from approximately 2 million in 2000 to nearly 2.4 million in 2008.^{71,124}

2. Add variables on incarceration history to general population studies. Although many studies examine the prevalence of psychiatric disorders in incarcerated populations,^{13-16,74,122,125-127} few focus on the effect of incarceration on psychiatric disorders. We suggest that general population epidemiologic surveys add the following variables: number of incarcerations, age at time of incarceration, length of incarcerations, and experiences in "community corrections" (parole, probation, and community supervision). This strategy would generate necessary information on how disproportionate confinement of racial/ethnic minorities affects health disparities in psychiatric disorders and related outcomes.

3. Include females in longitudinal studies of delinquents. The sex differences observed in our study underscore that findings for males may not generalize to

females. Yet, most longitudinal studies of delinquents exclude females or sample too few to analyze sex differences (summary tables available on request). Future studies must include females and collect data on pregnancy, childbirth, and child-rearing. We will then have the requisite empirical foundation to improve sex-specific mental health services, especially needed now that females constitute an increasing proportion (now 30%¹²⁸) of juvenile arrests.

4. Examine variables that affect trajectories of disorder in high-risk youth. Few studies of high-risk youth examine trajectories of disorder; fewer still examine how potentially modifiable risk and protective factors predict trajectories of disorder. Future studies should investigate how social, cognitive, and biological factors interact to affect trajectories. For example, advances in neuroscience research provide unique opportunities to investigate how developmental differences in emotion regulation interact with “turning points” to alter trajectories.¹²⁹⁻¹³¹

IMPLICATIONS FOR MENTAL HEALTH POLICY

1. Focus on delinquent males. In recent years, innovative programs funded by the Office of Juvenile Justice and Delinquency Prevention¹³²—such as Girl Talk,¹³³ and Girl Scouts in Detention Centers—addressed the needs of delinquent females.¹³⁴⁻¹³⁶ The mental health system must now improve services for males, who constitute 70% of juvenile arrests and 85% of youth in correctional facilities.^{17,128} Our findings demonstrate that interventions for substance use and disruptive behavior disorders are especially needed. Comprehensive interventions, such as functional family therapy,¹³⁷ multidimensional treatment foster care,¹³⁸ and multisystemic therapy,¹³⁹ can be effective. Continued development and dissemination of these programs can further reduce illegal behaviors and provide cost-effective alternatives to incarceration.¹⁴⁰

2. Assess and treat substance use disorders in correctional facilities and after release. Irrespective of sex or race/ethnicity, alcohol and drug use disorders were among the most common and persistent disorders. However, the need for services far exceeds their availability. Approximately one-half of youth in juvenile correctional facilities^{141,142} and approximately three-quarters of youth in adult jails and prisons who need substance abuse treatment do not receive it.¹⁴¹ Incarcerated adults fare much worse: summarizing Department of Justice statistics,^{143,144} a recent study published in *JAMA*¹⁴⁵ concluded that 80% to 85% of adult prisoners who needed treatment for drug abuse did not receive it. After release back to the community, services may be difficult to obtain. The Substance Abuse and Mental Health Services Administration reports, for example, that fewer than 10% of juveniles¹⁴⁶ and adults¹⁴⁷ with an “alcohol use problem” received specialty services in the past year.

Despite the potential of health care reform, the law may not improve mental health services for persons like our participants, who may frequently cycle through correctional facilities. Health care reform has no benefit to prisoners because federal law prohibits Medicaid payments for individuals—juvenile or adult—who are in-

mates of public institutions.¹⁴⁸ Incarceration disrupts community treatment and Medicaid benefits.¹⁴⁹

Moreover, the Patient Protection and Affordable Care Act (PPACA) has limitations. First, although the Supreme Court upheld most provisions of the PPACA,¹⁵⁰ the law does not improve the deteriorating public health infrastructure,^{151,152} where many impoverished persons would receive treatment^{153,154} after being released from corrections. Second, although the PPACA expands Medicaid benefits to persons with incomes up to 133% of the federal poverty level, the Supreme Court ruled that states could opt out.¹⁵⁵ The effect of the Supreme Court’s ruling is not yet known. Many persons like our participants are poor enough to qualify for Medicaid even without the expansion. Moreover, states have strong financial incentives to expand Medicaid.^{156,157} Thus, as of this writing, it is unclear how many additional people would actually receive health care coverage. Since the Supreme Court’s decision, the Congressional Budget Office¹⁵⁸ projected that 30 million “nonelderly residents”—such as undocumented immigrants and persons with low incomes who fail to enroll in Medicaid^{152,153}—will remain uninsured in 2016. Therefore, we must improve services and access to care in corrections and in the community, where virtually all detained juveniles and incarcerated adults will eventually return.

Submitted for Publication: April 4, 2011; final revision received November 18, 2011; accepted November 23, 2011.

Author Affiliations: Health Disparities and Public Policy (Drs Teplin, Welty, and Abram) and Division of Psychology (Dr Washburn), Department of Psychiatry and Behavioral Sciences (Drs Teplin, Welty, Abram, Dulcan, and Washburn), and Department of Preventive Medicine (Dr Welty), Northwestern University Feinberg School of Medicine, Chicago, Illinois; Department of Child and Adolescent Psychiatry, Ann & Robert H. Lurie Children’s Hospital of Chicago (Dr Dulcan); and Alexian Brothers Behavioral Health Hospital, Hoffman Estates, Illinois (Dr Washburn).

Correspondence: Linda A. Teplin, PhD, Health Disparities and Public Policy, Department of Psychiatry and Behavioral Sciences, Northwestern University Feinberg School of Medicine, 710 N Lake Shore Dr, Ste 900, Chicago, IL 60611 (healthdisparities@northwestern.edu).

Author Contributions: Dr Teplin had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. All other authors had full access to all study data.

Financial Disclosure: Dr Dulcan receives book royalties from American Psychiatric Publishing, Inc. She sat on the Attention-Deficit/Hyperactivity Disorder and Reader Disorder and Strattera advisory boards of Eli Lilly and Company, most recently in 2009. She is a consultant for Care Management Technologies, Inc.

Funding/Support: This work was supported by grants R01DA019380, R01DA022953, and R01DA028763 from the National Institute on Drug Abuse; grants R01MH54197 and R01MH59463 from the National Institute of Mental Health (Division of Services and Intervention Research and Center for Mental Health Research on AIDS); and grants 1999-JE-FX-1001, 2005-JL-FX-0288, and 2008-JF-FX-0068 from the Office of Juvenile

Justice and Delinquency Prevention. Major funding was also provided by the National Institute on Alcohol Abuse and Alcoholism; the National Institutes of Health (NIH) Office of Behavioral and Social Sciences Research, Substance Abuse and Mental Health Services Administration (Center for Mental Health Services, Center for Substance Abuse Prevention, and Center for Substance Abuse Treatment); the NIH Center on Minority Health and Health Disparities; the Centers for Disease Control and Prevention (National Center for Injury Prevention and Control and National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention); the NIH Office of Research on Women's Health; the NIH Office of Rare Diseases, Department of Labor, Department of Housing and Urban Development; the William T. Grant Foundation; and the Robert Wood Johnson Foundation. Additional funds were provided by the John D. and Catherine T. MacArthur Foundation, The Open Society Institute, and The Chicago Community Trust.

Online-Only Material: The eMethods and eTables are available at <http://www.archgenpsychiatry.com>.

Additional Contributions: Celia Fisher, PhD, provided invaluable advice on the project. Elena de la Garza Bassett, MS; Lynda Carey, MA; Jennifer Hershfield, MA; Kathryn Jacoby, BA; and Nichole Olson, MS, provided research assistance and editorial assistance. We thank our participants for their time and willingness to participate, as well as the Cook County Juvenile Temporary Detention Center, Cook County Department of Corrections, and Illinois Department of Corrections for their cooperation. We also thank Mark Heyrman, JD, and the anonymous reviewers, who provided insightful and incisive comments on earlier drafts.

REFERENCES

- Davis DL, Bean GJ, Schumacher JE, Stringer TL. Prevalence of emotional disorders in a juvenile justice institutional population. *Am J Forensic Psychol*. 1991; 9(1):5-17.
- Forehand R, Wierson M, Frame C, Kempton T, Armistead L. Juvenile delinquency entry and persistence: do attention problems contribute to conduct problems? *J Behav Ther Exp Psychiatry*. 1991;22(4):261-264.
- Eppright TD, Kashani JH, Robison BD, Reid JC. Comorbidity of conduct disorder and personality disorders in an incarcerated juvenile population. *Am J Psychiatry*. 1993;150(8):1233-1236.
- Rohde P, Mace DE, Seeley JR. The association of psychiatric disorders with suicide attempts in a juvenile delinquent sample. *Crim Behav Ment Health*. 1997; 7:187-200. doi:10.1002/cbm.172.
- Steiner H, Garcia IG, Matthews Z. Posttraumatic stress disorder in incarcerated juvenile delinquents. *J Am Acad Child Adolesc Psychiatry*. 1997;36(3):357-365.
- Timmons-Mitchell J, Brown C, Schulz SC, Webster SE, Underwood LA, Semple WE. Comparing the mental health needs of female and male incarcerated juvenile delinquents. *Behav Sci Law*. 1997;15(2):195-202.
- Cauffman E, Feldman SS, Waterman J, Steiner H. Posttraumatic stress disorder among female juvenile offenders. *J Am Acad Child Adolesc Psychiatry*. 1998; 37(11):1209-1216.
- Duclos CW, Beals J, Novins DK, Martin C, Jewett CS, Manson SM. Prevalence of common psychiatric disorders among American Indian adolescent detainees. *J Am Acad Child Adolesc Psychiatry*. 1998;37(8):866-873.
- Erwin BA, Newman E, McMackin RA, Morrissey C, Kaloupek DG. PTSD, malevolent environment, and criminality among criminally involved male adolescents. *Crim Justice Behav*. 2000;27(2):196-215. doi:10.1177/0093854800027002004.
- Piszka SR, Sherman JO, Barrow MV, Irick S. Affective disorder in juvenile offenders: a preliminary study. *Am J Psychiatry*. 2000;157(1):130-132.
- Shelton D. Emotional disorders in young offenders. *J Nurs Scholarsh*. 2001;33(3):259-263.
- Wasserman GA, McReynolds LS, Lucas CP, Fisher P, Santos L. The voice DISC-IV with incarcerated male youths: prevalence of disorder. *J Am Acad Child Adolesc Psychiatry*. 2002;41(3):314-321.
- Domalanta DD, Risser WL, Roberts RE, Risser JM. Prevalence of depression and other psychiatric disorders among incarcerated youths. *J Am Acad Child Adolesc Psychiatry*. 2003;42(4):477-484.
- Fazel S, Doll H, Långström N. Mental disorders among adolescents in juvenile detention and correctional facilities: a systematic review and meta-regression analysis of 25 surveys. *J Am Acad Child Adolesc Psychiatry*. 2008;47(9): 1010-1019.
- Teplin LA, Abram KM, McClelland GM, Dulcan MK, Mericle AA. Psychiatric disorders in youth in juvenile detention. *Arch Gen Psychiatry*. 2002;59(12): 1133-1143.
- Karnik NS, Soller M, Redlich A, Silverman M, Kraemer HC, Haapanen R, Steiner H. Prevalence of and gender differences in psychiatric disorders among juvenile delinquents incarcerated for nine months. *Psychiatr Serv*. 2009;60(6):838-841.
- Sickmund M, Sladky TJ, Kang W. Census of juveniles in residential placement databook. <http://www.ojdp.gov/ojstatbb/ezacjrp/>. Accessed August 9, 2012.
- Wareham J, Dembo R. A longitudinal study of psychological functioning among juvenile offenders: a latent growth model analysis. *Crim Justice Behav*. 2007; 34(2):259-273. doi:10.1177/0093854806289828.
- Dixon A, Howie P, Starling J. Psychopathology in female juvenile offenders. *J Child Psychol Psychiatry*. 2004;45(6):1150-1158.
- Gover AR. Childhood sexual abuse, gender, and depression among incarcerated youth. *Int J Offender Ther Comp Criminol*. 2004;48(6):683-696.
- Dembo R, Jainchill N, Turner C, Fong C, Farkas S, Childs K. Levels of psychopathy and its correlates: a study of incarcerated youths in three states. *Behav Sci Law*. 2007;25(5):717-738.
- Perron BE, Howard MO. Prevalence and correlates of traumatic brain injury among delinquent youths. *Crim Behav Ment Health*. 2008;18(4):243-255.
- Hiscoke UL, Långström N, Ottosson H, Grann M. Self-reported personality traits and disorders (DSM-IV) and risk of criminal recidivism: a prospective study. *J Pers Disord*. 2003;17(4):293-305.
- Vermeiren R, Schwab-Stone M, Ruchkin V, De Clippele A, Deboutte D. Predicting recidivism in delinquent adolescents from psychological and psychiatric assessment. *Compr Psychiatry*. 2002;43(2):142-149.
- Randall J, Henggeler SW, Pickrel SG, Brondino MJ. Psychiatric comorbidity and the 16-month trajectory of substance-abusing and substance-dependent juvenile offenders. *J Am Acad Child Adolesc Psychiatry*. 1999;38(9):1118-1124.
- Douglas KS, Epstein ME, Poythress NG. Criminal recidivism among juvenile offenders: testing the incremental and predictive validity of three measures of psychopathic features. *Law Hum Behav*. 2008;32(5):423-438.
- Harrington RC, Kroll L, Rothwell J, McCarthy K, Bradley D, Bailey S. Psychosocial needs of boys in secure care for serious or persistent offending. *J Child Psychol Psychiatry*. 2005;46(8):859-866.
- Craig TKJ, Hodson S. Homeless youth in London, II: accommodation, employment and health outcomes at 1 year. *Psychol Med*. 2000;30(1):187-194.
- Meyer SE, Carlson GA, Youngstrom E, Ronsaville DS, Martinez PE, Gold PW, Hakak R, Radke-Yarrow M. Long-term outcomes of youth who manifested the CBCL-pediatric bipolar disorder phenotype during childhood and/or adolescence. *J Affect Disord*. 2009;113(3):227-235.
- Mason WA, Kosterman R, Hawkins JD, Herrenkohl TI, Lengua LJ, McCauley E. Predicting depression, social phobia, and violence in early adulthood from childhood behavior problems. *J Am Acad Child Adolesc Psychiatry*. 2004;43(3): 307-315.
- Cohen P, Chen H, Crawford TN, Brook JS, Gordon K. Personality disorders in early adolescence and the development of later substance use disorders in the general population. *Drug Alcohol Depend*. 2007;88(suppl 1):S71-S84.
- Fothergill KE, Ensminger ME. Childhood and adolescent antecedents of drug and alcohol problems: a longitudinal study. *Drug Alcohol Depend*. 2006; 82(1):61-76.
- Fothergill KE, Ensminger ME, Green KM, Crum RM, Robertson J, Juon H-S. The impact of early school behavior and educational achievement on adult drug use disorders: a prospective study. *Drug Alcohol Depend*. 2008;92(1-3): 191-199.
- Crum RM, Juon H-S, Green KM, Robertson J, Fothergill K, Ensminger M. Educational achievement and early school behavior as predictors of alcohol-use disorders: 35-year follow-up of the Woodlawn Study. *J Stud Alcohol*. 2006; 67(1):75-85.
- Egeland JA, Shaw JA, Endicott J, Pauls DL, Allen CR, Hostetter AM, Sussex JN. Prospective study of prodromal features for bipolarity in well Amish children. *J Am Acad Child Adolesc Psychiatry*. 2003;42(7):786-796.
- Weissman MM, Warner V, Wickramaratne P, Moreau D, Olfson M. Offspring of depressed parents: 10 years later. *Arch Gen Psychiatry*. 1997;54(10):932-940.
- Clark DB, Parker AM, Lynch KG. Psychopathology and substance-related problems during early adolescence: a survival analysis. *J Clin Child Psychol*. 1999; 28(3):333-341.
- Barnow S, Schuckit M, Smith T, Freyberger HJ. Predictors of attention problems for the period from pre-teen to early teen years. *Psychopathology*. 2006; 39(5):227-235.
- King KM, Chassin L. A prospective study of the effects of age of initiation of alcohol and drug use on young adult substance dependence. *J Stud Alcohol Drugs*. 2007;68(2):256-265.
- King KM, Chassin L. Adolescent stressors, psychopathology, and young adult substance dependence: a prospective study. *J Stud Alcohol Drugs*. 2008; 69(5):629-638.
- Kirisci L, Tarter RE, Reynolds M, Vanyukov M. Individual differences in childhood neurobehavior disinhibition predict decision to desist substance use dur-

- ing adolescence and substance use disorder in young adulthood: a prospective study. *Addict Behav.* 2006;31(4):686-696.
42. Kirisci L, Tarter R, Mezzich A, Vanyukov M. Developmental trajectory classes in substance use disorder etiology. *J Psychol Addict Behav.* 2007;21(3):287-296.
 43. Nigg JT, Wong MM, Martel MM, Jester JM, Puttler LI, Glass JM, Adams KM, Fitzgerald HE, Zucker RA. Poor response inhibition as a predictor of problem drinking and illicit drug use in adolescents at risk for alcoholism and other substance use disorders. *J Am Acad Child Adolesc Psychiatry.* 2006;45(4):468-475.
 44. Buu A, Dipiazza C, Wang J, Puttler LI, Fitzgerald HE, Zucker RA. Parent, family, and neighborhood effects on the development of child substance use and other psychopathology from preschool to the start of adulthood. *J Stud Alcohol Drugs.* 2009;70(4):489-498.
 45. Reynolds MD, Tarter R, Kirisci L, Kirillova G, Brown S, Clark DB, Gavaler J. Testosterone levels and sexual maturation predict substance use disorders in adolescent boys: a prospective study. *Biol Psychiatry.* 2007;61(11):1223-1227.
 46. Trim RS, Meehan BT, King KM, Chassin L. The relation between adolescent substance use and young adult internalizing symptoms: findings from a high-risk longitudinal sample. *Psychol Addict Behav.* 2007;21(1):97-107.
 47. Costello EJ, Angold A, Burns BJ, Stangl DK, Tweed DL, Erkanli A, Worthman CM. The Great Smoky Mountains Study of Youth: goals, design, methods, and the prevalence of DSM-III-R disorders. *Arch Gen Psychiatry.* 1996;53(12):1129-1136.
 48. Copeland WE, Shanahan L, Costello EJ, Angold A. Childhood and adolescent psychiatric disorders as predictors of young adult disorders. *Arch Gen Psychiatry.* 2009;66(7):764-772.
 49. Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen H-U, Kendler KS. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: results from the National Comorbidity Survey. *Arch Gen Psychiatry.* 1994;51(1):8-19.
 50. Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry.* 2005;62(6):617-627.
 51. Merikangas KR, He JP, Burstein M, Swanson SA, Avenevoli S, Cui L, Benjet C, Georgiades K, Swendsen J. Lifetime prevalence of mental disorders in U.S. adolescents: results from the National Comorbidity Survey Replication—Adolescent Supplement (NCS-A). *J Am Acad Child Adolesc Psychiatry.* 2010;49(10):980-989.
 52. Grant BF, Stinson FS, Dawson DA, Chou SP, Dufour MCD, Compton WC, Pickering RP, Kaplan K. Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry.* 2004;61(8):807-816.
 53. Lewinsohn PM, Rohde P, Klein DN, Seeley JR. Natural course of adolescent major depressive disorder. I: continuity into young adulthood. *J Am Acad Child Adolesc Psychiatry.* 1999;38(1):56-63.
 54. Lewinsohn PM, Rohde P, Seeley JR, Klein DN, Gotlib IH. Natural course of adolescent major depressive disorder in a community sample: predictors of recurrence in young adults. *Am J Psychiatry.* 2000;157(10):1584-1591.
 55. Loeber R, Farrington DP, Stouthamer-Loeber M, Van Kammen WB. *Antisocial Behavior and Mental Health Problems: Explanatory Factors in Childhood and Adolescence.* Mahwah, NJ: Lawrence Erlbaum Associates Publishers; 1998.
 56. Briggs-Gowan MJ, Owens PL, Schwab-Stone ME, Leventhal JM, Leaf PJ, Horwitz SM. Persistence of psychiatric disorders in pediatric settings. *J Am Acad Child Adolesc Psychiatry.* 2003;42(11):1360-1369.
 57. Costello EJ, Angold A, Keeler GP. Adolescent outcomes of childhood disorders: the consequences of severity and impairment. *J Am Acad Child Adolesc Psychiatry.* 1999;38(2):121-128.
 58. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry.* 2005;62(6):593-602.
 59. Costello EJ, Mustillo S, Erkanli A, Keeler G, Angold A. Prevalence and development of psychiatric disorders in childhood and adolescence. *Arch Gen Psychiatry.* 2003;60(8):837-844.
 60. Grant BF, Dawson DA, Stinson FS, Chou SP, Dufour MC, Pickering RP. The 12-month prevalence and trends in DSM-IV alcohol abuse and dependence: United States, 1991-1992 and 2001-2002. *Drug Alcohol Depend.* 2004;74(3):223-234.
 61. Blanco C, Okuda M, Wright C, Hasin DS, Grant BF, Liu S-M, Olfson M. Mental health of college students and their non-college-attending peers: results from the National Epidemiologic Study on Alcohol and Related Conditions. *Arch Gen Psychiatry.* 2008;65(12):1429-1437.
 62. Pardini D, White HR, Stouthamer-Loeber M. Early adolescent psychopathology as a predictor of alcohol use disorders by young adulthood. *Drug Alcohol Depend.* 2007;88(suppl 1):S38-S49.
 63. Olin TM, Klein DN, Lewinsohn PM, Rohde P, Seeley JR. Longitudinal associations between depressive and anxiety disorders: a comparison of two trait models. *Psychol Med.* 2008;38(3):353-363.
 64. Wiesner M. A longitudinal latent variable analysis of reciprocal relations between depressive symptoms and delinquency during adolescence. *J Abnorm Psychol.* 2003;112(4):633-645.
 65. Lewinsohn PM, Rohde P, Seeley JR, Baldwin CL. Gender differences in suicide attempts from adolescence to young adulthood. *J Am Acad Child Adolesc Psychiatry.* 2001;40(4):427-434.
 66. Office of Human Research Protections. *OHRP Guidance on the Involvement of Prisoners in Research.* Bethesda, MD: US Dept of Health and Human Services; 2003.
 67. Snyder HN, Sickmund M. *Juvenile Offenders and Victims: 1999 National Report.* Washington, DC: Office of Juvenile Justice and Delinquency Prevention; 1999. NCJ 178257.
 68. US Department of Health and Human Services. *Healthy People 2010.* Washington, DC: US Dept of Health and Human Services; 2000.
 69. US Department of Health and Human Services. *Healthy People 2020.* <http://www.healthypeople.gov/>. Accessed December, 2010.
 70. Smedley BD, Stith AY, Nelson AR. *Unequal Treatment: Confronting Racial and Ethnic Disparities in Health Care.* Washington, DC: Institute of Medicine of the National Academies; 2003.
 71. West HC. *Prison Inmates at Midyear 2009—Statistical Tables.* Washington, DC: US Dept of Justice, Office of Justice Programs, Bureau of Justice Statistics; 2010.
 72. US Census Bureau. Resident population by race, Hispanic origin, and single years of age: 2009: Table 20. Monthly resident population estimates by age, sex, race and Hispanic origin for the United States: April 1, 2000 to December 1, 2010. <http://www.census.gov/compendia/statab/2012/tables/12s0011.pdf>. Accessed March 18, 2011.
 73. Abram KM, Teplin LA, Charles DR, Longworth SL, McClelland GM, Dulcan MK. Posttraumatic stress disorder and trauma in youth in juvenile detention. *Arch Gen Psychiatry.* 2004;61(4):403-410.
 74. Abram KM, Teplin LA, McClelland GM, Dulcan MK. Comorbid psychiatric disorders in youth in juvenile detention. *Arch Gen Psychiatry.* 2003;60(11):1097-1108.
 75. Federal policy for the protection of human subjects: notices and rules: part 2. *Fed Regist.* 1991;56(117):28002-28032.
 76. Shaffer D, Fisher P, Dulcan MK, Davies M, Piacentini J, Schwab-Stone ME, Lahey BB, Bourdon K, Jensen PS, Bird HR, Canino G, Regier DA. The NIMH Diagnostic Interview Schedule for Children Version 2.3 (DISC-2.3): description, acceptability, prevalence rates, and performance in the MECA Study: methods for the Epidemiology of Child and Adolescent Mental Disorders Study. *J Am Acad Child Adolesc Psychiatry.* 1996;35(7):865-877.
 77. Fisher PW, Shaffer D, Piacentini JC, Lapkin J, Kafantaris V, Leonard H, Herzog DB. Sensitivity of the Diagnostic Interview Schedule for Children, 2nd edition (DISC-2.1) for specific diagnoses of children and adolescents. *J Am Acad Child Adolesc Psychiatry.* 1993;32(3):666-673.
 78. Piacentini J, Shaffer D, Fisher PW, Schwab-Stone M, Davies M, Gioia P. The Diagnostic Interview Schedule for Children—Revised Version (DISC-R), III: concurrent criterion validity. *J Am Acad Child Adolesc Psychiatry.* 1993;32(3):658-665.
 79. Schwab-Stone M, Fisher PW, Piacentini J, Shaffer D, Davies M, Briggs M. The Diagnostic Interview Schedule for Children—Revised Version (DISC-R), II: test-retest reliability. *J Am Acad Child Adolesc Psychiatry.* 1993;32(3):651-657.
 80. Schwab-Stone ME, Shaffer D, Dulcan MK, Jensen PS, Fisher P, Bird HR, Goodman SH, Lahey BB, Lichtman JH, Canino G, Rubio-Stipec M, Rae DS. Criterion validity of the NIMH Diagnostic Interview Schedule for Children Version 2.3 (DISC-2.3). *J Am Acad Child Adolesc Psychiatry.* 1996;35(7):878-888.
 81. Shaffer D, Schwab-Stone M, Fisher PW, Cohen P, Piacentini J, Davies M, Conners CK, Regier D. The Diagnostic Interview Schedule for Children—Revised Version (DISC-R), I: preparation, field testing, interrater reliability, and acceptability. *J Am Acad Child Adolesc Psychiatry.* 1993;32(3):643-650.
 82. Shaffer D, Fisher P, Lucas C. The Diagnostic Interview Schedule for Children (DISC). In: Hilsenroth MJ, Segal DL, eds. *Comprehensive Handbook of Psychological Assessment.* Vol 2. Hoboken, NJ: John Wiley & Sons; 2003:256-270.
 83. Shaffer D, Fisher P, Lucas CP, Dulcan MK, Schwab-Stone ME. NIMH Diagnostic Interview Schedule for Children Version IV (NIMH DISC-IV): description, differences from previous versions, and reliability of some common diagnoses. *J Am Acad Child Adolesc Psychiatry.* 2000;39(1):28-38.
 84. Canino G, Shrout PE, Rubio-Stipec M, Bird HR, Bravo M, Ramirez R, Chavez L, Alegria M, Bauermeister JJ, Hohmann A, Ribera J, Garcia P, Martinez-Tobias A. The DSM-IV rates of child and adolescent disorders in Puerto Rico: prevalence, correlates, service use, and the effects of impairment. *Arch Gen Psychiatry.* 2004;61(1):85-93.
 85. Robins LN, Cottler LB, Bucholz KK, Compton WM. *Diagnostic Interview Schedule for DSM-IV (DIS-IV).* St. Louis, MO: Washington University; 1995.
 86. Compton WM, Cottler LB. The Diagnostic Interview Schedule (DIS). In: Hilsenroth M, Segal DL, eds. *Comprehensive Handbook of Psychological Assessment.* Vol 2. Hoboken, NJ: John Wiley & Sons; 2004:153-162.
 87. Korn E, Graubard B. *Analysis of Health Surveys.* New York, NY: John Wiley & Sons; 1999.
 88. Cochran WG. *Sampling Techniques.* New York, NY: John Wiley & Sons; 1977.
 89. Levy PS, Lemeshow S. *Sampling of Populations: Methods and Applications.* New York, NY: John Wiley & Sons; 1999.
 90. Zeger SL, Liang KY. Longitudinal data analysis for discrete and continuous outcomes. *Biometrics.* 1986;42(1):121-130.
 91. Molenberghs G, Verbeke G. *Models for Discrete Longitudinal Data.* New York, NY: Springer; 2006.
 92. *Stata Release 11* [computer program]. College Station, TX: StataCorp LP; 2009.
 93. Moffitt TE, Caspi A, Harrington H, Milne BJ. Males on the life-course-persistent and adolescence-limited antisocial pathways: follow-up at age 26 years. *Dev Psychopathol.* 2002;14(1):179-207.
 94. Teplin LA, Abram KM, McClelland GM, Washburn JJ, Pikus AK. Detecting mental disorder in juvenile detainees: who receives services. *Am J Public Health.* 2005;95(10):1773-1780.
 95. Sampson RJ, Laub JH. Crime and deviance over the life course: the salience of adult social bonds. *Am Sociol Rev.* 1990;55(5):609-627.
 96. Kreeger DA, Matsuoda RL, Eroshva EA. Motherhood and criminal desistance

- in disadvantaged neighborhoods. *Criminology*. 2010;48(1):221-258. doi:10.1111/j.1745-9125.2010.00184.x.
97. Hope TL, Wilder EI, Watt TT. The relationships among adolescent pregnancy, pregnancy resolution, and juvenile delinquency. *Social Q*. 2003;44(4):555-576. doi:10.1111/j.1533-8525.2003.tb00525.x.
 98. Thornberry TP, Wei EH, Stouthamer-Loeber M, Van Dyke J. *Teenage Fatherhood and Delinquent Behavior*. Washington, DC: US Dept of Justice, Office of Juvenile Justice and Delinquency Prevention; 2000.
 99. NCS-R dataset. Collaborative Psychiatric Epidemiology Surveys website. <http://www.icpsr.umich.edu/CPEs>. Accessed March 15, 2011.
 100. Puzanchara C, Kang W. Easy access to juvenile court statistics: 1985-2008. <http://www.ojjdp.gov/ojstatbb/ezajcs/>. Accessed June 18, 2012.
 101. Sickmund M, Sladky TJ, Kang W. Census of juveniles in residential placement databook. <http://www.ojjdp.gov/ojstatbb/ezajcrp/>. Accessed June 18, 2012.
 102. Minton TD. *Jail Inmates at Midyear 2010—Statistical Tables*. Washington, DC: US Dept of Justice; 2011.
 103. Beckett K, Nyrop K, Pflingst L. Race, drugs, and policing: understanding disparities in drug delivery arrests. *Criminology*. 2006;44(1):105-137. doi:10.1111/j.1745-9125.2006.00044.x.
 104. Beckett K, Nyrop K, Pflingst L, Bowen M. Drug use, drug possession arrests, and the question of race: lessons from Seattle. *Soc Probl*. 2005;52(3):419-441 <http://www.plu.edu/~renfroddg/doc/srebeckett.pdf>. Accessed June 18, 2012.
 105. Moore LD, Elkavich A. Who's using and who's doing time: incarceration, the war on drugs, and public health. *Am J Public Health*. 2008;98(5):782-786.
 106. Gorvin I. *Targeting Blacks: Drug Law Enforcement and Race in the United States*. New York, NY: Human Rights Watch; 2008.
 107. Kakade M, Duarte CS, Liu X, Fuller CJ, Drucker E, Hoven CW, Fan B, Wu P. Adolescent substance use and other illegal behaviors and racial disparities in criminal justice system involvement: findings from a US national survey. *Am J Public Health*. 2012;301(7):1307-1310.
 108. National comorbidity survey. Harvard School of Medicine website. <http://www.hcp.med.harvard.edu/ncs/diagnosis.php>. Accessed March 16, 2011.
 109. NCS and NCS-R data. Harvard School of Medicine website. http://www.hcp.med.harvard.edu/ncs/ncs_data.php. Accessed March 16, 2011.
 110. Costello EJ, Copeland W, Angold A. Trends in psychopathology across the adolescent years: what changes when children become adolescents, and when adolescents become adults? *J Child Psychol Psychiatry*. 2011;52(10):1015-1025.
 111. Wittchen HU, Nelson CB, Lachner G. Prevalence of mental disorders and psychosocial impairments in adolescents and young adults. *Psychol Med*. 1998;28(1):109-126.
 112. Jaffee SR, Moffitt TE, Caspi A, Fombonne E, Poulton R, Martin J. Differences in early childhood risk factors for juvenile-onset and adult-onset depression. *Arch Gen Psychiatry*. 2002;59(3):215-222.
 113. Richardson LP, Davis R, Poulton R, McCauley E, Moffitt TE, Caspi A, Connell F. A longitudinal evaluation of adolescent depression and adult obesity. *Arch Pediatr Adolesc Med*. 2003;157(8):739-745.
 114. Moffitt TE, Harrington H, Caspi A, Kim-Cohen J, Goldberg D, Gregory AM, Poulton R. Depression and generalized anxiety disorder: cumulative and sequential comorbidity in a birth cohort followed prospectively to age 32 years. *Arch Gen Psychiatry*. 2007;64(6):651-660.
 115. Newman DL, Moffitt TE, Caspi A, Magdol L, Silva PA, Stanton WR. Psychiatric disorder in a birth cohort of young adults: prevalence, comorbidity, clinical significance, and new case incidence from ages 11 to 21. *J Consult Clin Psychol*. 1996;64(3):552-562.
 116. Kessler RC, Walters EE. Epidemiology of DSM-III-R major depression and minor depression among adolescents and young adults in the National Comorbidity Survey. *Depress Anxiety*. 1998;7(1):3-14.
 117. Herschi T, Gottfredson M. Age and the explanation of crime *Am J. Social*. 1983;89(3):552-584 <http://www.jstor.org/stable/2779005>. Accessed June 18, 2012.
 118. Moffitt TE. Adolescence-limited and life-course-persistent antisocial behavior: a developmental taxonomy. *Psychol Rev*. 1993;100(4):674-701.
 119. Arnett JJ. The developmental context of substance use in emerging adulthood. *J Drug Issues*. 2005;35(2):235-253.
 120. White HR, Jackson K. Social and psychological influences on emerging adult drinking behavior. *Alcohol Res Health*. 2004-2005;28(4):182-190 <http://pubs.niaaa.nih.gov/publications/arh284/182-190.pdf>. Accessed June 18, 2012.
 121. Berzin SC, De Marco AC. Understanding the impact of poverty on critical events in emerging adulthood. *Youth Soc*. 2010;42(2):278-300. doi:10.1177/0044118X09351909.
 122. Fazel S, Danesh J. Serious mental disorder in 23000 prisoners: a systematic review of 62 surveys. *Lancet*. 2002;359(9306):545-550.
 123. Wasserman GA, McReynolds LS, Craig SS, Keating JM, Jones SA. Psychiatric disorder, comorbidity, and suicidal behavior in juvenile justice youth. *Crim Justice Behav*. 2010;37(12):1361-1376. doi:10.1177/0093854810382751.
 124. Sickmund M. *Juveniles in Residential Placement, 1997-2008: Office of Juvenile Justice and Delinquency Prevention*. Washington, DC: US Dept of Justice; 2010.
 125. Teplin LA, Abram KM, McClelland GM. Prevalence of psychiatric disorders among incarcerated women, I: pretrial jail detainees. *Arch Gen Psychiatry*. 1996;53(6):505-512.
 126. Jordan BK, Schlenger WE, Fairbank JA, Caddell JM. Prevalence of psychiatric disorders among incarcerated women, II: convicted felons entering prison. *Arch Gen Psychiatry*. 1996;53(6):513-519.
 127. Steadman HJ, Osher FC, Robbins PC, Case B, Samuels S. Prevalence of serious mental illness among jail inmates. *Psychiatr Serv*. 2009;60(6):761-765.
 128. Puzanchara C, Adams B. *Juvenile Arrests 2009*. Washington, DC: Office of Juvenile Justice and Delinquency Prevention, US Dept of Justice; 2011. NCJ 236477.
 129. Feder A, Nestler EJ, Charney DS. Psychobiology and molecular genetics of resilience. *Nat Rev Neurosci*. 2009;10(6):446-457.
 130. Drabant EM, McRae K, Manuck SB, Hariri AR, Gross JJ. Individual differences in typical reappraisal use predict amygdala and prefrontal responses. *Biol Psychiatry*. 2009;65(5):367-373.
 131. Wager TD, Davidson ML, Hughes BL, Lindquist MA, Ochsner KN. Prefrontal-subcortical pathways mediating successful emotion regulation. *Neuron*. 2008;59(6):1037-1050.
 132. US Department of Justice. *Correctional Populations in the United States, 1998*. Washington, DC: US Dept of Justice; 2001. NCJ 192929.
 133. Shaffner L. *A Study of GIRLS LINK Collaborative, Part One: The Evaluation of the GIRLS LINK Collaborative*. Chicago, IL: Cook County Bureau of Public Safety and Judicial Coordination; 2002.
 134. Slowikowski J. *In Focus: Girls' Delinquency*. Washington, DC: US Dept of Justice, Office of Juvenile Justice and Delinquency Prevention; 2010.
 135. Greene Peters and Associates. Guiding principles for promising female programming: an inventory of best practices. <http://www.ojjdp.gov/pubs/principles/contents.html>. Accessed March 22, 2005.
 136. Sherman FT. *Pathways to Juvenile Detention Reform: Detention Reform and Girls*. Baltimore, MD: Annie E. Casey Foundation; 2005.
 137. Gordon DA, Arbuthnot J, Gustafson KE, McGreen P. Home-based behavioral-systems family therapy with disadvantaged juvenile delinquents. *Am J Fam Ther*. 1988;16(3):243-255. doi:10.1080/01926188808250729.
 138. Chamberlain P, Leve LD, Degarmo DS. Multidimensional treatment foster care for girls in the juvenile justice system: 2-year follow-up of a randomized clinical trial. *J Consult Clin Psychol*. 2007;75(1):187-193.
 139. Henggeler SW, Clingempeel WG, Brondino MJ, Pickrel SG. Four-year follow-up of multisystemic therapy with substance-abusing and substance-dependent juvenile offenders. *J Am Acad Child Adolesc Psychiatry*. 2002;41(7):868-874.
 140. Aos S, Phillips P, Barnoski B, Lieb R. *The Comparative Costs and Benefits of Programs to Reduce Crime*. Olympia: Washington State Institute for Public Policy; 2001.
 141. Mulvey EP, Schubert CA, Chung HL. Service use after court involvement in a sample of serious adolescent offenders. *Child Youth Serv Rev*. 2007;29(4):518-544.
 142. Sedlak AJ, McPherson KS. *Youth's Needs and Services: Findings from the Survey of Youth in Residential Placement*. Washington, DC: US Dept of Justice, Office of Juvenile Justice and Delinquency Prevention; 2010.
 143. Karberg JC, James DJ. *Substance Dependence, Abuse, and Treatment of Jail Inmates, 2002*. Washington, DC: US Dept of Justice; 2005.
 144. Mumola CJ, Karberg JC. *Drug Use and Dependence, State and Federal Prisoners, 2004*. Washington, DC: US Dept of Justice; 2006.
 145. Chandler RK, Fletcher BW, Volkow ND. Treating drug abuse and addiction in the criminal justice system: improving public health and safety. *JAMA*. 2009;301(2):183-190.
 146. Substance Abuse and Mental Health Services Administration. *Results From the 2009 National Survey on Drug Use and Health: volume I: Summary of National Findings*. Rockville, MD: Office of Applied Studies, Substance Abuse and Mental Health Services Administration; 2010. NSDUH series H-38A, HHS publication No. SMA 10-4856 Findings.
 147. 2009 National Survey on Drug Use and Health: detailed tables. <http://www.samhsa.gov/data/NSDUH/2k10ResultsTables/NSDUHTables2008R/HTM/TOC.htm>. Accessed October 4, 2010.
 148. Continuation of full benefits in certain cases of medical confinement. 20 CFR §416.212.
 149. Freudenberg N, Daniels J, Crum M, Perkins T, Richie BE. Coming home from jail: the social and health consequences of community reentry for women, male adolescents, and their families and communities. *Am J Public Health*. 2005;95(10):1725-1736.
 150. National Federation of Independent Business v Sebelius, 567 US 2012 (2012).
 151. Jacobson PD, Gostin LO. Restoring health to health reform. *JAMA*. 2010;304(1):85-86.
 152. Gostin LO, Jacobson PD, Record KL, Hardcastle LE. Restoring health to health reform: integrating medicine and public health to advance the population's well-being. *Univ PA Law Rev*. 2011;159(6):1777-1823.
 153. Katz MH. Future of the safety net under health reform. *JAMA*. 2010;304(6):679-680.
 154. Redlener I, Grant R. America's safety net and health care reform—what lies ahead? *N Engl J Med*. 2009;361(23):2201-2204.
 155. Congressional Budget Office. Updated estimates for the insurance coverage provisions of the Affordable Care Act. March 2012. <http://cbo.gov/sites/default/files/cbofiles/attachments/03-13-Coverage%20Estimates.pdf>. Accessed July 30, 2012.
 156. Hochman M, Katz MH. The Supreme Court decision: a victory for uninsured patients. *Arch Intern Med*. In press.
 157. McDonough JE. The road ahead for the Affordable Care Act. *N Engl J Med*. 2012;367(3):199-201.
 158. Congressional Budget Office. Estimates for the insurance coverage provisions of the Affordable Care Act updated for the recent Supreme Court decision. July 2012. <http://www.cbo.gov/sites/default/files/cbofiles/attachments/43472-07-24-2012-CoverageEstimates.pdf>. Accessed July 30, 2012.