

Sex Differences in the Prevalence and Detection of Depressive and Anxiety Disorders in General Health Care Settings

Report From the World Health Organization Collaborative Study on Psychological Problems in General Health Care

Richard Gater, MD; Michele Tansella, MD; Ailsa Korten, BSc; Bea G. Tiemens, MA; Venos G. Mavreas, MD, PhD; Michael O. Olatawura, MD

Background: Understanding the relevance of biological and social factors to sex differences in the prevalence and detection of depressive and anxiety disorders has been impaired by the lack of standardized research methods across cultures.

Method: Prevalence rates of depressive and anxiety disorders were assessed using a 2-stage design from 26 969 patients attending for primary care in 15 centers from 4 continents. Logistic regression analysis was used to examine sex differences in prevalence and detection across centers.

Results: Odds ratios for women compared with men of current depression (1.60; 95% confidence interval [CI], 1.37-1.86) and agoraphobia or panic (1.63; 95% CI, 1.18-2.20) were consistent across centers. The odds ratio for generalized anxiety varied among centers: 3 groups of centers were identified with odds ratios of 0.46 (95% CI, 0.27-0.78), 1.34 (95% CI, 1.08-1.66), and 3.09 (95% CI, 1.60-

5.89). There was no sex difference in the detection of depressive and anxiety disorders by physicians across centers.

Conclusions: The absence of a sex-by-center effect for current depression and agoraphobia or panic disorder is consistent with biological and psychosocial factors, either interacting or working alone, that have a similar final effect across cultures. It does not support the idea that sex differences in prevalence are caused by local psychosocial factors that vary from country to country. The variation in the odds ratio for generalized anxiety disorder offers some support to the idea that there are local differences between the centers contributing to the sex difference in rates. Patients' sex does not appear to affect the likelihood of current depression and anxiety being detected by primary care physicians.

Arch Gen Psychiatry. 1998;55:405-413

From the School of Psychiatry and Behavioural Sciences, University of Manchester, Manchester, England (Dr Gater); the Institute of Psychiatry, Università di Verona, Verona, Italy (Dr Tansella); the National Health and Medical Research Council Psychiatric Epidemiology Research Centre, The Australian National University, Canberra, Australia (Ms Korten); the Department of Social Psychiatry, University of Groningen, Groningen, the Netherlands (Ms Tiemens); the University Mental Health Research Institute, Eginition Hospital, Athens, Greece (Dr Mavreas); and the Department of Psychiatry, University College Hospital, Ibadan, Nigeria (Dr Olatawura).

A FEMALE preponderance of depressive disorders has been a consistent observation in community care surveys.^{1,2} The female-male ratio for current depression calculated from reported rates in the community often falls between 1.3 and 2.8.³⁻¹¹ The female-male ratio for anxiety disorders calculated from published community rates varies from 0.7 in Canberra, Australia,⁴ to 4.5 in London, England,⁵ and 4.8 in Santander, Spain.⁷ Studies in primary care settings have seldom reported prevalence by sex for individual disorders, but when this has been done, prevalence rates were higher in women than in men for all categories of depressive disorder, whereas rates of anxiety were lower in women than in men.¹² Previous studies have reported on the detection by primary care physicians of depressive and anxiety disorders in men and women.¹³⁻¹⁵

It is difficult to interpret variations between prevalence and detection rates from different studies because of differences in survey methods, particularly the methods of sample selection and diagnosis. Nevertheless, the investigation of cross-cultural variations in the sex ratio of rates of disorders can contribute to our understanding of the causative factors underlying the observed difference in rates. Biological factors, experience, and circumstances are continuously interacting and influencing each other in emotional development and expression and in the cause of mental disorders.¹⁶⁻¹⁹ Previous research has produced evidence that sex differences in the rates of mental disorders are related to the different social roles of men and women.²⁰⁻²⁶ If such factors are important and vary among cultures, they are likely to be reflected in sex ratios that vary among cultures.

METHODS

The World Health Organization Psychological Problems in General Health Care study has investigated the form, frequency, management, and outcome of common psychological disorders in primary care patients. Participating centers are in Ankara, Turkey; Athens, Greece; Bangalore, India; Berlin, Germany; Groningen, the Netherlands; Ibadan, Nigeria; Mainz, Germany; Manchester, England; Nagasaki, Japan; Paris, France; Rio de Janeiro, Brazil; Santiago, Chile; Seattle, Wash; Shanghai, China; and Verona, Italy. The choice of centers was made taking into account the need to select from a variety of countries, the availability of previous community and primary care surveys for comparison, and the capability of centers to carry out research using standardized instruments and epidemiological methods. The general health care clinics were selected to be prototypically representative of primary health care services in their country, and, wherever possible, atypical or university-based clinics were avoided. The participating clinics varied in type, organization, and the sociodemography, presenting symptoms, and physical and mental disorders of their patients (**Table 1**). Full descriptions of each of the centers and of the study methods have been published elsewhere.²⁷

SAMPLING PROCEDURE

The study used a 2-stage sampling procedure: a first-stage screening followed by a second-stage standardized in-person interview of a selected sample. In the first stage, consecutive patients, between the ages of 18 and 65 years, who were attending primary care clinics were administered the 12-item General Health Questionnaire (GHQ)²⁸ by a research worker. Respondents completed this written screening questionnaire, or, if they were not able to read or write,

the questionnaire was administered verbally following a standard protocol. For the second stage, a stratified random sample was selected for interview. Samples were drawn from 3 GHQ strata: 100% of patients with a high GHQ score, a random sample of 35% of medium scorers, and a random sample of 10% of low scorers. Based on the GHQ distribution for each sex at each center, high scorers were defined as the highest 20% at the center, medium scorers were the next 20%, and the remaining 60% were classified as low GHQ scorers.

Second-Stage Assessment

For the second-stage assessment, we used the primary care version of the Composite International Diagnostic Interview (CIDI).²⁹ The CIDI was selected because there are explicit diagnostic algorithms for *International Statistical Classification of Diseases, 10th Revision (ICD-10)*,³⁰ and *DSM-III-R*³¹ diagnoses; it has high interrater reliability; and it has been translated, backtranslated, and tested in the different languages relevant to the centers involved in the study. Several other instruments were also included in the second-stage interview to collect information on the main symptom, pathways to care, physical and psychological comorbidity, social disability, and self-rated health status.³² The assessments were carried out by interviewers who received uniform training and met a high standard of interrater reliability.²⁷

For each patient selected, the primary care physician who treated the patient completed a brief Physician's Encounter Form to indicate his or her assessment of the patient's current physical and psychological status and the management of the case. The primary care physician was asked to rate each patient's mental health status on a 5-point scale, according to whether the physician thought the patient was completely normal (1 point); had some psychological symptoms but was not mentally ill (2 points);

The World Health Organization study of Psychological Problems in General Health Care used a standardized method of recruitment and assessment across 15 different centers. It offers an ideal opportunity to investigate sex differences in the prevalence of depressive and anxiety disorders in primary care across different cultures. In this article, we will focus on sex differences in the current rates of depressive and anxiety disorders across the centers. We will address the following questions:

- Are there sex differences in the current prevalence of depressive and anxiety disorders in primary care? Are there differences in the sex ratio of these disorders between centers?
- Are there sex differences in the detection of depressive and anxiety disorders in primary care? Are there differences between centers in the detection of these disorders according to the sex of the patient?

RESULTS

In total, from the 15 centers, 26 969 primary care patients were eligible for screening, of whom 25 916 (96.1%) completed the 12-item GHQ (**Table 2**). From all cen-

ters, 62% of those screened were women, ranging from 47% in Paris to 73% in Ibadan. Second-stage interviews were completed on 5438 patients, representing 64.9% of those eligible. There was some variation between centers in the response rate for the second-stage interview. The method of weighting study data, however, was adjusted to take account of nonresponse according to sex and GHQ score stratum for each center.

CURRENT DEPRESSIVE EPISODE

The average prevalence for an *ICD-10* current depressive episode among men was 7.1% (95% confidence interval [CI], 6.0%-8.3%) compared with 12.5% (95% CI, 11.5%-13.7%) among women (**Table 3**). Most centers had higher rates in women, and this reached significance at the 5% level in Bangalore, Paris, Rio de Janeiro, and Santiago.

The results of the LR analyses, with the current depressive episode as the dependent variable for all centers combined, are shown in **Table 4**. The sex-by-center interaction was not significant (LR, $\chi^2_{14} = 20.53$; $P = .11$). Therefore, the odds of depression for women com-

or had a case of mild (3 points), moderate (4 points), or severe (5 points) psychological disorder. The physicians made their ratings using all the information normally available to them at the index consultation, but they were not aware of the results of the screening test.

Case Definition and Diagnostic Classification

Responses to the CIDI were used to make *ICD-10* diagnoses for the current depressive episode, generalized anxiety disorder (GAD), and agoraphobia or panic disorder using a computer algorithm based on the *ICD-10* Diagnostic Criteria for Research.³⁰ Prevalence rates were calculated using weights that reflected the probabilities of selection into the sample according to sex and GHQ score, adjusted for nonresponse.²⁷ The weighting procedure took account of the differences in cutoff points and response rates across centers. Therefore, the reported rates are unbiased estimates of the prevalence of each disorder among all consecutive patients attending at the respective primary care sites.

STATISTICAL ANALYSIS

For each center, prevalence rates for men and women were compared using the Fisher exact test. To test whether sex differences were consistent across centers, logistic regression (LR) analysis was used with the disorder of interest as the dependent variable, and sex, center, and sex-by-center interaction were entered as independent variables. A significant sex-by-center interaction term indicates that the centers differ in the sex ratio of the disorder. Post hoc comparisons of coefficients were used to show any centers that were outliers. The effects of possible confounders were tested one by one. Results are presented in the form of odds ratios (ORs) for women compared with men when all other variables in the model are held constant. Model 1 contains only sex and center as independent variables, so the OR presented represents the

increase in the odds of the disorder for women compared with men, adjusting for center. Model 2 includes sex, center, and GHQ stratum. All other models presented in the tables include the independent variables sex, center, GHQ stratum, and the confounding variable of interest. Confounding variables are included in the model singly because small numbers in individual cells lead to unstable estimates if all are included simultaneously. Data were unweighted for all LR analyses.

The following possible confounding variables, which vary considerably between centers and are associated with sex, were considered: comorbid disorders, GHQ stratum, age (coded 18-24, 25-44, and 45-65 years), education (<9, 9-13, and >13 years), marital status (never married; currently married; or widowed, separated, or divorced), employment status (employed, unemployed, or keeping house), and number of children (0 or 1; and ≥ 2). The samples were representative of the primary care services investigated but may not have been representative of the communities from which they came. Therefore, apparent sex differences in rates among patients attending primary care clinics could be accounted for by variations in sex differences in care-seeking behavior. To examine this possibility, the time since the last appointment was included in the LR analyses as a proxy measure of care seeking.³³

Logistic regression models were also used to investigate whether primary care physicians differ across centers in their detection rates by sex of the patients. The dependent variable was the presence of a diagnosis of depressive or anxiety disorder according to *ICD-10* criteria, as determined by the primary care physician, and independent variables were the CIDI diagnosis, sex, center, and their interaction terms. The possible effects of confounding variables were also investigated. The pattern of significant interaction terms determines how detection rates vary by sex and disorder across centers. Unless otherwise stated, α levels of .05 have been used.

pared with men could be estimated across all centers simultaneously: this was found to be 1.60 (95% CI, 1.37-1.86). When adjusted for other possible confounding variables, the ORs varied only slightly, as shown in Table 4. The interaction terms of each possible confounder with sex was tested for each model and found to be nonsignificant for all, except parity. The odds of depression among women with 2 or more children was slightly higher than for women with no children or 1 child (OR=1.57 compared with 1.41; $P=.02$). The size of the difference, although statistically significant, is slight. Although most potential confounders were significantly associated with a major depressive episode, the association was independent of the sex of the patient.

GENERALIZED ANXIETY DISORDER

The average rate of GAD for men is 5.7% (95% CI, 4.7%-6.8%) and for women is 9.2% (95% CI, 8.3%-10.3%), but there appears to be considerable variation in the rates across centers (Table 3). In most centers, the rate is higher among women than among men, and there was a significantly higher prevalence of GAD among women in

Berlin, Groningen, and Rio de Janeiro. In 3 centers—Ibadan, Nagasaki, and Seattle—there is a tendency for women to have lower rates than men. In some centers, relative to the average, there are particularly high rates in women (eg, Rio de Janeiro and Santiago), whereas in other centers, there are particularly low rates in men (eg, Groningen and Verona).

In the LR analysis with GAD as the dependent variable, the sex-by-center interaction was significant (LR, $\chi^2_{14} = 27.4$; $P=.02$). Therefore, the odds of GAD for women compared with men could not be estimated across all centers. This interaction remained significant after adjustments for the GHQ stratum, comorbid current depressive episode, and age (LR, $\chi^2_{14} = 23.7$; $P=.049$). A post hoc examination of the interaction terms showed that the centers could be split into 3 different groups. Group 1, comprising Groningen and Verona, had much higher odds of GAD in women than in men (3.09; 95% CI, 1.60-5.89). Group 2 contained most of the centers: Ankara, Athens, Bangalore, Berlin, Mainz, Manchester, Paris, Rio de Janeiro, Santiago, and Shanghai. Women were more likely than men to be diagnosed as having GAD, but the OR (1.34; 95% CI, 1.08-1.66) was smaller than for the

Table 1. Characteristics of Centers and Their Patients*

Center	Participating Sites and Facilities				Sociodemography of		
	Type	Location	Cost Coverage	Length of Consultation, min	Female, %	Mean Age, y	>4 Years of School, %
Ankara, Turkey	Primary health care unit	Semiurban	Government	12	67	34	71
Athens, Greece	Health center; 2 polyclinics	Semirural; urban	Government	10	65	44	87
Bangalore, India	Health center; 2 primary care centers	Urban; rural (2)	Government (2); private	5	50	32	56
Berlin, Germany	GP offices	Urban	Insurance	15	59	41	100
Groningen, the Netherlands	Family practices	Urban/semiurban	Insurance	10	60	38	100
Ibadan, Nigeria	Outpatient clinics of 2 general hospitals	Urban	Missionary and private	10	73	34	64
Mainz, Germany	GP offices	Urban	Insurance	12	57	37	100
Manchester, England	GP offices	Urban	Government	10	69	40	98
Nagasaki, Japan	Outpatient department of national and municipal hospitals	Urban	Insurance and government	10	51	50	100
Paris, France	GP offices and private clinics	Urban	Insurance	15	47	38	99
Rio de Janeiro, Brazil	Outpatient clinic of university hospital	Urban	Government	20	73	46	55
Santiago, Chile	Primary care clinics	Urban	Government	10	71	45	75
Seattle, Wash	Health maintenance organized clinics	Urban	Insurance	15	67	39	100
Shanghai, China	Neighborhood hospital, district hospital	Urban	Government	10	63	46	74
Verona, Italy	GP offices	Urban	Government	15	64	43	95

*GP indicates general practice; ICD-10, International Statistical Classification of Diseases, 10th Revision.

Table 2. Response Rates for Each Center*

Center	GHQ Completed, No.	GHQ Response Rate, %	Second-Stage Interview Completed, No.	Second-Stage Response Rate, %
Ankara, Turkey	1307	99.5	400	97.1
Athens, Greece	1610	92.1	196	42.7
Bangalore, India	1366	99.4	398	83.4
Berlin, Germany	2364	92.2	400	43.1
Groningen, the Netherlands	1271	96.3	340	69.0
Ibadan, Nigeria	926	100.0	269	87.6
Mainz, Germany	2460	91.0	400	36.8
Manchester, England	1523	94.0	428	71.1
Nagasaki, Japan	1555	98.8	336	73.8
Paris, France	2096	97.7	405	61.9
Rio de Janeiro, Brazil	2795	99.7	393	46.2
Santiago, Chile	1453	99.8	274	46.0
Seattle, Wash	1962	93.0	373	61.3
Shanghai, China	1673	99.5	576	98.8
Verona, Italy	1555	95.7	250	54.7
Totals	25 916	96.5	5438	64.9

*GHQ indicates General Health Questionnaire.

centers in group 1. Group 3 consisted of the centers in Ibadan, Seattle, and Nagasaki, where women were significantly less likely than men to be diagnosed as having GAD (OR, 0.46; 95% CI, 0.27-0.78).

When adjusted for other possible confounding variables, the ORs varied only slightly (**Table 5**). The interaction terms of each possible confounder with sex were tested and found to be nonsignificant except for marital

status, which presented a complex picture. To interpret the coefficients, separate analyses were done for the group of patients who were currently married and a second group of patients who were single, widowed, separated, or divorced ("not currently married"). Odds ratios for women being diagnosed as having GAD compared with men were obtained. For the group 1 centers, the OR of GAD for married women compared with married men is 2.4 (95%

Study Sample		Presenting Symptom		Physical Comorbidity		Prevalence of Mental Disorder		
Married, %	Not Employed, %	Psychological, %	Physical or Pain, %	Any Physical Illness, %	Physical Illness, at Least Moderate Severity, %	ICD-10 Cases, %	Current Depression, %	Generalized Anxiety, %
74	6	3	85	66	24	18	12	1
61	19	2	55	41	8	22	6	15
61	5	1	53	48	16	24	9	9
54	20	4	68	67	27	25	6	9
63	14	13	65	14	3	29	16	6
78	4	2	82	48	13	10	4	3
53	6	4	79	75	31	31	11	8
50	31	9	67	40	6	26	17	7
81	14	1	44	55	19	15	3	5
44	8	11	65	52	10	31	14	12
56	18	8	59	57	19	38	16	23
47	19	13	50	72	18	54	30	19
48	7	3	38	38	9	20	6	2
86	17	0.2	79	41	9	10	4	2
71	15	6	52	62	21	12	5	4

CI, 1.1-5.3), whereas among those not currently married, the OR for women compared with men is higher at 5.2 (95% CI, 1.5-18.8). For the group 2 centers, the OR of GAD for women compared with men was significant only for those who were currently married (OR, 1.65; 95% CI, 1.22-2.24). Among those not currently married, the OR for women compared with men was 0.87 (95% CI, 0.62-1.21). Finally, for the group 3 centers, there was no sex-by-marital status interaction and no differences in the GAD according to marital status.

AGORAPHOBIA AND PANIC DISORDER

The rates for agoraphobia and panic disorder were lower than those for the other diagnoses, and these 2 disorders were combined and analyzed together. The overall average prevalence of agoraphobia or panic disorder was 1.5% (95% CI, 1.0%-2.1%) in men and 2.8% (95% CI, 2.2%-3.4%) in women (Table 3). In only 1 center (Paris) were the rates significantly higher in women. Elsewhere the comparison of rates within each center did not reveal any significant differences in the rates in men compared with women, although the small number involved limits the power of these comparisons.

The results of the LR analyses, with agoraphobia or panic disorder (or both) as the dependent variable, are shown in **Table 6**. The sex-by-center interaction was not significant (LR, $\chi^2_{14} = 14.54$; $P = .41$), and the odds of agoraphobia and panic disorder for women compared with men could be estimated across all centers si-

multaneously. This was found to be 1.63 (95% CI, 1.18-2.20). When adjusted for other possible confounding variables, the OR varied only slightly. The interaction terms of each possible confounder with sex was tested and found to be nonsignificant.

DETECTION OF CIDI CASES BY PRIMARY CARE PHYSICIANS

When each center was analyzed separately, in none were there significant differences in the detection rates of ICD-10 depressive and anxiety disorders for male patients compared with female patients. A series of LR analyses were made with detection by the primary care physician as the dependent variable and sex, center, CIDI diagnosis, and GHQ stratum and their interaction terms as independent variables. Because there were significant interaction terms between the diagnostic variables, sex and center, the analysis was conducted for 2 groups separately: 1 group of all subjects given a CIDI diagnosis of 1 or more of the depressive and anxiety disorders and a second group of those who received none of these diagnoses. For these separate LR analyses, interaction terms involving a center were not significant, and therefore an analysis for all centers combined could be made. Among subjects with a CIDI diagnosis, there is no difference between men and women in the odds of being given a psychological diagnosis by the primary care physician, controlling for center and GHQ stratum (OR for detection for women compared with men, 0.94; 95% CI, 0.79-1.21). Among those subjects who did not re-

Table 3. Prevalence by Sex and Sex Ratios of Composite International Diagnostic Interview–Derived ICD-10 Depressive and Anxiety Disorders for Each Center*

Center	Current Depressive Episode				Generalized
	No. †	Prevalence, %‡		Sex Ratio	
		Women	Men		
Ankara, Turkey	105	12.5	9.8	1.3	7
Athens, Greece	34	6.4	6.5	1.0	44
Bangalore, India	72	13.3	4.8	2.8§	66
Berlin, Germany	62	7.7	3.7	2.1	60
Groningen, the Netherlands	121	17.9	13.0	1.4	47
Ibadan, Nigeria	23	3.8	5.3	0.7	15
Mainz, Germany	62	12.3	9.8	1.3	46
Manchester, England	154	18.3	13.9	1.3	62
Nagasaki, Japan	25	2.8	2.3	1.2	30
Paris, France	132	18.7	9.3	2.0§	79
Rio de Janeiro, Brazil	119	19.6	5.8	3.4§	112
Santiago, Chile	117	36.8	11.2	3.3§	73
Seattle, Wash	61	6.5	6.0	1.1	18
Shanghai, China	53	4.4	3.3	1.3	25
Verona, Italy	34	5.5	3.2	1.7	21
Totals	1174	12.5	7.1	1.8§	705

*ICD-10 indicates International Statistical Classification of Diseases, 10th Revision; question mark, unknown.

†Number of patients with the disorder interviewed.

‡Estimated percentage of consecutive patients attending the center who have the disorder.

§P ≤ .01.

||P ≤ .05.

Table 4. Results of the Logistic Regression Analyses With the Current Depressive Episode as the Dependent Variable for All Centers*

Model†	Odds Ratio (95% CI) of Current Depressive Episode for Women Compared With Men
Sex + center	1.60 (1.37-1.86)
Model 1 + GHQ stratum	1.52 (1.29-1.75)
Model 2 + GAD	1.52 (1.29-1.52)
Model 2 + age	1.52 (1.30-1.78)
Model 2 + marital status‡	1.48 (1.26-1.74)
Model 2 + parity‡	§
Model 2 + employment‡	1.51 (1.28-1.79)
Model 2 + education	1.49 (1.28-1.75)
Model 2 + time since last visit	1.51 (1.28-1.76)

*CI indicates confidence interval; GHQ, General Health Questionnaire; and GAD, generalized anxiety disorder.

†Variables included in the model as independent terms. Interaction terms of each confounding variable with sex were tested and found to be nonsignificant, except for parity by sex.

‡Main effect nonsignificant.

§The odds ratio (95% CI) for women with 0 to 1 child compared with men is 1.41 (1.19-1.67) and for women with ≥ 2 children compared with men is 1.57 (1.29-1.91).

ceive a CIDI diagnosis, however, the odds of a woman being reported by the physician as having a psychological diagnosis are slightly higher than that for a man (OR, 1.20; 95% CI, 1.01-1.43). When the estimates of the odds ratios are adjusted for confounding variables (age, education, marital status, parity, employment, and frequency of consultation), the individual ORs vary only slightly. Confounding variables were related to the odds of detection, but no interaction terms between these and sex were found. That is, the detection of a case of a psychological disorder asso-

ciated with sex does not alter according to whether confounding variables were present.

COMMENT

The odds of a current depressive episode for women compared with men across all centers were estimated to be 1.60, and for agoraphobia and panic disorder, the OR was 1.63. The absence of a sex-by-center effect suggests that the causes of a higher rate in women for these disorders are having a similar resultant effect in all centers. This does not support the idea that the sex difference in rates is caused by local psychosocial effects that vary from one society to another. It is compatible with biological or psychosocial factors, either interacting or working alone, that have a similar final effect across cultures.

In contrast, the odds for men and women of GAD vary between centers. In some centers, Groningen and Verona (group 1), the odds of having generalized anxiety are 3 times higher for women than for men. In most centers (group 2), there is a modest OR of 1.3, and in a few centers (group 3), the odds for women are half those for men. These findings offer support to the idea that there are differences between the centers that contribute to the sex difference in rates of GAD, such as differences in the social roles and experiences of women and men. Our findings also suggest different effects of marital status for men and women when these 3 groups of centers are analyzed separately. An important reason for carrying out this post hoc separation of the centers into 3 groups is to try to present the results in a manner that allows a nonstatistical evaluation. The grouping of centers, however, does not fall into a pattern that is culturally recognizable, and the health care centers do not have characteristics that

Anxiety Disorder				Agoraphobia or Panic Disorder			
Prevalence, %‡			Sex Ratio	No.†	Prevalence, %‡		
Women	Men				Women	Men	Sex Ratio
1.1	0.5		2.2	7	1.7	0.5	3.4
16.1	12.5		1.3	7	1.6	1.5	1.1
10.8	6.2		1.7	9	1.0	1.1	0.9
11.9	5.0		2.4	13	3.5	0.9	3.9
9.1	2.5		3.6	30	5.0	2.0	2.5
2.2	4.9		0.4	5	0.5	1.5	0.3
10.0	5.1		2.0	20	3.4	2.1	1.6
8.1	4.9		1.7	48	5.6	4.0	1.4
3.6	6.5		0.6	2	0.2	0.2	1.0
14.1	9.9		1.4	20	5.6	1.5	3.7
25.8	14.1		1.8	23	2.1	4.1	0.5
21.8	11.0		2.0	25	5.5	1.6	3.4
1.4	3.7		0.4	16	3.0	2.8	1.1
2.1	1.7		1.2	3	0.3	0.0	?
4.9	1.5		3.3	8	2.9	0.4	7.3
9.2	5.7		1.6§	236	2.8	1.5	1.9§

Table 5. Results of the Logistic Regression Analyses With Generalized Anxiety Disorder (GAD) as a Dependent Variable for All Centers*

Model†	Odds Ratio (95% CI) of GAD for Women Compared With Men			
	Groningen and Verona	Ankara, Athens, Bangalore, Berlin, Mainz, Manchester, Paris, Rio, Santiago, and Shanghai		Ibadan, Nagasaki, and Seattle
Sex + center	3.09 (1.60-5.89)	1.34 (1.08-1.66)	0.46 (0.27-0.78)	
Model 1 + GHQ stratum	2.91 (1.48-5.71)	1.27 (1.03-1.57)	0.48 (0.28-0.82)	
Model 2 + depressive episode	2.73 (1.39-5.38)	1.16 (0.93-1.44)	0.48 (0.28-0.83)	
Model 2 + age	3.01 (1.55-5.87)	0.49 (0.27-0.78)	1.26 (1.01-1.56)	
Model 2 + marital status	‡	§	0.47 (0.27-0.67)	
Model 2 + parity	3.01 (1.55-5.87)	1.35 (1.09-1.69)	0.49 (0.29-0.84)	
Model 2 + employment	2.88 (1.25-6.51)	1.29 (1.01-1.62)	0.53 (0.30-0.96)	
Model 2 + education	2.81 (1.44-5.49)	1.25 (1.01-1.55)	0.50 (0.29-0.84)	
Model 2 + time since last visit	2.74 (1.39-5.39)	1.27 (1.02-1.57)	0.48 (0.28-0.82)	

*CI indicates confidence interval; GHQ, General Health Questionnaire. See Table 1 for country names for cities.

†Variables are included in the model as independent terms. Interaction terms of each confounding variable with sex was tested and found to be nonsignificant, except for marital status by sex.

‡Marital status-by-sex interaction is significant. For those currently married, the odds ratio for women compared with men is 2.40 (95% CI, 1.06-5.26). For those not currently married, the odds ratio for women compared with men is 5.24 (95% CI, 1.46-18.77).

§Marital status-by-sex interaction is significant. For those currently married, the odds ratio for women compared with men is 1.65 (95% CI, 1.22-2.24). For those not currently married, the odds ratio for women compared with men is 0.87 (95% CI, 0.62-1.21).

fall into these 3 groups. Although the status of GAD as an independent diagnosis has been questioned,³⁴⁻³⁶ there is no reason to suppose that in the different centers this disorder has been assessed less consistently than the other disorders.

There are no intercenter differences in the odds of detection for women compared with men. In fact, the detection of ICD-10 cases of depressive and anxiety disorders is similar for men and women in every center. Therefore, sex does not appear to affect the likelihood of

depressive and anxiety disorders being detected by primary care physicians.

The Psychological Problems in General Health Care study provides an excellent opportunity to look for differences across centers because the same method has been used in 15 different centers around the world. The analysis is complex, however, and it is important not to read too much into the data. This study recruited from primary care practices, which may not reflect the situation in the community. Because the time since the last ap-

Table 6. Results of the Logistic Regression Analyses With Agoraphobia or Panic Disorder as the Dependent Variable for All Centers*

Model†	Odds Ratio (95% CI) of Current Agoraphobia or Panic Disorder for Women Compared With Men
Sex + center	1.63 (1.18-2.20)
Model 1 + GHQ stratum	1.53 (1.12-2.12)
Model 2 + depressive episode	1.41 (1.04-2.04)
Model 2 + age	1.55 (1.12-2.15)
Model 2 + marital status	1.54 (1.09-2.10)
Model 2 + parity	1.54 (1.10-2.15)
Model 2 + employment	1.52 (1.07-2.16)
Model 2 + education	1.54 (1.11-2.13)
Model 2 + time since last visit	1.55 (1.12-2.15)

*CI indicates confidence interval; GHQ, General Health Questionnaire.

†Variables included in the model as independent terms. Interaction terms of each confounding variable with gender were tested and found to be nonsignificant.

pointment did not make a significant contribution to the analysis, however, we conclude that sex differences in the frequency of care seeking did not account for our findings. The small number of patients identified in each center means that the estimates of prevalence may be unstable. This is apparent in the large CIs of the ORs indicating sex differences in prevalence. Our analysis has assumed that the CIDI and ICD-10 criteria were identifying cases of mental disorder with similar accuracy in men and women in different countries, but men and women could respond differently to the interview in some centers compared with others. Previous research has suggested that varying the number of symptoms required for a diagnosis of depression alters the ratio of women to men.^{37,38} Thus, our findings may to some extent be the product of sex differences in the conceptualization and reporting of symptoms of depression and anxiety, rather than real differences in the rates of the disorders. The analysis also assumes that nonresponse was ignorable insofar as it was unrelated to the variables of interest. Finally, the finding that the ORs varied only slightly when adjusted for sociodemographic variables may simply reflect the limits of our sociodemographic variables.

Future cross-cultural research assessing sex differences in rates of disorder should consider a wider range of demographic and social variables along with other variables that have been identified elsewhere as relevant in the cause of depression and anxiety, such as genetic factors, childhood and adult adverse experiences, and the pattern of social support. Consideration should also be given to the longitudinal investigation of sex differences in the attribution of illness onset to social sources and the symbolic linking of symptoms to life context in various cultures.³⁹

Accepted for publication October 31, 1997.

This study is based on data and experience obtained during the World Health Organization (WHO) Project on Psychological Problems in General Health Care, a project sponsored by the WHO, by the participating field research

centers, and by an educational grant from Delagrangre Laboratories and Synthelabo Pharmacie, Paris, France.

Investigators from the participating centers are: Orhan M. Ozturk, MD, and Murat S. Rezaki, MD, Ankara, Turkey; Costas Stefanis, MD, and Venetsanos Mavreas, MD, PhD, Athens, Greece; S. M. Channabasavanna, MD, and T. Sriram, MD, Bangalore, India; Hanfried Helmchen, MD, and Michael Linden, PhD, MD, Berlin, Germany; Wim van den Brink, MD, PhD, and Bea Tiemens, MA, Groningen, the Netherlands; Michael O. Olatawura, MD, and Oye Gureje, MD, Ibadan, Nigeria; Otto Benkert, MD, and Wolfgang Maier, MD, Mainz, Germany; Richard Gater, MD, and Steven Kisely, MSc, Manchester, England; Yoshiyumi Nakane, MD, and Sunichirou Michitsuji, MD, Nagasaki, Japan; Yves Lecrubier, MD, and Patrice Boyer, MD, Paris, France; Jorge-Alberto Costa e Silva, MD, and Louis Villano, MD, MSc, Rio de Janeiro, Brazil; Ramon Florenzano Urzua, MD, and Julia Acuno Rojas, MD, Santiago, Chile; Gregory E. Simon, MD, and Michael Von Korff, DSc, Seattle, Wash; Yan He-Qin, MD, and Zaio Shi Fu, MD, Shanghai, People's Republic of China; and Michele Tansella, MD, and Cesario Bellantuono, MD, Verona, Italy. The Advisory Group consisted of the following members: Dr Silva; David P. Goldberg, MD, Manchester; Dr Lecrubier; Johan Ormel, PhD, Groningen; Dr Korff; Hans-Uli Wittchen, PhD, Munich, Germany; T. Bedhiran Üstün, MD, PhD, WHO Headquarters, Geneva, Switzerland; and Norman Sartorius (chair), MD, PhD, WHO Headquarters.

Reprints: Richard Gater, MD, School of Psychiatry and Behavioural Sciences, Rawnsley Building, Manchester Royal Infirmary, Oxford Road, Manchester M13-9WL, England.

REFERENCES

- Weissman MM, Klerman GL. Sex differences and the epidemiology of depression. *Arch Gen Psychiatry*. 1977;34:98-111.
- Piccinelli M, Gomez-Homen F. *Gender Differences in the Epidemiology of Affective Disorders and Schizophrenia*. Geneva, Switzerland: World Health Organization; 1997.
- Orley J, Wing JK. Psychiatric disorders in two African villages. *Arch Gen Psychiatry*. 1979;36:513-520.
- Henderson S, Duncan-Jones P, Byrne DG, Scott R, Adcock S. Psychiatric disorder in Canberra: a standardised study of prevalence. *Acta Psychiatr Scand*. 1979; 60:355-374.
- Bebbington P, Hurry J, Tennant C, Sturt E, Wing JK. Epidemiology of mental disorders in Camberwell. *Psychol Med*. 1981;11:561-579.
- Mavreas VG, Beis A, Mouyias A, Rignon F, Lyketsos GC. Prevalence of psychiatric disorders in Athens. *Soc Psychiatry*. 1986;21:172-181.
- Vásquez-Barquero JL, Díez-Manrique JF, Peña C, Aldama J, Samaniego Rodríguez C, Menéndez Arango J, Mirapeix C. A community health survey in Cantabria: a general description of morbidity. *Psychol Med*. 1986;17:227-241.
- Wells JE, Bushnell JA, Hornblow AR, Joyce PR, Oakley-Browne MA. Christchurch psychiatric epidemiology study, I: methodology and lifetime prevalence for specific psychiatric disorders. *Aust N Z J Psychiatry*. 1989;23:315-326.
- Regier DA, Narrow WE, Rae DS. The epidemiology of anxiety disorders: the Epidemiological Catchment Area (ECA) experience. *J Psychiatr Res*. 1990;24 (suppl 2):3-14.
- Lehtinen V, Joukamaa M, Lahtela K, Raitasalo R, Jyrkinen E, Maatela J, Aromaa A. Prevalence of mental disorders among adults in Finland: basic results from the Mini Finland Health Survey. *Acta Psychiatr Scand*. 1990;81:418-425.
- Lee CK, Kwak YS, Yamamoto J, Rhee H, Kim YS, Han JH, Choi JO, Lee YO. Psychiatric epidemiology in Korea, I: gender and age differences in Seoul. *J Nerv Ment Dis*. 1990;178:242-251.
- Barrett JE, Barrett JA, Oxman TE, Gerber P. The prevalence of psychiatric disorders in a primary care practice. *Arch Gen Psychiatry*. 1988;45:1100-1106.
- Goldberg D, Huxley P. *Common Mental Disorders: A Bio-Social Model*. London, England: Routledge; 1992.

14. Marks JN, Goldberg DP, Hillier VF. Determinants of the ability of general practitioners to detect psychiatric illness. *Psychol Med.* 1979;9:337-353.
15. Boardman AP. The General Health Questionnaire and the detection of emotional disorder by general practitioners. *Br J Psychiatry.* 1987;150:373-381.
16. Kendler KS. Genetic epidemiology in psychiatry. *Arch Gen Psychiatry.* 1995;52:895-899.
17. Kendler KS. Adversity, stress, and psychopathology: a psychiatric genetic perspective. *Int J Methods Psychiatr Res.* 1995;5:163-170.
18. Schore AN. *Affect Regulation and the Origin of the Self: The Neurobiology of Emotional Development.* Hillsdale, NJ: Lawrence A Erlbaum Associates; 1994.
19. Eisenberg L. The social construction of the human brain. *Am J Psychiatry.* 1995;152:1563-1575.
20. Rosenfield S. Sex differences in depression: do women always have higher rates? *J Health Soc Behav.* 1980;21:33-42.
21. Anenshel CS, Frerichs RR, Clark VA. Family roles and sex differences in depression. *J Health Soc Behav.* 1981;22:379-393.
22. Gore S, Mangione TW. Social roles, sex roles, and psychological distress: additive and interactive models of sex differences. *J Health Soc Behav.* 1983;24:300-312.
23. Cleary PD, Mechanic D. Sex differences in psychological distress among married people. *J Health Soc Behav.* 1983;24:111-121.
24. Lennon MC. Sex differences in distress: the impact of gender and work roles. *J Health Soc Behav.* 1987;28:290-305.
25. Ruble DN, Greulich F, Pomerantz EM, Gochberg B. The role of gender-related processes in the development of sex differences in self-evaluation and depression. *J Affect Dis.* 1993;29:97-128.
26. Glass J, Fujimoto T. Housework, paid work, and depression among husbands and wives. *J Health Soc Behav.* 1994;35:179-191.
27. Von Korff M, Üstün TB. Methods of the WHO Collaborative Study on "Psychological Problems in General Health Care." In: Üstün TB, Sartorius N, eds. *Mental Illness in General Health Care: An International Study.* New York, NY: John Wiley & Sons Inc; 1995:19-38.
28. Goldberg D, Williams P. *A User's Guide to the General Health Questionnaire.* Windsor, England: NFER-NELSON Publishing Co Ltd; 1988.
29. Wittchen HU, Robins LN, Cottler LB, Sartorius N, Burke JD, Regier DA. Participants in the Multicenter World Health Organization/Alcohol, Drug Abuse, and Mental Health Administration field trials: cross-cultural feasibility, reliability, and sources of variance of the Composite International Diagnostic Interview (CIDI). *Br J Psychiatry.* 1991;159:645-653.
30. World Health Organization. *International Classification of Diseases and Related Health Problems. 10th Revision (ICD-10): Diagnostic Criteria for Research.* Geneva, Switzerland: World Health Organization; 1993.
31. American Psychiatric Association, Committee on Nomenclature and Statistics. *Diagnostic and Statistical Manual of Mental Disorders, Third Edition, Revised.* Washington, DC: American Psychiatric Association; 1987.
32. Sartorius N, Üstün TB, Costa e Silva JA, Goldberg D, Lecrubier Y, Ormel J, Von Korff M, Wittchen HU. An international study of psychological problems in primary care: preliminary report from the World Health Organization collaborative project on "Psychological Problems in General Health Care." *Arch Gen Psychiatry.* 1993;50:819-824.
33. Laska EM, Meisner M, Siegel C. Estimating the size of a population from a single sample. *Biometrics.* 1988;44:461-472.
34. Sanderson WC, Wetzler S. Chronic anxiety and generalized anxiety disorder: issues in comorbidity. In: Rapee RM, Barlow DH, eds. *Chronic Anxiety: Generalized Anxiety Disorder and Mixed Anxiety-Depression.* New York, NY: Guilford Press; 1991:119-135.
35. Andrews G. Panic and generalized anxiety disorders. *Curr Opin Psychiatry.* 1993;6:191-194.
36. Starcevic V, Uhlenhuth EH, Fallon S, Pathak D. Personality dimensions in panic disorder and generalized anxiety disorder. *J Affect Disord.* 1996;37:75-79.
37. Ernst C, Angst J. The Zurich Study, XII: sex differences in depression: evidence from longitudinal epidemiological data. *Eur Arch Psychiatry Clin Neurosci.* 1992;241:222-230.
38. Kessler RC, McGonagle KA, Swartz M, Blazer DG, Nelson CB. Sex and depression in the National Comorbidity Survey, I: lifetime prevalence, chronicity and recurrence. *J Affect Disord.* 1993;29:85-96.
39. Ware NC, Kleinman A. Culture and somatic experience: the social course of illness in neurasthenia and chronic fatigue syndrome. *Psychosom Med.* 1992;54:546-560.

Correction

Error in Table Data. In the article titled "First-Episode Schizophrenic Psychosis Differs From First-Episode Affective Psychosis and Controls in P300 Amplitude Over Left Temporal Lobe," published in the February 1998 issue of the ARCHIVES (*Arch Gen Psychiatry.* 1998;55:173-180), the reported value for the P (1-way ANOVA) for the age characteristic given in Table 2 on page 175 is incorrect. The correct P value for the age characteristic is .066.